

EFB 1369
 FAS1574C1

Drug Facts (continued)

Warnings: Do not use if you are allergic to omeprazole. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have: had heartburn over 3 months. This may be a sign of a more serious condition; heartburn with lightheadedness, sweating or dizziness; chest pain or tender pain with soreness of breast, swelling, or spreading to arms, neck or shoulders, or lightheadedness; frequent weight loss; unexplained weight loss; frequent wheezing, particularly with heartburn; frequent chest pain.

Ask a doctor or pharmacist before use if you are taking: prescription anti-fungal or anti-yeast medicines; warfarin (blood-thinning medicine); diazepam (anxiety medicine); digoxin (heart medicine); tacrolimus (immune system medicine); atazanavir (medication for HIV infection).

Stop use and ask a doctor if: your heartburn continues or worsens; you need to take this product for more than 14 days; you need to take more than 1 course of treatment every 4 months.

Other Information: Keep the carton and package insert. They contain important information. Store at 20-25°C (68-77°F). Keep product out of high heat and humidity. Protect product from moisture.

Directions: Adults 18 years of age and older (every 24 hours): This product is to be used once a day (every 24 hours). It may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours. Take every day for 14 days. Swallow 1 tablet with a glass of water before eating in the morning. Do not take more than 1 tablet a day. Do not chew or crush the tablets. Do not take for more than 14 days unless directed by your doctor. Do not use for more than 14 days unless directed by a doctor.

Repeat 14-Day Course (if needed): You may repeat a 14-day course every 4 months every 4 months unless directed by a doctor.

Children under 18 years of age: Ask a doctor.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Drug Facts (continued)

Use: Treats frequent heartburn (occurs 2 or more days a week). Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.

Warnings: Do not use if you are allergic to omeprazole. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have: had heartburn over 3 months. This may be a sign of a more serious condition; heartburn with lightheadedness, sweating or dizziness; chest pain or tender pain with soreness of breast, swelling, or spreading to arms, neck or shoulders, or lightheadedness; frequent weight loss; unexplained weight loss; frequent wheezing, particularly with heartburn; frequent chest pain.

Ask a doctor or pharmacist before use if you are taking: prescription anti-fungal or anti-yeast medicines; warfarin (blood-thinning medicine); diazepam (anxiety medicine); digoxin (heart medicine); tacrolimus (immune system medicine); atazanavir (medication for HIV infection).

Stop use and ask a doctor if: your heartburn continues or worsens; you need to take this product for more than 14 days; you need to take more than 1 course of treatment every 4 months.

Other Information: Keep the carton and package insert. They contain important information. Store at 20-25°C (68-77°F). Keep product out of high heat and humidity. Protect product from moisture.

Directions: Adults 18 years of age and older (every 24 hours): This product is to be used once a day (every 24 hours). It may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours. Take every day for 14 days. Swallow 1 tablet with a glass of water before eating in the morning. Do not take more than 1 tablet a day. Do not chew or crush the tablets. Do not take for more than 14 days unless directed by your doctor. Do not use for more than 14 days unless directed by a doctor.

Repeat 14-Day Course (if needed): You may repeat a 14-day course every 4 months every 4 months unless directed by a doctor.

Children under 18 years of age: Ask a doctor.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Drug Facts (continued)

Warnings: Do not use if you are allergic to omeprazole. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have: had heartburn over 3 months. This may be a sign of a more serious condition; heartburn with lightheadedness, sweating or dizziness; chest pain or tender pain with soreness of breast, swelling, or spreading to arms, neck or shoulders, or lightheadedness; frequent weight loss; unexplained weight loss; frequent wheezing, particularly with heartburn; frequent chest pain.

Ask a doctor or pharmacist before use if you are taking: prescription anti-fungal or anti-yeast medicines; warfarin (blood-thinning medicine); diazepam (anxiety medicine); digoxin (heart medicine); tacrolimus (immune system medicine); atazanavir (medication for HIV infection).

Stop use and ask a doctor if: your heartburn continues or worsens; you need to take this product for more than 14 days; you need to take more than 1 course of treatment every 4 months.

Other Information: Keep the carton and package insert. They contain important information. Store at 20-25°C (68-77°F). Keep product out of high heat and humidity. Protect product from moisture.

Directions: Adults 18 years of age and older (every 24 hours): This product is to be used once a day (every 24 hours). It may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours. Take every day for 14 days. Swallow 1 tablet with a glass of water before eating in the morning. Do not take more than 1 tablet a day. Do not chew or crush the tablets. Do not take for more than 14 days unless directed by your doctor. Do not use for more than 14 days unless directed by a doctor.

Repeat 14-Day Course (if needed): You may repeat a 14-day course every 4 months every 4 months unless directed by a doctor.

Children under 18 years of age: Ask a doctor.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Drug Facts (continued)

Use: Treats frequent heartburn (occurs 2 or more days a week). Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.

Warnings: Do not use if you are allergic to omeprazole. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have: had heartburn over 3 months. This may be a sign of a more serious condition; heartburn with lightheadedness, sweating or dizziness; chest pain or tender pain with soreness of breast, swelling, or spreading to arms, neck or shoulders, or lightheadedness; frequent weight loss; unexplained weight loss; frequent wheezing, particularly with heartburn; frequent chest pain.

Ask a doctor or pharmacist before use if you are taking: prescription anti-fungal or anti-yeast medicines; warfarin (blood-thinning medicine); diazepam (anxiety medicine); digoxin (heart medicine); tacrolimus (immune system medicine); atazanavir (medication for HIV infection).

Stop use and ask a doctor if: your heartburn continues or worsens; you need to take this product for more than 14 days; you need to take more than 1 course of treatment every 4 months.

Other Information: Keep the carton and package insert. They contain important information. Store at 20-25°C (68-77°F). Keep product out of high heat and humidity. Protect product from moisture.

Directions: Adults 18 years of age and older (every 24 hours): This product is to be used once a day (every 24 hours). It may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours. Take every day for 14 days. Swallow 1 tablet with a glass of water before eating in the morning. Do not take more than 1 tablet a day. Do not chew or crush the tablets. Do not take for more than 14 days unless directed by your doctor. Do not use for more than 14 days unless directed by a doctor.

Repeat 14-Day Course (if needed): You may repeat a 14-day course every 4 months every 4 months unless directed by a doctor.

Children under 18 years of age: Ask a doctor.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)



Omeprazole Delayed Release Tablets 20mg

Safety Feature - Do not use if printed tablet blister unit is open or torn.



Treats Frequent Heartburn!

Omeprazole Delayed Release Tablets 20mg

Acid Reducer

14 Tablets
 One 14-day course of treatment

CODE AREA

Manufactured by: **TEVA** LTD., Southern Industrial Zone Or Akiva 30600, Israel

Made in Israel

Drug Facts (continued)

Warnings: Do not use if you are allergic to omeprazole. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have: had heartburn over 3 months. This may be a sign of a more serious condition; heartburn with lightheadedness, sweating or dizziness; chest pain or tender pain with soreness of breast, swelling, or spreading to arms, neck or shoulders, or lightheadedness; frequent weight loss; unexplained weight loss; frequent wheezing, particularly with heartburn; frequent chest pain.

Ask a doctor or pharmacist before use if you are taking: prescription anti-fungal or anti-yeast medicines; warfarin (blood-thinning medicine); diazepam (anxiety medicine); digoxin (heart medicine); tacrolimus (immune system medicine); atazanavir (medication for HIV infection).

Stop use and ask a doctor if: your heartburn continues or worsens; you need to take this product for more than 14 days; you need to take more than 1 course of treatment every 4 months.

Other Information: Keep the carton and package insert. They contain important information. Store at 20-25°C (68-77°F). Keep product out of high heat and humidity. Protect product from moisture.

Directions: Adults 18 years of age and older (every 24 hours): This product is to be used once a day (every 24 hours). It may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours. Take every day for 14 days. Swallow 1 tablet with a glass of water before eating in the morning. Do not take more than 1 tablet a day. Do not chew or crush the tablets. Do not take for more than 14 days unless directed by your doctor. Do not use for more than 14 days unless directed by a doctor.

Repeat 14-Day Course (if needed): You may repeat a 14-day course every 4 months every 4 months unless directed by a doctor.

Children under 18 years of age: Ask a doctor.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Drug Facts (continued)

Warnings: Do not use if you are allergic to omeprazole. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have: had heartburn over 3 months. This may be a sign of a more serious condition; heartburn with lightheadedness, sweating or dizziness; chest pain or tender pain with soreness of breast, swelling, or spreading to arms, neck or shoulders, or lightheadedness; frequent weight loss; unexplained weight loss; frequent wheezing, particularly with heartburn; frequent chest pain.

Ask a doctor or pharmacist before use if you are taking: prescription anti-fungal or anti-yeast medicines; warfarin (blood-thinning medicine); diazepam (anxiety medicine); digoxin (heart medicine); tacrolimus (immune system medicine); atazanavir (medication for HIV infection).

Stop use and ask a doctor if: your heartburn continues or worsens; you need to take this product for more than 14 days; you need to take more than 1 course of treatment every 4 months.

Other Information: Keep the carton and package insert. They contain important information. Store at 20-25°C (68-77°F). Keep product out of high heat and humidity. Protect product from moisture.

Directions: Adults 18 years of age and older (every 24 hours): This product is to be used once a day (every 24 hours). It may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours. Take every day for 14 days. Swallow 1 tablet with a glass of water before eating in the morning. Do not take more than 1 tablet a day. Do not chew or crush the tablets. Do not take for more than 14 days unless directed by your doctor. Do not use for more than 14 days unless directed by a doctor.

Repeat 14-Day Course (if needed): You may repeat a 14-day course every 4 months every 4 months unless directed by a doctor.

Children under 18 years of age: Ask a doctor.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Bold Cond. Oblique

TITLE (continued): 8 pt Helvetica Bold Cond. Oblique

Oblique

HEADING: 8 pt Helvetica Medium Cond

HEADING: 8 pt Helvetica Bold Cond. Oblique

SUBHEADING: 6 pt Helvetica Bold Cond.

TEXT: 6 pt Helvetica Medium Cond


LEADING: 5.9/6.5/6.6 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6 pt Helvetica Bold Cond.

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt



Omeprazole Delayed Release Tablets 20mg

Safety Feature -
Do not use if printed tablet blister unit is open or torn.

NDC (30) 400-00



Treats **Frequent** Heartburn!

Omeprazole Delayed Release Tablets 20mg

Acid Reducer

Omeprazole Delayed Release Tablets 20mg

42 Tablets
Three 14-day courses of treatment

Made in Israel
Manufactured by:
DEBCEL LTD.
Southern Industrial Zone
Or Akiva 30550, Israel

Drug Facts

Active ingredient (in each tablet)

Omeprazole delayed-release tablet, 20 mg

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- diazepam (anxiety medicine) ■ digoxin (heart medicine)
- tacrolimus (immune system medicine)
- atazanavir (medicine for HIV infection)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months ▶

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people got complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not chew or crush the tablets
- do not crush tablets in food
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age; ask a doctor

Other information

- read the directions, warnings, and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture ▼

Drug Facts (continued)

Inactive ingredients carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260