

084A



Batch

EXP

**CellCept®**  
(mycophenolate mofetil tablets)  
**500 mg tablets**  
NDC 0004-0260-01

NDC 0004-0260-01

**CellCept®**  
(mycophenolate mofetil tablets)  
**500 mg tablets**

Usual dosage: For dosage recommendations and other important prescribing information, read accompanying insert.

Dispense in tight, light-resistant container as defined in USP/NF.

**Store at 25°C (77°F); excursions permitted to 15°–30°C (59°–86°F).**

**Note to pharmacist:**  
Tamper-evident bottle enclosed.



Made in Ireland  
Distributed by:  
**Roche Laboratories Inc.**  
Nutley, New Jersey 07110

NDC 0004-0260-01

**CellCept®**  
(mycophenolate mofetil tablets)  
**500 mg**

Each tablet contains 500 mg mycophenolate mofetil.

100 tablets

Attention Pharmacist: Dispense the accompanying Medication Guide to each patient. For additional Medication Guides call 1-800-617-8191 or visit [www.rocheusa.com/products/CellCept](http://www.rocheusa.com/products/CellCept).

**Rx only**




3 0004-0260-014

27899636(A)



084A



**CellCept® Intravenous**  
(mycophenolate mofetil hydrochloride for injection)  
**500 mg**  
**4 Vials**  
NDC 0004-0298-09

NDC 0004-0298-09

**CellCept® Intravenous**  
(mycophenolate mofetil hydrochloride for injection)

CellCept Intravenous 500 mg vials must be reconstituted with 14 mL 5% Dextrose Injection USP. Further dilution of the content of the reconstituted vials with 5% Dextrose Injection USP is required to produce the infusion solution at the final concentration of 6 mg/mL. Refer to package insert for complete directions.

**OTHER INFUSION SOLUTIONS ARE NOT RECOMMENDED FOR DILUTION.**

USUAL DOSAGE: For dosage recommendations and other important prescribing information, read accompanying insert.

Store powder and reconstituted/infusion solutions at 25°C (77°F); excursions permitted to 15°–30°C (59°–86°F).

**COMMENCEMENT OF TREATMENT TO OCCUR WITHIN 4 HOURS FROM RECONSTITUTION AND DILUTION.**

NDC 0004-0298-09

**CellCept® Intravenous**  
(mycophenolate mofetil hydrochloride for injection)

**500 mg**

**4 Vials**

Made in Ireland

Distributed by:  
**Roche Laboratories Inc.**  
Nutley, New Jersey 07110

NDC 0004-0298-09

**CellCept® Intravenous**  
(mycophenolate mofetil hydrochloride for injection)

**500 mg**

**FOR INTRAVENOUS INFUSION ONLY**

Each vial contains the equivalent of 500 mg mycophenolate mofetil (as the hydrochloride salt), 25 mg polysorbate 80, and 5 mg citric acid.

**R<sub>x</sub> only**

Attention Pharmacist: Dispense the accompanying Medication Guide to each patient. For additional Medication Guides call 1-800-617-8191 or visit [www.rocheusa.com/products/CellCept](http://www.rocheusa.com/products/CellCept).

**4 Vials**



N 0004-0298-09 3

27899625(A)  
3 0 0 0 5 5 2 B

093A

NDC 0004-0259-43  
**CellCept®**  
(mycophenolate  
mofetil capsules)  
**250 mg capsules**

Batch

EXP

NDC 0004-0259-43


**CellCept®**  
(mycophenolate  
mofetil capsules)  
**250 mg capsules**

Usual dosage: For dosage  
recommendations and other  
important prescribing information,  
read accompanying insert.

Dispense in tight container  
as defined in USP/NF.

**Store at 25°C (77°F);  
excursions permitted to  
15°–30°C (59°–86°F).**

**Note to pharmacist:**  
Tamper-evident bottle enclosed.



Made in Ireland  
Distributed by:  
**Roche Laboratories Inc.**  
Nutley, New Jersey 07110

NDC 0004-0259-43

**CellCept®**  
(mycophenolate  
mofetil capsules)  
**250 mg**

Each capsule contains  
250 mg mycophenolate mofetil.

**500 capsules**

**Package not child-resistant**

**Attention Pharmacist:** Dispense the  
accompanying Medication Guide  
to each patient. For additional Medication  
Guides call 1-800-617-8191 or visit  
[www.rocheusa.com/products/CellCept](http://www.rocheusa.com/products/CellCept).


**R<sub>x</sub> only**



27899631(A)



093A



092A

NDC 0004-0259-01  
**CellCept**<sup>®</sup>  
 (mycophenolate  
 mofetil capsules)  
**250 mg capsules**

Batch

EXP

NDC 0004-0259-01  
**CellCept**<sup>®</sup>  
 (mycophenolate  
 mofetil capsules)  
**250 mg capsules**

Usual dosage: For dosage  
 recommendations and other  
 important prescribing information,  
 read accompanying insert.

Store at 25°C (77°F);  
 excursions permitted to  
 15°–30°C (59°–86°F).

**Note to pharmacist:**  
 Tamper-evident bottle enclosed.



Made in Ireland  
 Distributed by:  
**Roche Laboratories Inc.**  
 Nutley, New Jersey 07110


NDC 0004-0259-01  
**CellCept**<sup>®</sup>  
 (mycophenolate  
 mofetil capsules)  
**250 mg**

Each capsule contains  
 250 mg mycophenolate mofetil.

100 capsules

Attention Pharmacist: Dispense the  
 accompanying Medication Guide  
 to each patient. For additional Medication  
 Guides call 1-800-617-8191 or visit  
[www.rocheusa.com/products/CellCept](http://www.rocheusa.com/products/CellCept).

**R** only



0004-0259-018

27899629(A)



092A



NDC 0004-0261-29  
**CellCept®**  
**Oral Suspension**  
 (mycophenolate mofetil  
 for oral suspension)  
 LOT  
 EXPIRES

NDC 0004-0261-29  
**CellCept®**  
**Oral Suspension**  
 (mycophenolate mofetil  
 for oral suspension)  
**200 mg/mL**

Net contents after constitution: 175 mL,  
 containing 35.0 grams mycophenolate mofetil.  
 Deliverable volume after constitution:  
 160-165 mL.

Phenylketonurics: Contains 0.56 mg  
 phenylalanine per mL after constitution.  
 Usual dosage: For dosage recommendations,  
 handling and other important prescribing  
 information, read accompanying insert.

**Store dry powder at 25°C (77°F);  
 excursions permitted to 15°–30°C  
 (59°–86°F).**

**Store constituted suspension at  
 25°C (77°F); excursions permitted  
 to 15°–30°C (59°–86°F).**

**Storage in a refrigerator at 2°–8°C  
 (36°–46°F) is acceptable.  
 Do not freeze.**

Shake closed bottle well before using.  
 Discard any unused portion 60 days  
 after constitution.

**To the pharmacist:**

Tap the closed bottle several  
 times to loosen the powder.  
 Measure 94 mL of water in a  
 graduated cylinder.  
 Add approximately half the total  
 amount of water for constitution  
 to the bottle and shake the  
 closed bottle well for about  
 1 minute.

Add the remainder of water and  
 shake the closed bottle well for  
 about 1 minute.

Remove the child-resistant cap  
 and push bottle adapter into  
 neck of bottle.

Close bottle with child-resistant  
 cap tightly. This will assure the  
 proper seating of the bottle  
 adapter in the bottle, and  
 child-resistant status of the cap.

Dispense with patient instructions  
 and oral dispensers.

See enclosed patient instructions for  
 use of oral dispenser.

Contains 225 mL bottle with bottle  
 adapter and 2 oral dispensers.

Made in Ireland

Distributed by:  
**Roche Laboratories Inc.**  
 Nutley, New Jersey 07110

NDC 0004-0261-29

**CellCept®**  
**Oral Suspension**  
 (mycophenolate mofetil  
 for oral suspension)

**200 mg/mL**

Each mL contains  
 200 mg mycophenolate mofetil  
 after constitution.

**Attention Pharmacist: Dispense the  
 accompanying Medication  
 Guide to each patient. For additional Medication  
 Guides call 1-800-617-8191 or visit  
[www.rocheusa.com/products/CellCept](http://www.rocheusa.com/products/CellCept).**

**R<sub>x</sub> only**



27899633(A)





NDC 0004-0260-43  
**CellCept®**  
(mycophenolate  
mofetil tablets)  
**500 mg tablets**  
EXP  
Batch

NDC 0004-0260-43

**CellCept®**  
(mycophenolate  
mofetil tablets)  
**500 mg tablets**

Usual dosage: For dosage  
recommendations and other  
important prescribing information,  
read accompanying insert.

Dispense in tight, light-resistant  
container as defined in USP/NF.

**Store at 25°C (77°F);  
excursions permitted to  
15°–30°C (59°–86°F).**

**Note to pharmacist:**  
Tamper-evident bottle enclosed.

Made in Ireland  
Distributed by:  
**Roche Laboratories Inc.**  
Nutley, New Jersey 07110

NDC 0004-0260-43

**CellCept®**  
(mycophenolate  
mofetil tablets)  
**500 mg**

Each tablet contains  
500 mg mycophenolate mofetil.

**500 tablets**


**Package not child-resistant**

**Attention Pharmacist:** Dispense the  
accompanying Medication Guide  
to each patient. For additional Medication  
Guides call 1-800-617-8191 or visit  
[www.rocheusa.com/products/CellCept](http://www.rocheusa.com/products/CellCept).

**R<sub>x</sub> only**



27899627(A)



024A

Dispense in tight, light-resistant container as defined in USP/NF.

Store at 25°C (77°F); excursions permitted to 15°–30°C (59°–86°F).

Usual dosage: For dosage recommendations and other important prescribing information, read accompanying insert.

Made in Ireland

Distributed by:  
**Roche Laboratories Inc.**  
Nutley, New Jersey 07110

NDC 0004-0260-01

**CellCept®**  
(mycophenolate mofetil tablets)

**500 mg**

Each tablet contains 500 mg mycophenolate mofetil.

**100 tablets**  only

Attention Pharmacist: Dispense the accompanying Medication Guide to each patient.



EXP  
Batch



27899637



NDC 0004-0298-30

**CellCept®**  
**Intravenous**  
(mycophenolate  
mofetil hydrochloride  
for injection)

**500 mg****R<sub>x</sub> only****FOR INTRAVENOUS  
INFUSION ONLY**

Each vial contains the  
equivalent of 500 mg  
mycophenolate mofetil  
(as the hydrochloride salt),  
25 mg polysorbate 80, and  
5 mg citric acid.

**USUAL DOSAGE:** See  
package insert.

**Attention Pharmacist: Dispense  
the accompanying Medication  
Guide to each patient.**

**Store powder at 25°C (77°F);  
excursions permitted to  
15°–30°C (59°–86°F).**

Made in Ireland


Distributed by:  
**Roche Laboratories Inc.**  
Nutley, New Jersey 07110

27899624



(01) 103 0004 0298 30 4

3 0 0 0 5 5 3 B





Dispense in tight container as defined in USP/NF.  
**Package not child-resistant.**  
Store at 25°C (77°F); excursions permitted to 15°–30°C (59°–86°F).  
Usual dosage: For dosage recommendations and other important prescribing information, read accompanying insert.

Made in Ireland  
Distributed by:  
**Roche Laboratories Inc.**  
Nutley, New Jersey 07110

NDC 0004-0259-43

**CellCept®**  
(mycophenolate mofetil capsules)  
**250 mg**

Each capsule contains 250 mg mycophenolate mofetil.

**500 capsules**  **only**

Attention Pharmacist: Dispense the accompanying Medication Guide to each patient.



3 0004-0259-438

27899632



27899632

EXP  
Batch

Store at 25°C (77°F);  
excursions permitted to  
15°–30°C (59°–86°F).  
Usual dosage: For dosage  
recommendations and other  
important prescribing  
information, read  
accompanying insert.

Made in Ireland  
Distributed by:  
**Roche Laboratories Inc.**  
Nutley, New Jersey 07110

NDC 0004-0259-51

**CellCept®**  
(mycophenolate  
mofetil capsules)  
**250 mg**

Each capsule contains  
250 mg mycophenolate mofetil.

**120 capsules**  only

Attention Pharmacist: Dispense  
the accompanying Medication  
Guide to each patient.



EXP  
Batch

27899626



Store at 25°C (77°F);  
excursions permitted to  
15°–30°C (59°–86°F).  
Usual dosage: For dosage  
recommendations and other  
important prescribing  
information, read  
accompanying insert.

Made in Ireland  
Distributed by:  
**Roche Laboratories Inc.**  
Nutley, New Jersey 07110

NDC 0004-0259-01  
**CellCept®**  
(mycophenolate  
mofetil capsules)  
**250 mg**

Each capsule contains  
250 mg mycophenolate mofetil.

**100 capsules**   
Attention Pharmacist: Dispense  
the accompanying Medication  
Guide to each patient.



EXP  
Batch

27899630



Store dry powder at 25°C (77°F); excursions permitted to 15°–30°C (59°–86°F). Store constituted suspension at 25°C (77°F); excursions permitted to 15°–30°C (59°–86°F). Storage in a refrigerator at 2°–8°C (36°–46°F) is acceptable.

Do not freeze.

Shake closed bottle well before using.

Discard any unused portion 60 days after constitution.

Constitute Date:

LOT

EXP

27899634

NDC 0004-0261-29

# CellCept® Oral Suspension (mycophenolate mofetil for oral suspension)

## 200 mg/mL

Each mL contains  
200 mg mycophenolate mofetil  
after constitution.

**Attention Pharmacist: Dispense  
the accompanying Medication  
Guide to each patient.**

**R**only



Usual dosage: See package insert.  
Net contents after constitution: 175 mL,  
containing 35.0 grams mycophenolate mofetil.  
Deliverable volume after constitution: 160-165 mL.  
Phenylketonurics: Contains 0.56 mg  
phenylalanine per mL after constitution.  
**To the pharmacist:** Tap the closed bottle several  
times to loosen the powder. Measure 94 mL of water  
in a graduated cylinder. Add approximately half the  
total amount of water for constitution to the bottle  
and shake the closed bottle well for about 1 minute.  
Add the remainder of water and shake the closed  
bottle well for about 1 minute. Remove the child-  
resistant cap and push bottle adapter  
into neck of bottle. Close bottle with  
child-resistant cap tightly. This will  
assure the proper seating of the  
bottle adapter in the bottle, and  
child-resistant status of the cap.  
Dispense with patient instructions and  
oral dispensers. See enclosed patient  
instructions for use of oral dispenser.

Made in Ireland  
Distributed by:  
**Roche Laboratories Inc.**  
Nutley, New Jersey 07110



Dispense in tight, light-resistant container as defined in USP/NF.

**Package not child-resistant**

**Store at 25°C (77°F); excursions permitted to 15°–30°C (59°–86°F).**

Usual dosage: For dosage recommendations and other important prescribing information, read accompanying insert.

Made in Ireland

Distributed by:  
**Roche Laboratories Inc.**  
Nutley, New Jersey 07110

NDC 0004-0260-43

**CellCept®**  
**(mycophenolate mofetil tablets)**  
**500 mg**

Each tablet contains 500 mg mycophenolate mofetil.

**500 tablets** **R<sub>x</sub> only**

Attention Pharmacist: Dispense the accompanying Medication Guide to each patient.



27899628



EXP  
Batch

-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/

-----  
Ozlem Belen  
2/27/2009 05:13:38 PM