

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use **BICALUTAMIDE** safely and effectively. See full prescribing information for **BICALUTAMIDE** tablet, film coated for oral use.

INDICATIONS AND USAGE

- Bicalutamide tablets 50 mg are an androgen receptor inhibitor indicated for use in combination therapy with a luteinizing hormone-releasing hormone (LHRH) analog for the treatment of Stage II metastatic carcinoma of the prostate.
- Bicalutamide tablets 150 mg daily is not approved for use alone or with other treatments (see **CLINICAL STUDIES** (14.2)).

DOSEAGE AND ADMINISTRATION

The recommended dose for bicalutamide therapy in combination with an LHRH analog is one 50 mg tablet once daily (morning or evening), (2)

DOSEAGE FORMS AND STRENGTHS

- 50 mg tablets (3)

CONTRAINDICATIONS

- Hypersensitivity (4.1)
- Women (4.2)
- Pregnancy (4.3 and 5.1)

WARNINGS AND PRECAUTIONS

- Severe hepatic changes and hepatic failure have been observed rarely. Monitor serum transaminase levels daily after starting treatment with bicalutamide, at regular intervals for the first four months of treatment and periodically thereafter, and for symptoms of signs suggestive of hepatic dysfunction. Use bicalutamide with caution in patients with hepatic dysfunction. (5.1)
- Gynecomastric and breast pain have been reported during treatment with bicalutamide 50 mg when used as a single agent. (5.2)
- Bicalutamide is used in combination with a LHRH analog. LHRH agonists have been shown to cause a reduction in glucose tolerance in males. Consideration should be given to monitoring blood glucose in patients receiving bicalutamide in combination with LHRH agonists. (5.3)
- Monitoring Prostate Specific Antigen (PSA) is recommended. Evaluate for clinical progression if PSA increases. (5.4)

ADVERSE REACTIONS

Adverse reactions that occurred in more than 10% of patients receiving bicalutamide 50 mg daily were: hot flashes, pain (including general, back, pelvic and abdominal), asthenia, constipation, infection, nausea, peripheral neuropathy, dizziness, headache, somnolence, and anemia (8.1).

To report SUSPECTED ADVERSE REACTIONS, contact SANDOZ Inc. at 1-800-525-8747 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Bicalutamide is an inhibitor of CYP 3A4. Therefore, caution should be used when bicalutamide is co-administered with CYP 3A4 substrates. (7)
- Protein binding times should be closely monitored in patient already receiving coumestrol analogs who are started on bicalutamide. (7)

USE IN SPECIFIC POPULATIONS

Pediatric patients: Labeling describing pediatric clinical studies for bicalutamide is approved for AstraZeneca Pharmaceuticals LP's bicalutamide tablet. However, due to AstraZeneca Pharmaceuticals LP's marketing exclusivity rights, a description of these clinical studies is not approved for this bicalutamide labeling. (8.4)

See 17 for PATIENT COUNSELING INFORMATION

Proposed Patient Labeling Provided

FULL PRESCRIBING INFORMATION: CONTRAINDICATIONS**INDICATIONS AND USAGE****DOSEAGE AND ADMINISTRATION****DOSEAGE FORMS AND STRENGTHS****CONTRAINDICATIONS****WARNINGS AND PRECAUTIONS****ADVERSE REACTIONS****DRUG INTERACTIONS****USE IN SPECIFIC POPULATIONS****CLINICAL STUDIES****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****FULL PRESCRIBING INFORMATION****INDICATIONS AND USAGE****DOSEAGE AND ADMINISTRATION****DOSEAGE FORMS AND STRENGTHS****CONTRAINDICATIONS****WARNINGS AND PRECAUTIONS****ADVERSE REACTIONS****DRUG INTERACTIONS****USE IN SPECIFIC POPULATIONS****CLINICAL STUDIES****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING****17. PATIENT COUNSELING INFORMATION****DESCRIPTION****HOW SUPPLIED/STORAGE AND HANDLING**

Table 1. Incidence of Adverse Reactions (5% or Higher Treatment Group) Regardless of Causality

Body System Adverse Reaction	Treatment Group Number of Patients (%)	
	Placebo (n=407)	Plus LHRH Analog (n=407)
Body as a Whole		
Pain (General)	142 (35)	127 (31)
Back Pain	102 (25)	105 (26)
Aches	89 (22)	87 (21)
Pelvic Pain	85 (21)	70 (17)
Headache	57 (14)	57 (14)
Abdominal Pain	45 (11)	46 (11)
Chest Pain	34 (8)	38 (9)
Headache	29 (7)	27 (7)
Cardiovascular		
Hot Flashes	211 (53)	217 (53)
Hypertension	34 (8)	29 (7)
Digestive		
Constipation	87 (22)	69 (17)
Nausea	62 (15)	58 (14)
Diarrhea	49 (12)	107 (26)
Increased Liver Enzyme Test	37 (9)	48 (11)
Dyspepsia	30 (7)	23 (6)
Flatulence	26 (6)	22 (5)
Anorexia	25 (6)	28 (7)
Vomiting	24 (6)	35 (9)
Hematologic and Lymphatic		
Anemia	45 (11)	53 (13)
Metabolic and Nutritional		
Weight Loss	33 (8)	42 (10)
Weight Loss	30 (7)	39 (10)
Hypokalemia	26 (6)	27 (7)
Alkaline Phosphatase	22 (5)	24 (6)
Increased Weight Gain	22 (5)	18 (4)
Musculoskeletal		
Back Pain	37 (9)	43 (11)
Neck Pain	31 (8)	40 (10)
Arthritis	15 (4)	29 (7)
Myalgia	17 (4)	32 (8)
Nervous System		
Dizziness	41 (10)	35 (9)
Paresthesia	32 (8)	23 (6)
Insomnia	27 (7)	38 (10)
Headache	29 (7)	33 (8)
Depression	16 (4)	23 (6)
Respiratory System		
Cough	33 (8)	32 (8)
Cough Increased	33 (8)	24 (6)
Pharyngitis	32 (8)	23 (6)
Bronchitis	24 (6)	

(Continued)

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.
These are not all the possible side effects of bicalutamide tablets. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.
How should I store bicalutamide tablets?
Store at 20°-25°C (68°-77°F) (see USP Controlled Room Temperature).

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

General information about the safe and effective use of bicalutamide tablets.
Medicines are sometimes prescribed for purposes other than those listed in a patient information leaflet. Do not use bicalutamide tablets for a condition for which it was not prescribed. Do not give bicalutamide tablets to other people, even if they have the same symptoms that you have. It may harm them.

This patient information leaflet summarizes the most important information about bicalutamide tablets. If you would like more information about bicalutamide tablets talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about bicalutamide tablets that is written for health professionals. For more information call 1-800-525-8747.

What are the ingredients in bicalutamide tablets?
Active ingredients include: bicalutamide. Inactive ingredients include: corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, polyisobutyl 80, povidone, sodium starch glycolate, and titanium dioxide.

Sandoz Inc.
Princeton, NJ 08540

(Continued)

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.
These are not all the possible side effects of bicalutamide tablets. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.
How should I store bicalutamide tablets?
Store at 20°-25°C (68°-77°F) (see USP Controlled Room Temperature).

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

General information about the safe and effective use of bicalutamide tablets.
Medicines are sometimes prescribed for purposes other than those listed in a patient information leaflet. Do not use bicalutamide tablets for a condition for which it was not prescribed. Do not give bicalutamide tablets to other people, even if they have the same symptoms that you have. It may harm them.

This patient information leaflet summarizes the most important information about bicalutamide tablets. If you would like more information about bicalutamide tablets talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about bicalutamide tablets that is written for health professionals. For more information call 1-800-525-8747.

What are the ingredients in bicalutamide tablets?
Active ingredients include: bicalutamide. Inactive ingredients include: corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, polyisobutyl 80, povidone, sodium starch glycolate, and titanium dioxide.

Sandoz Inc.
Princeton, NJ 08540

(Continued)

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.
These are not all the possible side effects of bicalutamide tablets. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.
How should I store bicalutamide tablets?
Store at 20°-25°C (68°-77°F) (see USP Controlled Room Temperature).

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

General information about the safe and effective use of bicalutamide tablets.
Medicines are sometimes prescribed for purposes other than those listed in a patient information leaflet. Do not use bicalutamide tablets for a condition for which it was not prescribed. Do not give bicalutamide tablets to other people, even if they have the same symptoms that you have. It may harm them.

This patient information leaflet summarizes the most important information about bicalutamide tablets. If you would like more information about bicalutamide tablets talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about bicalutamide tablets that is written for health professionals. For more information call 1-800-525-8747.

What are the ingredients in bicalutamide tablets?
Active ingredients include: bicalutamide. Inactive ingredients include: corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, polyisobutyl 80, povidone, sodium starch glycolate, and titanium dioxide.

Sandoz Inc.
Princeton, NJ 08540

(Continued)

gen sensitive and responds to treatment that counteracts the effect of androgen and relieves the source of androgen.
When bicalutamide is combined with luteinizing hormone releasing hormone (LHRH) analogs, the suppression of serum testosterone induced by the LHRH analog is not affected. However, in clinical trials with bicalutamide as a single agent for prostate cancer, close to serum testosterone and estradiol have been noted.

In a subset of patients who have been treated with bicalutamide and an LHRH agonist, and who discontinue bicalutamide therapy due to progressive advanced prostate cancer, induction of Prostate Specific Antigen (PSA) after clinical improvement (androgen withdrawal phenomenon) may be observed.

12.3 Pharmacokinetics
Bicalutamide is well-absorbed following oral administration, although the absolute bioavailability is unknown. Co-administration of bicalutamide with food has no clinically significant effect on rate or extent of absorption.

Distribution
Bicalutamide is highly protein-bound (96%) (see DRUG INTERACTIONS 7).
Metabolism/Excretion: Bicalutamide is extensively metabolized. The 5 α -reduced isomer is metabolized primarily by glucuronidation. The 5 α -reduced isomer also undergoes glucuronidation but is predominantly excreted as an inactive metabolite following glucuronidation. Both the parent and metabolite glucuronides are eliminated in the urine and feces. The 5 α -metabolite is rapidly cleared relative to the 5 α -parent, with the 5 α -metabolite accounting for about 96% of total steady-state plasma levels.

There was no significant difference in time to objective tumor progression between treatment groups (see Figure 2).
Objective tumor progression was defined as the appearance of any bone metastasis or the worsening of any existing bone metastases on bone scan attributable to metastatic disease, or an increase by 25% or more of any existing tumor measurements. The hazard ratio for time to progression of bicalutamide plus LHRH analog to that of flutamide plus LHRH analog was 1.0 (95% confidence interval, 0.79 to 1.31).

Pharmacokinetics of the active enantiomer of bicalutamide in normal males and patients with prostate cancer are presented in Table 3.

Parameter	Mean	Standard Deviation
Normal Males (n=30)		
Agreement Coef. (C ₁₂ vs C ₁₈) (Linicity)	0.920	0.103
Single Dose Peak Concentration (ng/mL)	0.168	0.178
Single Dose Time to Peak (hours)	31.3	14.6
Half-life (hours)	5.8	2.29
Patients with Prostate Cancer (n=48)		
C ₁₂ (ng/mL)	0.939	3.304
C ₁₈ (ng/mL)		
C ₁₂ :C ₁₈ Mean Steady-State Concentration		

13. NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
Two-year oral carcinogenicity studies were conducted in both male and female rats and mice at doses of 5, 15 or 75 mg/kg/day of bicalutamide. A variety of tumor target organ effects were identified and were attributed to the androgenic activity of bicalutamide. In male rats, the most significant effects (Leydig) cell tumors in male rats at all dose levels (the steady-state plasma concentration with the 5 mg/kg/day bicalutamide approximately 235 times the specific concentrations) and ovarian adenocarcinomas in female rats at 75 mg/kg/day (approximately 1-1/2 times the human therapeutic concentration). There is no evidence of Leydig cell hyperplasia in patients; uterine tumors are not relevant to the human patient population.
A small increase in the incidence of hepatocellular carcinoma in male mice gives 75 mg/kg/day of bicalutamide approximately 4 times human therapeutic concentrations) and an increased incidence of benign thyroid follicular cell adenomas in mice given 5 mg/kg/day (approximately 23 human therapeutic concentrations) and above were recorded. These neoplastic changes were progressive or non-reversible changes related to hepatic enzyme induction observed in animal toxicity studies. Enzyme induction has not been observed following bicalutamide administration in man. There were no tumorigenic effects suggestive of genotoxic carcinogenesis.
A comprehensive battery of both *in vitro* and *in vivo* genotoxicity tests (yeast gene conversion, Ames, E. coli, CHO/HGPRT, human lymphocyte cytogenetic, mouse micronucleus, and *in vivo* mouse cytogenetic tests) has demonstrated that bicalutamide does not have genotoxic activity.
Administration of bicalutamide may lead to inhibition of spermatogenesis. The long-term effects of bicalutamide on male fertility have not been studied.
In male rats dosed at 250 mg/kg/day (approximately 5 times human therapeutic concentrations), the preovulatory interval and time to successful mating were increased in the first pairing but no effects on fertility following successful mating were seen. These effects were reversed by 7 weeks after the end of a 11-month period of dosing.
No effects on female rats dosed at 10, 50 and 250 mg/kg/day (approximately 2, 1, and 2 times human therapeutic concentrations, respectively) or their female offspring were observed. Administration of bicalutamide to pregnant females resulted in feminization of the male offspring leading to hypospadias at all dose levels. Affected male offspring were also born.

14. CLINICAL STUDIES

14.1 Bicalutamide Tablets 50 mg Daily in Combination with an LHRH-A
In a multicenter, double-blind, controlled clinical trial, 813 patients with previously untreated advanced prostate cancer were randomized to receive bicalutamide 50 mg daily (404 patients) or flutamide 250 mg (409 patients) three times a day, each in combination with LHRH analogs (either goserelin acetate or leuprolide acetate depot).
In an analysis conducted after a median follow-up of 169 weeks was reached, 215 (53.2%) patients treated with bicalutamide-LHRH analog therapy and 235 (57.5%) patients treated with flutamide-LHRH analog therapy had died. There was no difference in the cause of death between treatment groups (see Figure 1). The hazard ratio for time to death (survival) was 1.07 (95% confidence interval 0.72 to 1.50).

Figure 1. The Kaplan-Meier probability of death for both androgen treatment groups.





Figure 2. Kaplan-Meier curve for time to progression for both androgen treatment groups.



Quality of life was assessed with self-administered patient questionnaires on pain, social functioning, emotional well being, vitality, activity limitation, self disability, overall health, physical capacity, cognitive symptoms, and treatment related symptoms. Assessment of the Quality of Life questionnaires did not indicate consistent significant differences between the two treatment groups.

14.2 Safety Data from Clinical Studies using Bicalutamide 150 mg

Bicalutamide 150 mg is not approved for use as either alone or with other treatments.
Two identical multicenter, randomized, open-label trials comparing bicalutamide 150 mg daily monotherapy to castration were conducted in patients that had fully advanced (T3-4, N1, M0) or metastatic (M1) prostate cancer.

Monotherapy - M1 Group

Bicalutamide 150 mg daily is not approved for use in patients with M1 cancer of the prostate. Based on an interim analysis of the two trials for survival, the Data Safety Monitoring Board recommended that the bicalutamide treatment be discontinued in all M1 patients because the risk of death was 25% (HR 1.25, 95% CI 0.87 to 1.81) and 31% (HR 1.31, 95% CI 0.87 to 1.97) higher in the bicalutamide treated group compared to castration in the castration group, respectively.
Locally Advanced (T3-4, N1, M0) Group
Bicalutamide 150 mg daily is not approved for use in patients with locally advanced (T3-4, N1, M0) cancer of the prostate. Following discontinuation of all M1 patients, the trials continued with the T3-4, N1, M0 patients until study completion. In the lower (HR 0.92), the risk of death was 25% (HR 1.25, 95% CI 0.87 to 1.71) higher in the bicalutamide group and in the smaller (HR 1.45), the risk of death was 33% (HR 1.54, 95% CI 0.97 to 2.39) higher in the bicalutamide group.

In addition to the above two studies, there are three other on-going clinical studies that provide additional safety information for bicalutamide 150 mg, a dose that is not approved for use. These are two multicenter, randomized, double-blind, parallel group trials comparing bicalutamide 150 mg daily monotherapy (adjusted to placebo or under watchful waiting) with placebo, for death to time to disease progression, in a population of 813 patients with localized or locally advanced prostate cancer.
Bicalutamide 150 mg daily is not approved for use as therapy for patients with localized prostate cancer who are candidates for watchful waiting. Data from a planned subgroup analysis of two of these trials in 1627 patients with localized prostate cancer who were under watchful waiting, revealed a trend toward decreased survival in the bicalutamide arm after a median follow-up of 47.4 years versus 46.7 years in the placebo arm (24 (2.7%) deaths in the bicalutamide patients versus 279 (22.9%) deaths in the placebo treated patients) (localised prostate cancer) (see Table 1).

16. HOW SUPPLIED/STORAGE AND HANDLING

Bicalutamide tablets, for oral use, are available as:
50 mg: round, white, convex tablet core, debossed 32 on one side and 212 on the reverse side, film coated, white, and supplied as:
NDC 0781-5403-31 bottles of 30
NDC 0781-5403-10 bottles of 100
NDC 0781-5402-10 bottles of 1000
NDC 0781-5402-30 bottles of 30

17. Storage and Handling

Store at 20°-25°C (68°-77°F) (see USP Controlled Room Temperature).

18. PATIENT COUNSELING INFORMATION

Patients should be counseled that while taking bicalutamide tablets and the LHRH analog should be started at the same time and that they should not interrupt or stop taking these medications without consulting their physician.
During treatment with bicalutamide tablets, constipation has been reported, and those patients who experience this symptom should observe caution when driving or operating machinery or when performing tasks that require alertness.
Patients should be informed that diabetes, or loss of glycemic control in patients with pre-existing diabetes, has been reported during treatment

with LHRH agonists. Consideration should therefore be given to monitoring blood glucose in patients receiving bicalutamide tablets in combination with LHRH agonists.

PATIENT INFORMATION BICALUTAMIDE TABLETS

Read the Patient Information that comes with bicalutamide tablets before you start taking it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your healthcare provider about your medical condition or your treatment.

What is bicalutamide tablets?

Bicalutamide tablets is a prescription medicine called an androgen receptor inhibitor, used in combination with luteinizing hormone-releasing hormone (LHRH) medicines to treat stage IV metastatic prostate cancer. It is not known if bicalutamide tablets is safe and effective in children.

Who should not take bicalutamide tablets?

- Do not take bicalutamide tablets if:
 - you are allergic to any of the ingredients in bicalutamide tablets. See the end of this leaflet for a complete list of ingredients.
 - you are a woman.

What should I tell my healthcare provider before taking bicalutamide tablets?

Before you take bicalutamide tablets, tell your healthcare provider about all your other medications, including LHRH bicalutamide tablets, if you are pregnant or think you may be pregnant.

What should I take bicalutamide tablets with?

- Take a medicine to thin your blood. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take bicalutamide tablets with your medicine a blood thinner.
- Take bicalutamide tablets in combination with LHRH medicines.
- Take a medicine to control your blood sugar. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your blood pressure. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your cholesterol. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your diabetes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your constipation. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your pain. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your nausea. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your vomiting. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your diarrhea. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dizziness. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your headache. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your fatigue. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your weakness. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your loss of appetite. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your weight gain. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your hair loss. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry mouth. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry cough. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry skin. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry hair. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry eyes. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry nose. Ask your healthcare provider or pharmacist if you just refuse it your medicine a blood thinner.
- Take a medicine to control your dry throat. Ask