

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.  
Dissolves in any beverage.  
Distributed by Kremers Urban, LLC, Princeton, NJ 08540, USA

### Drug Facts (continued)

If pregnant or breast feeding, ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not take more than directed unless advised by your doctor
- the dosing cup supplied is marked to contain 17 grams of powder when filled to the indicated line
- adults and children 17 years of age and older:
  - fill to indicated line for the correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - use once daily
  - use no more than 7 days
- children 16 years of age or under: ask a doctor

### Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

**Inactive ingredients** none

### Questions or comments?

1-800-477-7877

NDC 62175-195-07

Original  
Prescription  
Strength

## Polyethylene Glycol 3350 Powder for Solution, Laxative

- \* Relieves Occasional Constipation
- \* Softens Stool

CAP is not a dosing CUP.

Use dosing CUP provided to measure dose.

NET WT 4.1 OZ (119 g) 7 ONCE-DAILY DOSES

### Drug Facts

Active ingredient (in each dose)	Purpose
Polyethylene Glycol 3350, 17 g (dosing cup filled to line)....	Laxative

### Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

### Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are** taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

### Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

4010359 Rev. 04/09



Exp.:

Lot: 

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

*Dissolves in Any Beverage*

### Drug Facts (continued)

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not take more than directed unless advised by your doctor
- the dosing cup supplied is marked to contain 17 grams of powder when filled to the indicated line
- adults and children 17 years of age and older:
  - fill to indicated line for correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - use once daily
  - use no more than 7 days
- children 16 years of age or under: ask a doctor

### Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

**Inactive ingredients** none

### Questions or comments?

1-800-477-7877

*Distributed by Kremers Urban, LLC, Princeton, NJ 08540 USA*

NDC 62175-195-15

Original  
Prescription  
Strength

# Polyethylene Glycol 3350 Powder for Solution, Laxative

\* *Relieves Occasional Constipation*

\* *Softens Stool*

**CAP** is not a dosing **CUP**.  
Use dosing **CUP** provided to measure dose.

NET WT 8.9 OZ (255 g)

15 ONCE-DAILY DOSES

*Polyethylene Glycol 3350 Powder for Solution increases frequency of bowel movements and softens the stool.*

### Drug Facts

**Active ingredient (in each dose)** Polyethylene Glycol 3350, 17 g (dosing cup filled to line)....**Purpose** Laxative

### Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are** taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week



Lot.:  
Exp.:

4010420 Rev. 04/09

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

*Dissolves in Any Beverage*

NDC 62175-195-31

**Original  
Prescription  
Strength**

*Polyethylene Glycol 3350 Powder for Solution  
increases frequency of bowel movements  
and softens the stool.*

### Drug Facts (continued)

If pregnant or breast feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not take more than directed unless advised by your doctor
- the dosing cup supplied is marked to contain 17 grams of powder when filled to the indicated line
- adults and children 17 years of age and older:
  - fill to indicated line for the correct dose (17 g)
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - use once daily
  - use no more than 7 days
- children 16 years of age or under: ask a doctor

### Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

**Inactive ingredients** none

### Questions or comments?

1-800-477-7877

*Distributed by Kremers Urban, LLC, Princeton, NJ 08540 USA*

# Polyethylene Glycol 3350 Powder for Solution, Laxative

*\* Relieves Occasional Constipation*

*\* Softens Stool*

**CAP** is not a dosing **CUP**.

Use dosing **CUP** provided to measure dose.

**NETWT 18.5 OZ (527 g) 31 ONCE-DAILY DOSES**

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

### Drug Facts

Active ingredient (in each dose)	Purpose
Polyethylene Glycol 3350, 17 g (dosing cup filled to line)....	Laxative

### Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are** taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week



Lot: Exp.:

4010421  
Rev. 04/09



FOLD AT LINE  
TEAR AT ARROW

NDC 62175-195-01

Original  
Prescription  
Strength

# Polyethylene Glycol 3350 Powder for Solution, Laxative

- \* *Relieves Occasional Constipation*
- \* *Softens Stool*

NET WT 0.5 OZ (17g)

1 Dose

## Drug Facts

Active ingredient (in each dose)	Purpose
Polyethylene Glycol 3350, 17 g.....	Laxative

### Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are** taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

## Drug Facts (continued)

**If pregnant or breast feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- **do not take more than directed unless advised by your doctor**
- adults and children 17 years of age and older:
  - stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - use once daily
  - use no more than 7 days
- children 16 years of age or under: ask a doctor

### Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil is open or broken

**Inactive ingredients** none

### Questions or comments?

1-800-477-7877



Distributed by Kremers Urban, LLC  
Princeton, NJ 08540 USA L5480 Rev. 04/09

Original  
Prescription  
Strength

# Polyethylene Glycol 3350 Powder for Solution, Laxative

NDC 62175-195-10

Original  
Prescription  
Strength

## Polyethylene Glycol 3350 Powder for Solution, Laxative

- \* Relieves Occasional Constipation
- \* Softens Stool
- \* Dissolves in Any Beverage

10 PACKETS, EACH  
PACKET NET WT 0.5 OZ (17g)

**10**  
ONCE-DAILY DOSES

Original  
Prescription  
Strength

## Polyethylene Glycol 3350 Powder for Solution, Laxative

- \* Relieves Occasional Constipation
- \* Softens Stool
- \* Dissolves in Any Beverage

CONTAINS 10 ONCE-DAILY DOSES  
EACH PACKET CONTAINS ONE DOSE NET WT 0.5 OZ (17G)

Original  
Prescription  
Strength

## Polyethylene Glycol 3350 Powder for Solution, Laxative

- \* Relieves Occasional Constipation
- \* Softens Stool
- \* Dissolves in Any Beverage



### Drug Facts

**Active ingredient (in each dose)** Purpose  
Polyethylene Glycol 3350, 17 g..... Laxative

#### Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

#### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol.  
**Do not use** if you have kidney disease, except under the advice and supervision of a doctor.

#### Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are** taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

#### Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

Distributed by Kremers Urban, LLC, Princeton, NJ 08540 USA

Control No.  
Exp. Date

Unvarnished Area



Barcode Area



CR5481  
Rev. 04/09