

Dissolves in Any Beverage



NDC 40032-312-07

Polyethylene Glycol 3350 Powder for Oral Solution

Laxative

7 ONCE-DAILY DOSES
NET WT 4.1 OZ (119 g)

- Relieves Occasional Constipation
- Softens Stool



Novel Laboratories, Inc. Somerset, NJ 08873

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool



9
40032-312-07
3

Drug Facts

Active Ingredient (in each dose)

Polyethylene Glycol 3350, 17 g (cap filled to line) Laxative

Purpose

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

PEEL HERE

Manufactured by:
Novel Laboratories, Inc.
Somerset, NJ 08873
NLF-312-07-00

Lot
Exp.

Drug Facts (continued)

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Drug Facts (continued)

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week
- side effects occur. You may report side effects to FDA at 1-800-FDA-1088

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a POISON CONTROL CENTER right away.

Drug Facts (continued)

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line
- adults and children 17 years of age and older:
 - fill to top of line in cap which is marked to indicate the correct dose (17 g)
 - Stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- Use once a day
- Use no more than 7 days
- Children 16 years of age or under: ask a doctor

Drug Facts (continued)

Other Information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive Ingredients none

Questions or comments?

1-800-FDA-1088 (1-800-332-1088) or www.fda.gov

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool

NDC 40032-312-08



Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a POISON CONTROL CENTER right away.

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line
- adults and children 17 years of age and older:
 - fill to top of line in cap which is marked to indicate the correct dose (17 g)
 - Stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - Use once a day
 - Use no more than 7 days
- Children 16 years of age or under: ask a doctor

Other Information

- store at 20° - 25°C (68°- 77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive Ingredients none

Questions or comments?

1-800-FDA-1088 (1-800-332-1088) or www.fda.gov



Novel Laboratories, Inc. Somerset, NJ 08873

Laxative

- Relieves Occasional Constipation
- Softens Stool

NET WT 8.3 OZ (238 g) 14 ONCE-DAILY DOSES

Dissolves in Any Beverage

Drug Facts

Active Ingredient (in each dose)

Polyethylene Glycol 3350, 17 g (cap filled to line) Laxative

Purpose

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week
- side effects occur. You may report side effects to FDA at 1-800-FDA-1088



N 3 40032-312-08 6

Lot
Exp.

NLB-312-08-00

