

FOR FDA SUBMISSION ONLY

Drug Facts (continued)

Active ingredients (in each tablet)

Fexofenadine HCl 60 mg
Pseudoephedrine HCl 120 mg

Purpose

Antihistamine
Nasal decongestant

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: sneezing, runny nose, itchy nose or throat, itchy, watery eyes, redness or swelling of the nasal passages

temporarily relieves sinus congestion and pressure

temporarily restores freer breathing through the nose

Drug Facts (continued)

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI)
- certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing

Ask a doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

Drug Facts (continued)

Directions

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

adults and children 12 years and over: take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours

children under 12 years of age: do not use

adults 65 years of age and older: ask a doctor

consumers with kidney disease: ask a doctor

Other information

- safety sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° and 25°C (68° and 77°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, stearic acid

Questions or comments?

call toll-free 1-800-633-1610 or www.allegra.com

NDC 41167 4310 2

Allegra-D®
12 Hour Allergy & Congestion

fexofenadine HCl 60 mg/antihistamine
pseudoephedrine HCl 120 mg/nasal decongestant
Extended Release Tablets

10 Tablets

12 HOUR

Allegra-D®
12 Hour Allergy & Congestion

The makers of Allegra® do not make store brand products.
The trade dress of this Allegra® package is subject to trademark protection.
Dist. By: Chatten, Inc. (part of the sanofi-aventis Group),
Chattanooga, TN 37409-0219 ©2010

50094535

0 41167 43102 3

Drug Facts (continued)

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI)
- certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing

Ask a doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

Drug Facts (continued)

Directions

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

adults and children 12 years and over: take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours

children under 12 years of age: do not use

adults 65 years of age and older: ask a doctor

consumers with kidney disease: ask a doctor

Other information

- safety sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° and 25°C (68° and 77°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, stearic acid

Questions or comments?

call toll-free 1-800-633-1610 or www.allegra.com

Allegra-D®
12 Hour Allergy & Congestion

12 HOUR

ORIGINAL PRESCRIPTION STRENGTH
NON DROWSY

RELIEF OF:

- ✓ Nasal Congestion
- ✓ Sinus Congestion & Pressure
- ✓ Runny Nose
- ✓ Sneezing
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

Drug Facts
(continued)

Warnings

- Do not use if you have ever had an allergic reaction to this product or any of its ingredients
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- If you have difficulty swallowing

Ask a doctor before use if you have

- heart disease
- thyroid disease
- glaucoma
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever.
- you get nervous, dizzy, or sleepless

Drug Facts (continued)

Do not use

- If you have ever had an allergic reaction to this product or any of its ingredients
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- If you have difficulty swallowing

Ask a doctor before use if you have

- heart disease
- thyroid disease
- glaucoma
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever.
- you get nervous, dizzy, or sleepless

Drug Facts (continued)

Warnings

- Do not use if you have ever had an allergic reaction to this product or any of its ingredients
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- If you have difficulty swallowing

Ask a doctor before use if you have

- heart disease
- thyroid disease
- glaucoma
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever.
- you get nervous, dizzy, or sleepless

Drug Facts
(continued)

Active ingredients (in each tablet)

- Fexofenadine HCl 60 mg
- Pseudoephedrine HCl 120 mg

Purpose

- Antihistamine
- Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: runny nose, sneezing, nasal congestion
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: itchy, watery eyes
- temporarily relieves sinus congestion and pressure
- temporarily restores free breathing through the nose

NDC 41167 4310 4

Allegra-D®
12 Hour Allergy & Congestion

fexofenadine HCl 60 mg/antihistamine
pseudoephedrine HCl 120 mg/nasal decongestant
Extended Release Tablets

20 Tablets

12 HOUR

**ORIGINAL PRESCRIPTION STRENGTH
NON DROWSY**

Allegra-D®
12 Hour Allergy & Congestion

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.
Dist. By: Chatten, Inc. (part of the sanofi-aventis Group), Chattanooga, TN 37409-0219 ©2010

50094538



0 4 1 1 6 7 4 3 1 0 4 7

Drug Facts (continued)

Directions

- do not divide, crush, chew or dissolve the tablet; swallow tablet whole
- adults and children 12 years and older: take 1 tablet with a glass of water every 12 hours on an empty stomach, do not take more than 2 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- safely sealed; do not use if carton is opened or if individual blister units are torn or opened
- store between 20° and 25°C (68° and 77°F)

Inactive ingredients

canbada wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, stearic acid

Questions or comments?

call toll-free 1-800-633-1610 or www.allegra.com

Drug Facts (continued)

Warnings

- Do not use if you have ever had an allergic reaction to this product or any of its ingredients
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- If you have difficulty swallowing

Ask a doctor before use if you have

- heart disease
- thyroid disease
- glaucoma
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever.
- you get nervous, dizzy, or sleepless

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER
01/24/2011