

03771600



NO INK
NO VARNISH

Drug Facts (continued)	Drug Facts (continued)
<ul style="list-style-type: none">■ frequent chest pain■ frequent wheezing, particularly if with heartburn■ unexplained weight loss■ nausea or vomiting■ stomach pain■ a sodium-restricted diet Ask a doctor or pharmacist before use if you are <ul style="list-style-type: none">■ taking blood-thinning medicines (blood-thinning medicines)■ prescription antibiotics or anti-fungal medicines■ digoxin (heart medicine)■ tacrolimus (immune system medicine)■ prescription antineoplastic medicines for HIV infection)■ any other prescription drugs. Sodium bicarbonate may interact with certain prescription drugs. Stop use and ask a doctor if <ul style="list-style-type: none">■ your heartburn continues or worsens■ you need to take this product for more than 14 days■ you need to take more than 1 course of treatment every 4 months If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	Directions <ul style="list-style-type: none">■ for adults 18 years of age and older■ this product is to be used once a day (every 24 hours), every day for 14 days■ if you take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours 14-Day Course of Treatment <ul style="list-style-type: none">■ swallow 1 capsule with a glass of water at least 1 hour before eating in the morning■ take every day for 14 days■ do not take more than 1 capsule a day■ do not chew or crush the capsule■ do not open capsule and sprinkle on food■ do not use for more than 14 days unless directed by your doctor Repeated 14-Day Courses (if needed) <ul style="list-style-type: none">■ you may repeat a 14-day course every 4 months■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor■ children under 18 years of age: ask a doctor Heartburn in children may sometimes be caused by a serious condition.

NO INK
NO VARNISH

1.8111

3.5434

Treats Frequent Heartburn

Zegerid[®] otc[®]

Tips for Managing Heartburn

- Do not lie flat or bend over soon after eating
- Do not eat late at night or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose-fitting clothing around your stomach

FREE STARTER PACK

NDC 11523-7285-5

Treats Frequent Heartburn

Zegerid[®] otc[®]

- *Omeprazole 20 mg/Acid Reducer*
- *Sodium Bicarbonate 1100 mg!*
- *Allows Absorption of this Omeprazole Product*

CAPSULES
2
Follow Samples with a 14-Day Course of Treatment

SAMPLE-NOT FOR SALE

Drug Facts (continued)
Other Information <ul style="list-style-type: none">■ each capsule contains: sodium 803 mg■ read the directions, warnings and accompanying label information before use■ store at 20°-25° (68°-77°)■ tamper-evident: do not use if the blue band around the capsule is missing or broken. Do not use if foil inner seal imprinted with "Sealed for your protection" is missing, open or broken.■ keep product out of high heat and humidity■ protect product from moisture Inactive Ingredients <ul style="list-style-type: none">■ croscarmellose sodium, FD&C blue No. 1, gelatin, croscarmellose sodium, pharmaceutical ink, polysorbate 80, titanium dioxide Questions or comments? <ul style="list-style-type: none">■ Call 1-888-4-ZEG-OTC (1-888-483-4822) between 8:00 AM and 5:00 PM Central Time, Monday through Friday or visit www.ZegeridOTC.com <p><small>© Copyright © Distributed by MSD Consumer Care, Inc., PO Box 377, Memphis, TN 38151 USA, a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ, USA. All rights reserved. Product of Spain. 37716-01 PKG00906</small></p> <p> recyclable carton</p>

LOT/EXP
Area
No ink
No Varnish

Drug Facts	Purpose
Active Ingredients (in each capsule) <ul style="list-style-type: none">Omeprazole 20 mg..... Acid reducerSodium Bicarbonate 1100 mg.......... Allows absorption of this omeprazole product	
Use <ul style="list-style-type: none">■ Treats frequent heartburn (occurs 2 or more days a week)■ not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.	
Warnings <p>Allergy alert: Do not use if you are allergic to omeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none">■ trouble or pain swallowing food■ vomiting with blood■ bloody or black stools <p>These may be signs of a serious condition. See your doctor.</p>	
Ask a doctor before use if you have <ul style="list-style-type: none">■ had heartburn over 3 months. This may be a sign of a more serious condition.■ heartburn with light-headedness, sweating or dizziness <ul style="list-style-type: none">■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness	

PRINTSIDE GLUE

FOLD

PRINTSIDE GLUE

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd, Bold Italic

Drug Facts (continued)

Drug Facts - 8 pt Helvetica Cnd, Bold Italic (continued) - 8 pt Helvetica Cnd.

Headings - 8 pt Helvetica Cnd, Bold Italic

Sub Heads - 6 pt Helvetica Cnd, Bold

Text - 6 pt Helvetica Cnd, minimum leading 6.5

Quota e Bullets - 3.5 pt Zaph Dingbats no compression

Border - 1 pt rule, 3/64" gap all around

Barline - 1 pt rule _____

Headline - 5 pt rule _____

Certified by: eh

APD- 2011-91886

06.06.11 - ju

1.8111 3.5434

03658905



NO INK
NO VARNISH

Drug Facts (continued)

- frequent heart pain
 - frequent wheezing, particularly with heartburn
 - unexplained weight loss
 - nausea or vomiting
 - stomach pain
 - a sodium-restricted diet
- Ask a doctor or pharmacist before use if you are taking**
- warfarin, dabigatran or clostazol (blood-thinning medicines)
 - prescription antihistalgic or anti-yeast medicines
 - diclofenac (arthritis medicine)
 - digoxin (heart medicine)
 - baclofen (muscle system medicine)
 - prescription antiretroviral medicines for HIV infection)
 - any other prescription drugs. Sodium bicarbonate may interact with certain prescription drugs.
- Stop use and ask a doctor if**
- you heartburn continues or worsens
 - you need to take this product for more than 14 days
 - you need to take more than 1 course of treatment every 4 months
- If pregnant or breast-feeding, ask a health professional before use.**
- Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

- Directions**
- for adults 18 years of age and older
 - this product is to be used once a day (every 24 hours), every day for 14 days
 - it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- 14-Day Course of Treatment**
- swallow 1 capsule with a glass of water at least 1 hour before eating in the morning
 - take every day for 14 days
 - do not take more than 1 capsule a day
 - do not chew or crush the capsule
 - do not open capsule and sprinkle on food
 - do not use for more than 14 days unless directed by your doctor
- Repeated 14-Day Courses (if needed)**
- you may repeat a 14-day course every 4 months
 - do not take for more than 14 days or more often than every 4 months unless directed by a doctor
 - children under 18 years of age, ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

- Other Information**
- each capsule contains sodium 303 mg
 - read the directions, warnings and accompanying label information before use
 - store at 20°-25°C (68°-77°F)
 - tamper-evident: do not use if the blue band around the capsule is missing or broken. Sealed for your protection
 - is missing, open or broken.
 - keep product out of high heat and humidity
 - protect product from moisture
- Inactive Ingredients**
- croscarmellose sodium, FD&C blue No. 1, gelatin, magnesium stearate, pharmaceutical ink, polysorbate 80, titanium dioxide
- Questions or comments?**
- Call 1-888-4-226-0100 (1-888-485-4662) between 8:00 AM and 3:00 PM Central Time, Monday through Friday or visit www.ZegeridOTC.com

© Copyright & Distributed by MSD Consumer Care, Inc., a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ 08802, USA. All rights reserved. Product of Spain. 35589-US/PK501722



NO INK
NO VARNISH



Treats Frequent Heartburn

Zegerid[®] OTC[®]

• Omeprazole 20 mg/Acid Reducer
• Sodium Bicarbonate 1100 mg/
Allows Absorption of this Omeprazole Product

14 CAPSULES
One 14-Day Course of Treatment



LDT/EXP
Area
No ink
No Varnish

Drug Facts

- | Active Ingredients (in each capsule) | Purpose |
|--------------------------------------|---|
| Omeprazole 20 mg..... | Acid reducer |
| Sodium Bicarbonate 1100 mg..... |Allows absorption of this omeprazole product |
- Use**
- treats frequent heartburn (occurs 2 or more days a week)
 - not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.
- Warnings**
- Allergy alert:** Do not use if you are allergic to omeprazole
- Do not use if you have:**
- trouble or pain swallowing food
 - vomiting with blood
 - bloody or black stools
- These may be signs of a serious condition. See your doctor.
- Ask a doctor before use if you have**
- had heartburn over 3 months. This may be a sign of a more serious condition.
 - heartburn with lightheadedness, sweating or dizziness
 - chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness

PRINTSIDE GLUE

FOLD

PRINTSIDE GLUE

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd, Bold Italic
Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd, Bold Italic (continued) - 5 pt Helvetica Cnd
Headings - 8 pt Helvetica Cnd, Bold Italic
Sub Heads - 6 pt Helvetica Cnd, Bold
Text - 6 pt Helvetica Cnd, m n min leading 6.5
Square Bullets - 3.5 pt Zaph D/ingpals no compression
Border - 1 pt rule, 3/64" gap all around
Barline - 1 pt rule _____
Hairline - .5 pt rule _____

Certified by: eb
APO #: 2011-91877
06.06.11 - ju

1.8111 [46.00]

4.7638 [121.00]

50585920



NO INK
NO VARNISH

Drug Facts	Purpose
Active Ingredients (in each capsule) Omeprazole 20 mg. Sodium Bicarbonate 1100 mg. Allows absorption of this omeprazole product.	Acid reducer
Use ■ Treats frequent heartburn (occurs 2 or more days a week). ■ Not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.	
Warnings Allergy alert: Do not use if you are allergic to omeprazole. Do not use if you have: ■ trouble or pain swallowing food ■ vomiting with blood ■ bloody or black stools These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ a sodium-restricted diet Ask a doctor or pharmacist before use if you are taking ■ warfarin, clopidogrel or diltiazem (blood-thinning medicines) ■ prescription antifungal or anti-yeast medicines	
Drug Facts (continued) ■ diltiazem (heart medicine) ■ dipyrin (heart medicine) ■ baclofen (muscle system medicine) ■ prescription antiretrovirals (medicines for HIV infection) ■ any other prescription drugs. Sodium bicarbonate may interact with certain prescription drugs. Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours 14-Day Course of Treatment ■ swallow 1 capsule with a glass of water at least 1 hour before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ do not chew or crush the capsule ■ do not open capsule and sprinkle on food ■ do not use for more than 14 days unless directed by your doctor Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor	

03658505



ATTACH LEGEND ON ALL
DIGITAL MECHANICALS

NO INK
NO VARNISH

STANDARD FORMAT

Helvetica Condensed

Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic

Drug Facts (continued)

Drug Facts - 8 pt Helvetica Cnd. Bold Italic

(continued) - 8 pt Helvetica Cnd.

Headings - 8 pt Helvetica Cnd. Bold Italic

Sub Heads - 6 pt Helvetica Cnd. Bold

Text - 6 pt Helvetica Cnd. minimum leading 6.5

Square Bullets - 3.5 pt Zapf Dingbats no compression

Drug Facts (continued) ■ children under 18 years of age, ask a doctor. ■ heartburn in children may sometimes be caused by a serious condition.	
Other Information ■ each capsule contains: sodium 303 mg. ■ read the directions, warnings and accompanying label information before use ■ store at 20°-25°C (68°-77°F) ■ temperature: do not use if the blue band around the capsule is missing or broken. Do not use if foil inner seal imprinted with "Sealed for your protection" is missing, open or broken. ■ keep product out of high heat and humidity ■ protect product from moisture	
Inactive Ingredients croscarmellose sodium, D1C blue ink, 1, gelatin, magnesium stearate, pharmaceutical ink, polystyrene 80, titanium dioxide	
Questions or comments? Call 1-888-4-ZEG-OTC (1-888-483-4882) between 8:00 AM and 5:00 PM Central Time, Monday through Friday or visit www.ZegeridOTC.com	

recyclable
carton

© Copyright & Distributed by MSD Consumer Care, Inc.,
a subsidiary of Merck & Co., Inc.,
Whitehouse Station, NJ USA.
All rights reserved. Product of Spain.
36585-05/PK601278

Treats Frequent Heartburn

Zegerid[®]

OTC[®]



42 CAPSULES

MDC 11523 7265 3

Treats Frequent Heartburn

Zegerid[®]

OTC[®]

42 CAPSULES

Three 14-Day Courses of Treatment

Treats Frequent Heartburn

Zegerid[®]

OTC[®]

42 CAPSULES

• Omeprazole 20 mg/Acid Reducer
• Sodium Bicarbonate 1100 mg/
Allows Absorption of this Omeprazole Product




Lot/EXP
Area
No Varnish
No Ink

3.5434 [90.00]



A - Top Panel




B - Back Panel




Printable Text Area




Base Panel




C - Insert 1 Top




D - Insert 1 Back




E - Insert 2 Top




F - Insert 2 Back




G - Insert 3 Top



H - Insert 3 Back




I - Insert 4 Top




J - Insert 4 Back




K - Insert 5 Top



L - Insert 5 Back



M - Insert 6 Top



N - Insert 6 Back

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic

Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd. Bold Italic (continued) - 8 pt Helvetica Cnd.

Headings - 8 pt Helvetica Cnd. Bold Italic

Sub Heads - 6 pt Helvetica Cnd. Bold

Text - 6 pt Helvetica Cnd, minimum leading 6.5

Square Bullets - 3.5 pt Zaph Dingbats no compression

Border - 1 pt rule, 3/64" gap all around

Barline - 1 pt rule

Hairline - .5 pt rule

Treats Frequent Heartburn

Zegerid[®] OTC[®]

- Omeprazole 20 mg/Acid Reducer
- Sodium Bicarbonate 1100 mg/
Allows Absorption of this Omeprazole Product

Drug Facts Active ingredients (in each capsule) Omeprazole 20 mg..... Sodium Bicarbonate 1100 mg.....Allows absorption of this omeprazole product Purpose Acid reducer Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect. Warnings Allergy alert: Do not use if you are allergic to omeprazole Do not use if you have: ■ trouble or pain swallowing food ■ vomiting with blood ■ bloody or black stools These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ a sodium-restricted diet Ask a doctor or pharmacist before use if you are taking ■ warfarin, clopidogrel or cilostazol (blood-thinning medicines) ■ prescription antifungal or anti-yeast medicines ■ diazepam (anxiety medicine) ■ digoxin (heart medicine) ■ tacrolimus (immune system medicine) ■ prescription antiretrovirals (medicines for HIV infection) ■ any other prescription drugs. Sodium bicarbonate may interact with certain prescription drugs.	Drug Facts (continued) Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Directions ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours 14-Day Course of Treatment ■ swallow 1 capsule with a glass of water at least 1 hour before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ do not chew or crush the capsule ■ do not open capsule and sprinkle on food ■ do not use for more than 14 days unless directed by your doctor Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months	Drug Facts (continued) ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. Other information ■ each capsule contains: sodium 303 mg ■ read the directions, warnings and accompanying label information before use ■ store at 20°-25°C (68°-77°F) ■ tamper-evident: do not use if the blue band around the capsule is missing or broken. Do not use if foil inner seal imprinted with "Sealed for your protection" is missing, open or broken. ■ keep product out of high heat and humidity ■ protect product from moisture Inactive ingredients croscarmellose sodium, FD&C blue No. 1, gelatin, magnesium stearate, pharmaceutical ink, polysorbate 80, titanium dioxide Questions or comments? Call 1-888-4-ZEG-OTC (1-888-493-4582) between 8:00 AM and 5:00 PM Central Time, Monday through Friday or visit www.ZegeridOTC.com
--	--	---

© Copyright & Distributed by MSD Consumer Care, Inc., PO Box 377, Memphis, TN 38151 USA, a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA. All rights reserved. Product of Spain.

NO INK NO VARNISH



recyclable carton

42 CAPSULES
Three 14-Day Courses of Treatment

Zegerid[®] OTC[®]

Treats Frequent Heartburn

- Omeprazole 20 mg/Acid Reducer
- Sodium Bicarbonate 1100 mg/
Allows Absorption of this Omeprazole Product

ITM# 807175

NDC 11529-7265-3

3 Pack

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER
07/25/2011