

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use BOTOX® safely and effectively. See full prescribing information for BOTOX.

### BOTOX (onabotulinumtoxinA)

Initial U.S. Approval: 1989

#### WARNING: Distant Spread of Toxin Effect

See full prescribing information for complete boxed warning. The effects of BOTOX and all botulinum toxin products may spread from the area of injection to produce symptoms consistent with botulinum toxin effects. These symptoms have been reported hours to weeks after injection. Swallowing and breathing difficulties can be life threatening and there have been reports of death. The risk of symptoms is probably greatest in children treated for spasticity but symptoms can also occur in adults, particularly in those patients who have underlying conditions that would predispose them to these symptoms.

#### RECENT MAJOR CHANGES

- Indications and Usage, Chronic Migraine (1.1) 10/2010
- Dosage and Administration, Chronic Migraine (2.2) 10/2010

#### INDICATIONS AND USAGE

BOTOX is an acetylcholine release inhibitor and a neuromuscular blocking agent indicated for:

- Prophylaxis of headaches in adult patients with chronic migraine ( $\geq 15$  days per month with headache lasting 4 hours a day or longer) (1.1)
- Treatment of upper limb spasticity in adult patients (1.2)
- Treatment of cervical dystonia in adult patients, to reduce the severity of abnormal head position and neck pain (1.3)
- Treatment of severe axillary hyperhidrosis that is inadequately managed by topical agents in adult patients (1.4)
- Treatment of blepharospasm associated with dystonia in patients  $\geq 12$  years of age (1.5)
- Treatment of strabismus in patients  $\geq 12$  years of age (1.5)

#### Important limitations:

- Safety and effectiveness have not been established for the prophylaxis of episodic migraine (14 headache days or fewer per month).
- Safety and effectiveness of BOTOX have not been established for the treatment of upper limb spasticity in pediatric patients, and for the treatment of lower limb spasticity in adult and pediatric patients.
- Safety and effectiveness of BOTOX for hyperhidrosis in body areas other than axillary have not been established.

#### DOSAGE AND ADMINISTRATION

- Indication specific dosage and administration recommendations should be followed; Do not exceed a total dose of 360 Units administered every 12 to 16 weeks or at longer intervals (2)
- See Preparation and Dilution Technique for instructions on BOTOX reconstitution, storage, and preparation before injection (2.1)
- Chronic Migraine: Recommended total dose 155 Units, as 0.1 mL (5 Units) injections per each site divided across 7 head/neck muscles (2.2)
- Upper Limb Spasticity: Select dose based on muscles affected, severity of muscle activity, prior response to treatment, and adverse event history; Electromyographic guidance recommended (2.3)
- Cervical Dystonia: Base dosing on the patient's head and neck position, localization of pain, muscle hypertrophy, patient response, and adverse event history; use lower initial dose in botulinum toxin naïve patients (2.4)
- Axillary Hyperhidrosis: 50 Units per axilla (2.5)

- Blepharospasm: 1.25 Units-2.5 Units into each of 3 sites per affected eye (2.6)
- Strabismus: 1.25 Units-2.5 Units initially in any one muscle (2.7)

#### DOSAGE FORMS AND STRENGTHS

Single-use, sterile 50 Units, 100 Units, or 200 Units vacuum-dried powder for reconstitution only with sterile, non-preserved 0.9% Sodium Chloride Injection USP prior to injection (3)

#### CONTRAINDICATIONS

- Hypersensitivity to any botulinum toxin preparation or to any of the components in the formulation (4.1, 5.3, 6.2)
- Infection at the proposed injection site (4.2)

#### WARNINGS AND PRECAUTIONS

- Potency Units of BOTOX not interchangeable with other preparations of botulinum toxin products (5.1, 11)
- Spread of toxin effects; swallowing and breathing difficulties can lead to death (5.2)
- Immediate medical attention may be required in cases of respiratory, speech or swallowing difficulties (5.2, 5.4)
- Concomitant neuromuscular disorder may exacerbate clinical effects of treatment (5.5)
- Use with caution in patients with compromised respiratory function (5.4, 5.6)
- Corneal exposure and ulceration (5.7)
- Retrobulbar hemorrhages and compromised retinal circulation (5.8)
- Bronchitis and upper respiratory tract infections in patients treated for upper limb spasticity (5.9)

#### ADVERSE REACTIONS

In controlled studies, the most commonly observed adverse reactions ( $\geq 5\%$  and  $>$ placebo) were:

- Chronic Migraine: neck pain, headache (6.1)
- Spasticity: pain in extremity (6.1)
- Cervical Dystonia: dysphagia, upper respiratory infection, neck pain, headache, increased cough, flu syndrome, back pain, rhinitis (6.1)
- Axillary Hyperhidrosis: injection site pain and hemorrhage, non-axillary sweating, pharyngitis, flu syndrome (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Allergan at 1-800-433-8871 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

#### DRUG INTERACTIONS

- Patients receiving concomitant treatment of BOTOX and aminoglycosides or other agents interfering with neuromuscular transmission (e.g., curare-like agents), or muscle relaxants, should be observed closely because the effect of BOTOX may be potentiated (7)

#### USE IN SPECIFIC POPULATIONS

- Pregnancy: Based on animal data, may cause fetal harm (8.1)
- Pediatric Use: Safety and efficacy are not established in patients under 18 years of age for the prophylaxis of headaches in chronic migraine, the treatment of upper limb spasticity and axillary hyperhidrosis, in patients under 16 years of age for the treatment of cervical dystonia, and in patients under 12 years of age for the treatment of blepharospasm and strabismus (8.4)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

Revised: XX/2011

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

- 1.1 Chronic Migraine
- 1.2 Upper Limb Spasticity
- 1.3 Cervical Dystonia
- 1.4 Primary Axillary Hyperhidrosis
- 1.5 Blepharospasm and Strabismus

### 2 DOSAGE AND ADMINISTRATION

- 2.1 Preparation and Dilution Technique
- 2.2 Chronic Migraine
- 2.3 Upper Limb Spasticity
- 2.4 Cervical Dystonia
- 2.5 Primary Axillary Hyperhidrosis
- 2.6 Blepharospasm
- 2.7 Strabismus

### 3 DOSAGE FORMS AND STRENGTHS

### 4 CONTRAINDICATIONS

- 4.1 Known Hypersensitivity to Botulinum Toxin
- 4.2 Infection at the Injection Site(s)

### 5 WARNINGS AND PRECAUTIONS

- 5.1 Lack of Interchangeability between Botulinum Toxin Products
- 5.2 Spread of Toxin Effect
- 5.3 Hypersensitivity Reactions
- 5.4 Dysphagia and Breathing Difficulties in Treatment of Cervical Dystonia
- 5.5 Pre-Existing Neuromuscular Disorders
- 5.6 Pulmonary Effects of **BOTOX** in Patients with Compromised Respiratory Status Treated for Spasticity
- 5.7 Corneal Exposure and Ulceration in Patients Treated with **BOTOX** for Blepharospasm
- 5.8 Retrobulbar Hemorrhages in Patients Treated with **BOTOX** for Strabismus
- 5.9 Bronchitis and Upper Respiratory Tract Infections in Patients Treated for Spasticity

### 5.10 Human Albumin and Transmission of Viral Diseases

### 6 ADVERSE REACTIONS

- 6.1 Clinical Studies Experience
- 6.2 Post-Marketing Experience
- 6.3 Immunogenicity

### 7 DRUG INTERACTIONS

### 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

### 10 OVERDOSAGE

### 11 DESCRIPTION

### 12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

### 13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### 14 CLINICAL STUDIES

- 14.1 Chronic Migraine
- 14.2 Upper Limb Spasticity
- 14.3 Cervical Dystonia
- 14.4 Primary Axillary Hyperhidrosis
- 14.5 Blepharospasm
- 14.6 Strabismus

### 16 HOW SUPPLIED/STORAGE AND HANDLING

### 17 PATIENT COUNSELING INFORMATION

- 17.1 Swallowing, Speaking or Breathing Difficulties, or Other Unusual Symptoms
- 17.2 Ability to Operate Machinery or Vehicles
- 17.3 Medication Guide

\* Sections or subsections omitted from the full prescribing information are not listed

## FULL PRESCRIBING INFORMATION

### Distant Spread of Toxin Effect

Postmarketing reports indicate that the effects of **BOTOX** and all botulinum toxin products may spread from the area of injection to produce symptoms consistent with botulinum toxin effects. These may include asthenia, generalized muscle weakness, diplopia, ptosis, dysphagia, dysphonia, dysarthria, urinary incontinence and breathing difficulties. These symptoms have been reported hours to weeks after injection. Swallowing and breathing difficulties can be life threatening and there have been reports of death. The risk of symptoms is probably greatest in children treated for spasticity but symptoms can also occur in adults treated for spasticity and other conditions, particularly in those patients who have underlying conditions that would predispose them to these symptoms. In unapproved uses, including spasticity in children, and in approved indications, cases of spread of effect have been reported at doses comparable to those used to treat cervical dystonia and at lower doses.

### 1 INDICATIONS AND USAGE

#### 1.1 Chronic Migraine

**BOTOX** (onabotulinumtoxinA) for injection is indicated for the prophylaxis of headaches in adult patients with chronic migraine ( $\geq 15$  days per month with headache lasting 4 hours a day or longer).

#### *Important limitations*

Safety and effectiveness have not been established for the prophylaxis of episodic migraine (14 headache days or fewer per month) in seven placebo-controlled studies.

#### 1.2 Upper Limb Spasticity

**BOTOX** is indicated for the treatment of upper limb spasticity in adult patients, to decrease the severity of increased muscle tone in elbow flexors (biceps), wrist flexors (flexor carpi radialis and flexor carpi ulnaris) and finger flexors (flexor digitorum profundus and flexor digitorum sublimis).

#### *Important limitations*

Safety and effectiveness of **BOTOX** have not been established for the treatment of other upper limb muscle groups, or for the treatment of lower limb spasticity. Safety and effectiveness of **BOTOX** have not been established for the treatment of spasticity in

pediatric patients under age 18 years. **BOTOX** has not been shown to improve upper extremity functional abilities, or range of motion at a joint affected by a fixed contracture. Treatment with **BOTOX** is not intended to substitute for usual standard of care rehabilitation regimens.

### **1.3 Cervical Dystonia**

**BOTOX** is indicated for the treatment of adults with cervical dystonia, to reduce the severity of abnormal head position and neck pain associated with cervical dystonia.

### **1.4 Primary Axillary Hyperhidrosis**

**BOTOX** is indicated for the treatment of severe primary axillary hyperhidrosis that is inadequately managed with topical agents.

#### *Important limitations*

The safety and effectiveness of **BOTOX** for hyperhidrosis in other body areas have not been established. Weakness of hand muscles and blepharoptosis may occur in patients who receive **BOTOX** for palmar hyperhidrosis and facial hyperhidrosis, respectively. Patients should be evaluated for potential causes of secondary hyperhidrosis (e.g., hyperthyroidism) to avoid symptomatic treatment of hyperhidrosis without the diagnosis and/or treatment of the underlying disease.

Safety and effectiveness of **BOTOX** have not been established for the treatment of axillary hyperhidrosis in pediatric patients under age 18.

### **1.5 Blepharospasm and Strabismus**

**BOTOX** is indicated for the treatment of strabismus and blepharospasm associated with dystonia, including benign essential blepharospasm or VII nerve disorders in patients 12 years of age and above.

## **2 DOSAGE AND ADMINISTRATION**

The potency Units of **BOTOX** (onabotulinumtoxinA) for injection are specific to the preparation and assay method utilized. They are not interchangeable with other preparations of botulinum toxin products and, therefore, units of biological activity of **BOTOX** cannot be compared to nor converted into units of any other botulinum toxin products assessed with any other specific assay method [*see Warnings and Precautions (5.1) and Description (11)*].

Injection specific dosage and administration recommendations should be followed. In treating adult patients for one or more indications, the maximum cumulative dose should generally not exceed 360 Units, in a 3 month interval.

The safe and effective use of **BOTOX** depends upon proper storage of the product, selection of the correct dose, and proper reconstitution and administration techniques. Physicians administering **BOTOX** must understand the relevant neuromuscular and/or orbital anatomy of the area involved and any alterations to the anatomy due to prior surgical procedures. An understanding of standard electromyographic techniques is also required for treatment of strabismus and of upper limb spasticity, and may be useful for the treatment of cervical dystonia.

Use caution when **BOTOX** treatment is used in the presence of inflammation at the proposed injection site(s) or when excessive weakness or atrophy is present in the target muscle(s).

### **2.1 Preparation and Dilution Technique**

**BOTOX** is supplied in single-use 50 Units, 100 Units, and 200 Units per vial. Prior to injection, reconstitute each vacuum-dried vial of **BOTOX** with sterile, non-preserved 0.9% Sodium Chloride Injection USP. Draw up the proper amount of diluent in the appropriate size syringe ([Dilution Table](#)), and slowly inject the diluent into the vial. Discard the vial if a vacuum does not pull the diluent into the vial. Gently mix **BOTOX** with the saline by rotating the vial. Record the date and time of reconstitution on the space on the label. **BOTOX** should be administered within 24 hours after reconstitution. During this time period, reconstituted **BOTOX** should be stored in a refrigerator (2° to 8°C).

**Dilution Instructions for BOTOX Vials (50 Units, 100 Units, and 200 Units)**

Diluent* Added to 50 Unit Vial	Resulting Dose Units per 0.1 mL	Diluent* Added to 100 Unit Vial	Resulting Dose Units per 0.1 mL	Diluent* Added to 200 Unit Vial	Resulting Dose Units per 0.1 mL
1 mL	5 Units	1 mL	10 Units	1 mL	20 Units
2 mL	2.5 Units	2 mL	5 Units	2 mL	10 Units
4 mL	1.25 Units	4 mL	2.5 Units	4 mL	5 Units
		8 mL	1.25 Units	8 mL	2.5 Units
				10 mL	2 Units

\*Preservative-free 0.9% Sodium Chloride Injection, USP Only

Note: These dilutions are calculated for an injection volume of 0.1 mL. A decrease or increase in the **BOTOX** dose is also possible by administering a smaller or larger injection volume - from 0.05 mL (50% decrease in dose) to 0.15 mL (50% increase in dose).





An injection of **BOTOX** is prepared by drawing into an appropriately sized sterile syringe an amount of the properly reconstituted toxin slightly greater than the intended dose. Air bubbles in the syringe barrel are expelled and the syringe is attached to an appropriate injection needle. Patency of the needle should be confirmed. A new, sterile needle and syringe should be used to enter the vial on each occasion for removal of **BOTOX**.

Reconstituted **BOTOX** should be clear, colorless, and free of particulate matter. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration and whenever the solution and the container permit.

**2.2 Chronic Migraine**

The recommended dilution is 200 Units/4 mL or 100 Units/2 mL, with a final concentration of 5 Units per 0.1 mL (see Dilution Table). The recommended dose for treating chronic migraine is 155 Units administered intramuscularly (IM) using a sterile 30-gauge, 0.5 inch needle as 0.1 mL (5 Units) injections per each site. Injections should be divided across 7 specific head/neck muscle areas as specified in the diagrams and Table 1 below. A 1 inch needle may be needed in the neck region for patients with thick neck muscles. With the exception of the procerus muscle, which should be injected at 1 site (midline), all muscles should be injected bilaterally with half the number of injection sites administered to the left, and half to the right side of the head and neck. The recommended re-treatment schedule is every 12 weeks.

Recommended injection sites for chronic migraine:



- A. Corrugator: 5 U each side
- B. Procerus: 5 U [one site]
- C. Frontalis: 10 U each side

D. Temporalis: 20 U each side

E. Occipitalis: 15 U each side

F. Cervical paraspinal: 10 U each side

G. Trapezius: 15 U each side

**Table 1: BOTOX Dosing by Muscle for Chronic Migraine**

Head/Neck Area	Recommended Dose (Number of Sites <sup>a</sup> )
Frontalis <sup>b</sup>	20 Units divided in 4 sites
Corrugator <sup>b</sup>	10 Units divided in 2 sites
Procerus	5 Units in 1 site
Occipitalis <sup>b</sup>	30 Units divided in 6 sites
Temporalis <sup>b</sup>	40 Units divided in 8 sites
Trapezius <sup>b</sup>	30 Units divided in 6 sites
Cervical Paraspinal Muscle Group <sup>b</sup>	20 Units divided in 4 sites
<b>Total Dose:</b>	<b>155 Units divided in 31 sites</b>

<sup>a</sup> Each IM injection site = 0.1 mL = 5 Units BOTOX

<sup>b</sup> Dose distributed bilaterally

### 2.3 Upper Limb Spasticity

Dosing in initial and sequential treatment sessions should be tailored to the individual based on the size, number and location of muscles involved, severity of spasticity, the presence of local muscle weakness, the patient's response to previous treatment, or adverse event history with **BOTOX**. In clinical trials, doses ranging from 75 Units to 360 Units were divided among selected muscles at a given treatment session.

**Table 2: BOTOX Dosing by Muscle for Upper Limb Spasticity**

Muscle	Recommended Dose Total Dosage (Number of Sites)
Biceps Brachii	100 Units-200 Units divided in 4 sites
Flexor Carpi Radialis	12.5 Units-50 Units in 1 site
Flexor Carpi Ulnaris	12.5 Units-50 Units in 1 site
Flexor Digitorum Profundus	30 Units-50 Units in 1 site
Flexor Digitorum Sublimis	30 Units-50 Units in 1 site

The recommended dilution is 200 Units/4 mL or 100 Units/2 mL with 0.9% non-preserved sterile saline (see Dilution Table). The lowest recommended starting dose should be used, and no more than 50 Units per site should generally be administered. An appropriately sized needle (e.g., 25-30 gauge) may be used for superficial muscles, and a longer 22 gauge needle may be used for deeper musculature. Localization of the involved muscles with electromyographic guidance or nerve stimulation techniques is recommended.

Repeat **BOTOX** treatment may be administered when the effect of a previous injection has diminished, but generally no sooner than 12 weeks after the previous injection. The degree and pattern of muscle spasticity at the time of re-injection may necessitate alterations in the dose of **BOTOX** and muscles to be injected.

### 2.4 Cervical Dystonia

The phase 3 study enrolled patients who had extended histories of receiving and tolerating **BOTOX** injections, with prior individualized adjustment of dose. The mean **BOTOX** dose administered to patients in the phase 3 study was 236 Units (25th to 75th percentile range of 198 Units to 300 Units). The **BOTOX** dose was divided among the affected muscles [see *Clinical Studies (14.3)*]. Dosing in initial and sequential treatment sessions should be tailored to the individual patient based on the patient's head and neck position, localization of pain, muscle hypertrophy, patient response, and adverse event history. The initial dose for a patient without prior use of **BOTOX** should be at a lower dose, with subsequent dosing adjusted based on individual response. Limiting the total dose injected into the sternocleidomastoid muscle to 100 Units or less may decrease the occurrence of dysphagia [see *Warnings and Precautions (5.2, 5.4, 5.5)*].

The recommended dilution is 200 Units/2 mL, 200 Units/4 mL, 100 Units/1 mL, or 100 Units/2 mL with 0.9% non-preserved sterile saline, depending on volume and number of injection sites desired to achieve treatment objectives (see Dilution Table). In general, no more than 50 Units per site should be administered. An appropriately sized needle (e.g., 25-30 gauge) may be used for superficial muscles, and a longer 22 gauge needle may be used for deeper musculature. Localization of the involved muscles with electromyographic guidance may be useful.

Clinical improvement generally begins within the first two weeks after injection with maximum clinical benefit at approximately six weeks post-injection. In the phase 3 study most subjects were observed to have returned to pre-treatment status by 3 months post-treatment.

## 2.5 Primary Axillary Hyperhidrosis

The recommended dose is 50 Units per axilla. The hyperhidrotic area to be injected should be defined using standard staining techniques, e.g., Minor's Iodine-Starch Test. The recommended dilution is 100 Units/4 mL with 0.9% preservative-free sterile saline (see [Dilution Table](#)). Using a 30 gauge needle, 50 Units of **BOTOX** (2 mL) is injected intradermally in 0.1 to 0.2 mL aliquots to each axilla evenly distributed in multiple sites (10-15) approximately 1-2 cm apart.


Repeat injections for hyperhidrosis should be administered when the clinical effect of a previous injection diminishes.

### *Instructions for the Minor's Iodine-Starch Test Procedure:*

Patients should shave underarms and abstain from use of over-the-counter deodorants or antiperspirants for 24 hours prior to the test. Patient should be resting comfortably without exercise, hot drinks, etc. for approximately 30 minutes prior to the test. Dry the underarm area and then immediately paint it with iodine solution. Allow the area to dry, then lightly sprinkle the area with starch powder. Gently blow off any excess starch powder. The hyperhidrotic area will develop a deep blue-black color over approximately 10 minutes.

Each injection site has a ring of effect of up to approximately 2 cm in diameter. To minimize the area of no effect, the injection sites should be evenly spaced as shown in Figure 1:

**Figure 1:**



Each dose is injected to a depth of approximately 2 mm and at a 45° angle to the skin surface, with the bevel side up to minimize leakage and to ensure the injections remain intradermal. If injection sites are marked in ink, do not inject **BOTOX** directly through the ink mark to avoid a permanent tattoo effect.

## 2.6 Blepharospasm

For blepharospasm, reconstituted **BOTOX** is injected using a sterile, 27-30 gauge needle without electromyographic guidance. The initial recommended dose is 1.25 Units-2.5 Units (0.05 mL to 0.1 mL volume at each site) injected into the medial and lateral pre-tarsal orbicularis oculi of the upper lid and into the lateral pre-tarsal orbicularis oculi of the lower lid. Avoiding injection near the levator palpebrae superioris may reduce the complication of ptosis. Avoiding medial lower lid injections, and thereby reducing diffusion into the inferior oblique, may reduce the complication of diplopia. Ecchymosis occurs easily in the soft eyelid tissues. This can be prevented by applying pressure at the injection site immediately after the injection.

The recommended dilution to achieve 1.25 Units is 50 Units/4 mL or 100 Units/8 mL; for 2.5 Units it is 50 Units/2 mL or 100 Units/4 mL (see [Dilution Table](#)).

In general, the initial effect of the injections is seen within three days and reaches a peak at one to two weeks post-treatment. Each treatment lasts approximately three months, following which the procedure can be repeated. At repeat treatment sessions, the dose may be increased up to two-fold if the response from the initial treatment is considered insufficient, usually defined as an effect that does not last longer than two months. However, there appears to be little benefit obtainable from injecting more than 5 Units per site. Some tolerance may be found when **BOTOX** is used in treating blepharospasm if treatments are given any more frequently than every three months, and is rare to have the effect be permanent.

The cumulative dose of **BOTOX** treatment for blepharospasm in a 30-day period should not exceed 200 Units.

## 2.7 Strabismus

**BOTOX** is intended for injection into extraocular muscles utilizing the electrical activity recorded from the tip of the injection needle as a guide to placement within the target muscle. Injection without surgical exposure or electromyographic guidance should not be attempted. Physicians should be familiar with electromyographic technique.

To prepare the eye for **BOTOX** injection, it is recommended that several drops of a local anesthetic and an ocular decongestant be given several minutes prior to injection.

Note: The volume of **BOTOX** injected for treatment of strabismus should be between 0.05-0.15 mL per muscle.

The initial listed doses of the reconstituted **BOTOX** [see *Dosage and Administration (2.1)*] typically create paralysis of the injected muscles beginning one to two days after injection and increasing in intensity during the first week. The paralysis lasts for 2-6 weeks and gradually resolves over a similar time period. Overcorrections lasting over six months have been rare. About one half of patients will require subsequent doses because of inadequate paralytic response of the muscle to the initial dose, or because of mechanical factors such as large deviations or restrictions, or because of the lack of binocular motor fusion to stabilize the alignment.

- I. Initial doses in Units. Use the lower listed doses for treatment of small deviations. Use the larger doses only for large deviations.
  - A. For vertical muscles, and for horizontal strabismus of less than 20 prism diopters: 1.25 Units-2.5 Units in any one muscle.
  - B. For horizontal strabismus of 20 prism diopters to 50 prism diopters: 2.5 Units-5 Units in any one muscle.
  - C. For persistent VI nerve palsy of one month or longer duration: 1.25 Units-2.5 Units in the medial rectus muscle.
- II. Subsequent doses for residual or recurrent strabismus.
  - A. It is recommended that patients be re-examined 7-14 days after each injection to assess the effect of that dose.
  - B. Patients experiencing adequate paralysis of the target muscle that require subsequent injections should receive a dose comparable to the initial dose.
  - C. Subsequent doses for patients experiencing incomplete paralysis of the target muscle may be increased up to two-fold compared to the previously administered dose.
  - D. Subsequent injections should not be administered until the effects of the previous dose have dissipated as evidenced by substantial function in the injected and adjacent muscles.
  - E. The maximum recommended dose as a single injection for any one muscle is 25 Units.

The recommended dilution to achieve 1.25 Units is 50 Units/4 mL or 100 Units/8 mL; for 2.5 Units it is 50 Units/2 mL or 100 Units/4 mL (see Dilution Table).

### 3 DOSAGE FORMS AND STRENGTHS

Single-use, sterile 50 Units, 100 Units, or 200 Units vacuum-dried powder for reconstitution only with sterile, non-preserved 0.9% Sodium Chloride Injection USP prior to injection [see *Dosage and Administration (2.1)*].

### 4 CONTRAINDICATIONS

#### 4.1 Known Hypersensitivity to Botulinum Toxin

**BOTOX** is contraindicated in patients who are hypersensitive to any botulinum toxin preparation or to any of the components in the formulation [see *Warnings and Precautions (5.3)*].

#### 4.2 Infection at the Injection Site(s)

**BOTOX** is contraindicated in the presence of infection at the proposed injection site(s).

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Lack of Interchangeability between Botulinum Toxin Products

**The potency Units of BOTOX are specific to the preparation and assay method utilized. They are not interchangeable with other preparations of botulinum toxin products and, therefore, units of biological activity of BOTOX cannot be compared to nor converted into units of any other botulinum toxin products assessed with any other specific assay method [see Description (11)].**

#### 5.2 Spread of Toxin Effect

Postmarketing safety data from **BOTOX** and other approved botulinum toxins suggest that botulinum toxin effects may, in some cases, be observed beyond the site of local injection. The symptoms are consistent with the mechanism of action of botulinum toxin and may include asthenia, generalized muscle weakness, diplopia, ptosis, dysphagia, dysphonia, dysarthria, urinary incontinence, and breathing difficulties. These symptoms have been reported hours to weeks after injection. Swallowing and breathing difficulties can be life threatening and there have been reports of death related to spread of toxin effects. The risk of the symptoms is probably greatest in children treated for spasticity but symptoms can also occur in adults treated for spasticity and other conditions, and particularly in those patients who have underlying conditions that would predispose them to these symptoms. In unapproved uses, including spasticity in children, and in approved indications, symptoms consistent with spread of toxin effect have been reported at doses comparable to or lower than doses used to treat cervical dystonia.

No definitive serious adverse event reports of distant spread of toxin effect associated with dermatologic use of **BOTOX/BOTOX Cosmetic** at the labeled dose of 20 Units (for glabellar lines) or 100 Units (for severe primary axillary hyperhidrosis) have been reported.

No definitive serious adverse event reports of distant spread of toxin effect associated with **BOTOX** for blepharospasm at the recommended dose (30 Units and below), strabismus, or for chronic migraine at the labeled doses have been reported.

### 5.3 Hypersensitivity Reactions

Serious and/or immediate hypersensitivity reactions have been reported. These reactions include anaphylaxis, serum sickness, urticaria, soft tissue edema, and dyspnea. If such a reaction occurs, further injection of **BOTOX** should be discontinued and appropriate medical therapy immediately instituted. One fatal case of anaphylaxis has been reported in which lidocaine was used as the diluent, and consequently the causal agent cannot be reliably determined.

### 5.4 Dysphagia and Breathing Difficulties in Treatment of Cervical Dystonia

Treatment with **BOTOX** and other botulinum toxin products can result in swallowing or breathing difficulties. Patients with pre-existing swallowing or breathing difficulties may be more susceptible to these complications. In most cases, this is a consequence of weakening of muscles in the area of injection that are involved in breathing or swallowing. When distant effects occur, additional respiratory muscles may be involved [see *Warnings and Precautions (5.2)*].

Deaths as a complication of severe dysphagia have been reported after treatment with botulinum toxin. Dysphagia may persist for several months, and require use of a feeding tube to maintain adequate nutrition and hydration. Aspiration may result from severe dysphagia and is a particular risk when treating patients in whom swallowing or respiratory function is already compromised.

Treatment of cervical dystonia with botulinum toxins may weaken neck muscles that serve as accessory muscles of ventilation. This may result in a critical loss of breathing capacity in patients with respiratory disorders who may have become dependent upon these accessory muscles. There have been postmarketing reports of serious breathing difficulties, including respiratory failure, in cervical dystonia patients.

Patients with smaller neck muscle mass and patients who require bilateral injections into the sternocleidomastoid muscle have been reported to be at greater risk for dysphagia. Limiting the dose injected into the sternocleidomastoid muscle may reduce the occurrence of dysphagia. Injections into the levator scapulae may be associated with an increased risk of upper respiratory infection and dysphagia.

Patients treated with botulinum toxin may require immediate medical attention should they develop problems with swallowing, speech or respiratory disorders. These reactions can occur within hours to weeks after injection with botulinum toxin [see *Warnings and Precautions (5.2) and Adverse Reactions (6.1)*].

### 5.5 Pre-Existing Neuromuscular Disorders

Individuals with peripheral motor neuropathic diseases, amyotrophic lateral sclerosis or neuromuscular junction disorders (e.g., myasthenia gravis or Lambert-Eaton syndrome) should be monitored particularly closely when given botulinum toxin. Patients with neuromuscular disorders may be at increased risk of clinically significant effects including severe dysphagia and respiratory compromise from typical doses of **BOTOX** [see *Adverse Reactions (6.1)*].

### 5.6 Pulmonary Effects of BOTOX in Patients with Compromised Respiratory Status Treated for Spasticity

Patients with compromised respiratory status treated with **BOTOX** for upper limb spasticity should be monitored closely. In a double-blind, placebo-controlled, parallel group study in patients with stable reduced pulmonary function (defined as FEV<sub>1</sub> 40-80% of predicted value and FEV<sub>1</sub>/FVC ≤ 0.75), the event rate in change of Forced Vital Capacity ≥15% or ≥20% was generally greater in patients treated with **BOTOX** than in patients treated with placebo (see Table 3).

**Table 3: Event rate per patient treatment cycle among patients with reduced lung function who experienced at least a 15% or 20% decrease in forced vital capacity from baseline at Week 1, 6, 12 post-injection with up to two treatment cycles with BOTOX or placebo**

	BOTOX 360 Units		BOTOX 240 Units		Placebo	
	≥15%	≥20%	≥15%	≥20%	≥15%	≥20%
Week 1	4%	0%	3%	0%	7%	3%
Week 6	7%	4%	4%	2%	2%	2%
Week 12	10%	5%	2%	1%	4%	1%

Differences from placebo were not statistically significant

In patients with reduced lung function, upper respiratory tract infections were also reported more frequently as adverse reactions in patients treated with **BOTOX** [see *Warnings and Precautions (5.9)*].

### 5.7 Corneal Exposure and Ulceration in Patients Treated with BOTOX for Blepharospasm

Reduced blinking from **BOTOX** injection of the orbicularis muscle can lead to corneal exposure, persistent epithelial defect, and corneal ulceration, especially in patients with VII nerve disorders. Vigorous treatment of any epithelial defect should be employed. This may require protective drops, ointment, therapeutic soft contact lenses, or closure of the eye by patching or other means.

### 5.8 Retrobulbar Hemorrhages in Patients Treated with BOTOX for Strabismus

During the administration of **BOTOX** for the treatment of strabismus, retrobulbar hemorrhages sufficient to compromise retinal circulation have occurred. It is recommended that appropriate instruments to decompress the orbit be accessible.

### 5.9 Bronchitis and Upper Respiratory Tract Infections in Patients Treated for Spasticity

Bronchitis was reported more frequently as an adverse reaction in patients treated for upper limb spasticity with **BOTOX** (3% at 251 Units-360 Units total dose), compared to placebo (1%). In patients with reduced lung function treated for upper limb spasticity, upper respiratory tract infections were also reported more frequently as adverse reactions in patients treated with **BOTOX** (11% at 360 Units total dose; 8% at 240 Units total dose) compared to placebo (6%).

### 5.10 Human Albumin and Transmission of Viral Diseases

This product contains albumin, a derivative of human blood. Based on effective donor screening and product manufacturing processes, it carries an extremely remote risk for transmission of viral diseases. A theoretical risk for transmission of Creutzfeldt-Jakob disease (CJD) is also considered extremely remote. No cases of transmission of viral diseases or CJD have ever been reported for albumin.

## 6 ADVERSE REACTIONS

The following adverse reactions to **BOTOX** (onabotulinumtoxinA) for injection are discussed in greater detail in other sections of the labeling:

- Spread of Toxin Effects [*see Warnings and Precautions (5.2)*]
- Hypersensitivity [*see Contraindications (4.1) and Warnings and Precautions (5.3)*]
- Dysphagia and Breathing Difficulties in Treatment of Cervical Dystonia [*see Warnings and Precautions (5.4)*]
- Bronchitis and Upper Respiratory Tract Infections in Patients Treated for Spasticity [*see Warnings and Precautions (5.9)*]

### 6.1 Clinical Studies Experience

Because clinical trials are conducted under widely varying conditions, the adverse reaction rates observed cannot be directly compared to rates in other trials and may not reflect the rates observed in clinical practice.

**BOTOX** and **BOTOX Cosmetic** contain the same active ingredient in the same formulation, but with different labeled Indications and Usage. Therefore, adverse events observed with the use of **BOTOX Cosmetic** also have the potential to be observed with the use of **BOTOX** and vice-versa.

In general, adverse events occur within the first week following injection of **BOTOX** and while generally transient, may have a duration of several months or longer. Localized pain, infection, inflammation, tenderness, swelling, erythema, and/or bleeding/bruising may be associated with the injection. Needle-related pain and/or anxiety may result in vasovagal responses (including e.g., syncope, hypotension), which may require appropriate medical therapy.

Local weakness of the injected muscle(s) represents the expected pharmacological action of botulinum toxin. However, weakness of nearby muscles may also occur due to spread of toxin [*see Warnings and Precautions (5.2)*].

#### *Chronic Migraine*

In double-blind placebo controlled chronic migraine pivotal efficacy trials (Study 1 and Study 2), the discontinuation rate was 12% in the **BOTOX** treated group and 10% in the placebo-treated group. Discontinuations due to an adverse event were 4% in the **BOTOX** group and 1% in the placebo group. The most frequent adverse events leading to discontinuation in the **BOTOX** group were neck pain, headache, worsening migraine, muscular weakness and eyelid ptosis.

The most frequently reported adverse reactions following injection of **BOTOX** for chronic migraine appear in Table 4.

#### **Table 4: Adverse Reactions Reported by $\geq 2\%$ of BOTOX treated Patients and More Frequent than in Placebo-treated Patients in Two Chronic Migraine Double-blind, Placebo-controlled Clinical Trials**

<b>Adverse Reactions by Body System</b>	<b>BOTOX 155 Units-195 Units (N=687)</b>	<b>Placebo (N=692)</b>
Nervous system disorders		
Headache	32 (5%)	22 (3%)
Migraine	26 (4%)	18 (3%)
Facial paresis	15 (2%)	0 (0%)
Eye disorders		
Eyelid ptosis	25 (4%)	2 (<1%)
Infections and Infestations		
Bronchitis	17 (3%)	11 (2%)
Musculoskeletal and connective tissue disorders		
Neck pain	60 (9%)	19 (3%)
Musculoskeletal stiffness	25 (4%)	6 (1%)
Muscular weakness	24 (4%)	2 (<1%)
Myalgia	21 (3%)	6 (1%)
Musculoskeletal pain	18 (3%)	10 (1%)
Muscle spasms	13 (2%)	6 (1%)
General disorders and administration site conditions		
Injection site pain	23 (3%)	14 (2%)
Vascular Disorders		
Hypertension	11 (2%)	7 (1%)

Other adverse events that occurred more frequently in the **BOTOX** group compared to the placebo group at a frequency less than 1% and potentially **BOTOX** related include: vertigo, dry eye, eyelid edema, dysphagia, eye infection, and jaw pain. Severe worsening of migraine requiring hospitalization occurred in approximately 1% of **BOTOX** treated patients in Study 1 and Study 2, usually within the first week after treatment, compared to 0.3% of placebo-treated patients.

#### *Upper Limb Spasticity*

The most frequently reported adverse reactions following injection of **BOTOX** for adult spasticity appear in Table 5.

**Table 5: Adverse Reactions Reported by ≥2% of BOTOX treated Patients and More Frequent than in Placebo-treated Patients in Adult Spasticity Double-blind, Placebo-controlled Clinical Trials**

<b>Adverse Reactions by Body System</b>	<b>BOTOX 251 Units- 360 Units (N=115)</b>	<b>BOTOX 150 Units- 250 Units (N=188)</b>	<b>BOTOX &lt;150 Units (N=54)</b>	<b>Placebo (N=182)</b>
Gastrointestinal disorder				
Nausea	3 (3%)	3 (2%)	1 (2%)	1 (1%)
General disorders and administration site conditions				
Fatigue	4 (3%)	4 (2%)	1 (2%)	0
Infections and infestations				
Bronchitis	4 (3%)	4 (2%)	0	2 (1%)
Musculoskeletal and connective tissue disorders				
Pain in extremity	7 (6%)	10 (5%)	5 (9%)	8 (4%)
Muscular weakness	0	7 (4%)	1 (2%)	2 (1%)

#### *Cervical Dystonia*

In cervical dystonia patients evaluated for safety in double-blind and open-label studies following injection of **BOTOX**, the most frequently reported adverse reactions were dysphagia (19%), upper respiratory infection (12%), neck pain (11%), and headache (11%).

Other events reported in 2-10% of patients in any one study in decreasing order of incidence include: increased cough, flu syndrome, back pain, rhinitis, dizziness, hypertonia, soreness at injection site, asthenia, oral dryness, speech disorder, fever, nausea, and drowsiness. Stiffness, numbness, diplopia, ptosis, and dyspnea have been reported.

Dysphagia and symptomatic general weakness may be attributable to an extension of the pharmacology of **BOTOX** resulting from the spread of the toxin outside the injected muscles [see *Warnings and Precautions (5.2, 5.4)*].

The most common severe adverse event associated with the use of **BOTOX** injection in patients with cervical dystonia is dysphagia with about 20% of these cases also reporting dyspnea [see *Warnings and Precautions (5.2, 5.4)*]. Most dysphagia is reported as mild or moderate in severity. However, it may be associated with more severe signs and symptoms [see *Warnings and Precautions (5.4)*].

Additionally, reports in the literature include a case of a female patient who developed brachial plexopathy two days after injection of 120 Units of **BOTOX** for the treatment of cervical dystonia, and reports of dysphonia in patients who have been treated for cervical dystonia.

#### *Primary Axillary Hyperhidrosis*

The most frequently reported adverse events (3-10% of adult patients) following injection of **BOTOX** in double-blind studies included injection site pain and hemorrhage, non-axillary sweating, infection, pharyngitis, flu syndrome, headache, fever, neck or back pain, pruritus, and anxiety.

The data reflect 346 patients exposed to **BOTOX** 50 Units and 110 patients exposed to **BOTOX** 75 Units in each axilla.

#### *Blepharospasm*

In a study of blepharospasm patients who received an average dose per eye of 33 Units (injected at 3 to 5 sites) of the currently manufactured **BOTOX**, the most frequently reported treatment-related adverse reactions were ptosis (21%), superficial punctate keratitis (6%), and eye dryness (6%).

Other events reported in prior clinical studies in decreasing order of incidence include: irritation, tearing, lagophthalmos, photophobia, ectropion, keratitis, diplopia, entropion, diffuse skin rash, and local swelling of the eyelid skin lasting for several days following eyelid injection.

In two cases of VII nerve disorder, reduced blinking from **BOTOX** injection of the orbicularis muscle led to serious corneal exposure, persistent epithelial defect, corneal ulceration and a case of corneal perforation. Focal facial paralysis, syncope, and exacerbation of myasthenia gravis have also been reported after treatment of blepharospasm.

#### *Strabismus*

Extraocular muscles adjacent to the injection site can be affected, causing vertical deviation, especially with higher doses of **BOTOX**. The incidence rates of these adverse effects in 2058 adults who received a total of 3650 injections for horizontal strabismus was 17%.

The incidence of ptosis has been reported to be dependent on the location of the injected muscles, 1% after inferior rectus injections, 16% after horizontal rectus injections and 38% after superior rectus injections.

In a series of 5587 injections, retrobulbar hemorrhage occurred in 0.3% of cases.

## **6.2 Post-Marketing Experience**

There have been spontaneous reports of death, sometimes associated with dysphagia, pneumonia, and/or other significant debility or anaphylaxis, after treatment with botulinum toxin [see *Warnings and Precautions (5.3, 5.4)*].

There have also been reports of adverse events involving the cardiovascular system, including arrhythmia and myocardial infarction, some with fatal outcomes. Some of these patients had risk factors including cardiovascular disease. The exact relationship of these events to the botulinum toxin injection has not been established.

New onset or recurrent seizures have also been reported, typically in patients who are predisposed to experiencing these events. The exact relationship of these events to the botulinum toxin injection has not been established.

The following events, not already addressed elsewhere in the package insert, have been reported since the drug has been marketed: abdominal pain; anorexia; brachial plexopathy; diarrhea; facial palsy; facial paresis; hyperhidrosis; hypoacusis; hypoesthesia; localized numbness; malaise; myalgia; paresthesia; pyrexia; radiculopathy; skin rash (including erythema multiforme, and psoriasiform eruption); tinnitus; vertigo; visual disturbances; and vomiting.

Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to botulinum toxin.

## **6.3 Immunogenicity**

As with all therapeutic proteins, there is a potential for immunogenicity. Formation of neutralizing antibodies to botulinum toxin type A may reduce the effectiveness of **BOTOX** treatment by inactivating the biological activity of the toxin.

In a long term, open-label study evaluating 326 cervical dystonia patients treated for an average of 9 treatment sessions with the current formulation of **BOTOX**, 4 (1.2%) patients had positive antibody tests. All 4 of these patients responded to **BOTOX** therapy at the time of the positive antibody test. However, 3 of these patients developed clinical resistance after subsequent treatment, while the fourth patient continued to respond to **BOTOX** therapy for the remainder of the study.

One patient among the 445 hyperhidrosis patients (0.2%), two patients among the 380 adult upper limb spasticity patients (0.5%), and no patients among 406 migraine patients with analyzed specimens showed the presence of neutralizing antibodies.

The data reflect the patients whose test results were considered positive or negative for neutralizing activity to **BOTOX** in a mouse protection assay. The results of these tests are highly dependent on the sensitivity and specificity of the assay. For these reasons, comparison of the incidence of neutralizing activity to **BOTOX** with the incidence reported to other products may be misleading.

The critical factors for neutralizing antibody formation have not been well characterized. The results from some studies suggest that **BOTOX** injections at more frequent intervals or at higher doses may lead to greater incidence of antibody formation. The potential for antibody formation may be minimized by injecting with the lowest effective dose given at the longest feasible intervals between injections.

## 7 DRUG INTERACTIONS

No formal drug interaction studies have been conducted with **BOTOX** (onabotulinumtoxinA) for injection.

Co-administration of **BOTOX** and aminoglycosides or other agents interfering with neuromuscular transmission (e.g., curare-like compounds) should only be performed with caution as the effect of the toxin may be potentiated.

Use of anticholinergic drugs after administration of **BOTOX** may potentiate systemic anticholinergic effects.

The effect of administering different botulinum neurotoxin products at the same time or within several months of each other is unknown. Excessive neuromuscular weakness may be exacerbated by administration of another botulinum toxin prior to the resolution of the effects of a previously administered botulinum toxin.

Excessive weakness may also be exaggerated by administration of a muscle relaxant before or after administration of **BOTOX**.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

Pregnancy Category C.

There are no adequate and well-controlled studies in pregnant women. **BOTOX** should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

When **BOTOX** (4, 8, or 16 Units/kg) was administered intramuscularly to pregnant mice or rats two times during the period of organogenesis (on gestation days 5 and 13), reductions in fetal body weight and decreased fetal skeletal ossification were observed at the two highest doses. The no-effect dose for developmental toxicity in these studies (4 Units/kg) is approximately 1½ times the average high human dose for upper limb spasticity of 360 Units on a body weight basis (Units/kg).

When **BOTOX** was administered intramuscularly to pregnant rats (0.125, 0.25, 0.5, 1, 4, or 8 Units/kg) or rabbits (0.063, 0.125, 0.25, or 0.5 Units/kg) daily during the period of organogenesis (total of 12 doses in rats, 13 doses in rabbits), reduced fetal body weights and decreased fetal skeletal ossification were observed at the two highest doses in rats and at the highest dose in rabbits. These doses were also associated with significant maternal toxicity, including abortions, early deliveries, and maternal death. The developmental no-effect doses in these studies of 1 Unit/kg in rats and 0.25 Units/kg in rabbits are less than the average high human dose based on Units/kg.

When pregnant rats received single intramuscular injections (1, 4, or 16 Units/kg) at three different periods of development (prior to implantation, implantation, or organogenesis), no adverse effects on fetal development were observed. The developmental no-effect level for a single maternal dose in rats (16 Units/kg) is approximately 3 times the average high human dose based on Units/kg.

### 8.3 Nursing Mothers

It is not known whether **BOTOX** is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when **BOTOX** is administered to a nursing woman.

## 8.4 Pediatric Use

### *Prophylaxis of Headaches in Chronic Migraine*

Safety and effectiveness in patients below the age of 18 years have not been established.

### *Spasticity*

Safety and effectiveness in patients below the age of 18 years have not been established.

### *Cervical Dystonia*

Safety and effectiveness in pediatric patients below the age of 16 years have not been established.

### *Blepharospasm and Strabismus*

Safety and effectiveness in pediatric patients below the age of 12 years have not been established.

### *Axillary Hyperhidrosis*

Safety and effectiveness in patients below the age of 18 years have not been established.

## 8.5 Geriatric Use

Clinical studies of **BOTOX** did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. There were too few patients over the age of 75 to enable any comparisons. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

## 10 OVERDOSAGE

Excessive doses of **BOTOX** (onabotulinumtoxinA) for injection may be expected to produce neuromuscular weakness with a variety of symptoms. Respiratory support may be required where excessive doses cause paralysis of respiratory muscles. In the event of overdose, the patient should be medically monitored for symptoms of excessive muscle weakness or muscle paralysis [*see Boxed Warning and Warnings and Precautions (5.2, 5.4)*]. Symptomatic treatment may be necessary.

Symptoms of overdose are likely not to be present immediately following injection. Should accidental injection or oral ingestion occur, the person should be medically supervised for several weeks for signs and symptoms of excessive muscle weakness or paralysis.

In the event of overdose, antitoxin raised against botulinum toxin is available from the Centers for Disease Control and Prevention (CDC) in Atlanta, GA. However, the antitoxin will not reverse any botulinum toxin-induced effects already apparent by the time of antitoxin administration. In the event of suspected or actual cases of botulinum toxin poisoning, please contact your local or state Health Department to process a request for antitoxin through the CDC. If you do not receive a response within 30 minutes, please contact the CDC directly at 1-770-488-7100. More information can be obtained at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5232a8.htm>.

## 11 DESCRIPTION

**BOTOX** (onabotulinumtoxinA) for injection is a sterile, vacuum-dried purified botulinum toxin type A, produced from fermentation of Hall strain *Clostridium botulinum* type A, and intended for intramuscular and intradermal use. It is purified from the culture solution by dialysis and a series of acid precipitations to a complex consisting of the neurotoxin, and several accessory proteins. The complex is dissolved in sterile sodium chloride solution containing Albumin Human and is sterile filtered (0.2 microns) prior to filling and vacuum-drying.

The primary release procedure for **BOTOX** uses a cell-based potency assay to determine the potency relative to a reference standard. The assay is specific to Allergan's products **BOTOX** and **BOTOX<sup>®</sup> Cosmetic**. One Unit of **BOTOX** corresponds to the calculated median intraperitoneal lethal dose (LD<sub>50</sub>) in mice. Due to specific details of this assay such as the vehicle, dilution scheme, and laboratory protocols, Units of biological activity of **BOTOX** cannot be compared to nor converted into Units of any other botulinum toxin or any toxin assessed with any other specific assay method. The specific activity of **BOTOX** is approximately 20 Units/nanogram of neurotoxin protein complex.

Each vial of **BOTOX** contains either 50 Units of *Clostridium botulinum* type A neurotoxin complex, 0.25 mg of Albumin Human, and 0.45 mg of sodium chloride; 100 Units of *Clostridium botulinum* type A neurotoxin complex, 0.5 mg of Albumin Human, and 0.9 mg of sodium chloride; or 200 Units of *Clostridium botulinum* type A neurotoxin complex, 1 mg of Albumin Human, and 1.8 mg of sodium chloride in a sterile, vacuum-dried form without a preservative.

## 12 CLINICAL PHARMACOLOGY

## 12.1 Mechanism of Action

**BOTOX** blocks neuromuscular transmission by binding to acceptor sites on motor or sympathetic nerve terminals, entering the nerve terminals, and inhibiting the release of acetylcholine. This inhibition occurs as the neurotoxin cleaves SNAP-25, a protein integral to the successful docking and release of acetylcholine from vesicles situated within nerve endings. When injected intramuscularly at therapeutic doses, **BOTOX** produces partial chemical denervation of the muscle resulting in a localized reduction in muscle activity. In addition, the muscle may atrophy, axonal sprouting may occur, and extrajunctional acetylcholine receptors may develop. There is evidence that reinnervation of the muscle may occur, thus slowly reversing muscle denervation produced by **BOTOX**.

When injected intradermally, **BOTOX** produces temporary chemical denervation of the sweat gland resulting in local reduction in sweating.

## 12.3 Pharmacokinetics

Using currently available analytical technology, it is not possible to detect **BOTOX** in the peripheral blood following intramuscular injection at the recommended doses.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

#### *Carcinogenesis*

Long term studies in animals have not been performed to evaluate the carcinogenic potential of **BOTOX**.

#### *Mutagenesis*

**BOTOX** was negative in a battery of in vitro (microbial reverse mutation assay, mammalian cell mutation assay, and chromosomal aberration assay) and in vivo (micronucleus assay) genetic toxicologic assays.

#### *Impairment of Fertility*

In fertility studies of **BOTOX** (4, 8, or 16 Units/kg) in which either male or female rats were injected intramuscularly prior to mating and on the day of mating (3 doses, 2 weeks apart for males, 2 doses, 2 weeks apart for females) to untreated animals, reduced fertility was observed in males at the intermediate and high doses and in females at the high dose. The no-effect doses for reproductive toxicity (4 Units/kg in males, 8 Units/kg in females) are approximately equal to the average high human dose for upper limb spasticity of 360 Units on a body weight basis (Units/kg).

## 14 CLINICAL STUDIES

### 14.1 Chronic Migraine

**BOTOX** was evaluated in two randomized, multi-center, 24-week, 2 injection cycle, placebo-controlled double-blind studies. Study 1 and Study 2 included chronic migraine adults who were not using any concurrent headache prophylaxis, and during a 28-day baseline period had  $\geq 15$  headache days lasting 4 hours or more, with  $\geq 50\%$  being migraine/probable migraine. In both studies, patients were randomized to receive placebo or 155 Units to 195 Units **BOTOX** injections every 12 weeks for the 2-cycle, double-blind phase. Patients were allowed to use acute headache treatments during the study. **BOTOX** treatment demonstrated statistically significant and clinically meaningful improvements from baseline compared to placebo for key efficacy variables (see Table 6).

**Table 6: Week 24 Key Efficacy Variables for Study 1 and Study 2**

Efficacy per 28 days	Study 1		Study 2	
	<b>BOTOX</b> (N=341)	<b>Placebo</b> (N=338)	<b>BOTOX</b> (N=347)	<b>Placebo</b> (N=358)
Change from baseline in frequency of headache days	-7.8*	-6.4	-9.2*	-6.9
Change from baseline in total cumulative hours of headache on headache days	-107*	-70	-134*	-95

\* Significantly different from placebo ( $p \leq 0.05$ )

Patients treated with **BOTOX** had a significantly greater mean decrease from baseline in the frequency of headache days at most timepoints from Week 4 to Week 24 in Study 1 (Figure 2), and all timepoints from Week 4 to Week 24 in Study 2 (Figure 3), compared to placebo-treated patients.

**Figure 2: Mean Change from Baseline in Number of Headache Days for Study 1**



## 14.2 Upper Limb Spasticity

The efficacy and safety of **BOTOX** for the treatment of upper limb spasticity were evaluated in three randomized, multi-center, double-blind, placebo-controlled studies.

Study 1 included 126 patients (64 **BOTOX** and 62 placebo) with upper limb spasticity (Ashworth score of at least 3 for wrist flexor tone and at least 2 for finger flexor tone) who were at least 6 months post-stroke. **BOTOX** (a total dose of 200 Units to 240 Units) and placebo were injected intramuscularly (IM) into the flexor digitorum profundus, flexor digitorum sublimis, flexor carpi radialis, flexor carpi ulnaris, and if necessary into the adductor pollicis and flexor pollicis longus (see Table 7). Use of an EMG/nerve stimulator was recommended to assist in proper muscle localization for injection. Patients were followed for 12 weeks.

**Table 7: Study Medication Dose and Injection Sites in Study 1**

Muscles Injected	Volume (mL)	BOTOX (Units)	Number of Injection Sites
<b>Wrist</b>			
Flexor Carpi Radialis	1	50	1
Flexor Carpi Ulnaris	1	50	1
<b>Finger</b>			1
Flexor Digitorum Profundus	1	50	
Flexor Digitorum Sublimis	1	50	1
<b>Thumb</b>			
Adductor Pollicis <sup>a</sup>	0.4	20	1
Flexor Pollicis Longus <sup>a</sup>	0.4	20	1

<sup>a</sup> injected only if spasticity is present in this muscle

The primary efficacy variable was wrist flexors muscle tone at week 6, as measured by the Ashworth score. The Ashworth Scale is a clinical measure of the force required to move an extremity around a joint, with a reduction in score clinically representing a reduction in the force needed to move a joint (i.e., improvement in spasticity).

Possible scores range from 0 to 4:

0 = No increase in muscle tone (none)

1 = Slight increase in muscle tone, giving a 'catch' when the limb was moved in flexion or extension (mild)

2 = More marked increase in muscle tone but affected limb is easily flexed (moderate)

3 = Considerable increase in muscle tone - passive movement difficult (severe)

4 = Limb rigid in flexion or extension (very severe).

Key secondary endpoints included Physician Global Assessment, finger flexors muscle tone, and thumb flexors tone at Week 6. The Physician Global Assessment evaluated the response to treatment in terms of how the patient was doing in his/her life using a scale from -4 = very marked worsening to +4 = very marked improvement. Study 1 results on the primary endpoint and the key secondary endpoints are shown in Table 8.

**Table 8: Primary and Key Secondary Endpoints by Muscle Group at Week 6 in Study 1**

	BOTOX (N=64)	Placebo (N=62)
Median Change from Baseline in Wrist Flexor Muscle Tone on the Ashworth Scale <sup>†a</sup>	-2.0*	0.0
Median Change from Baseline in Finger Flexor Muscle Tone on the Ashworth Scale <sup>††b</sup>	-1.0*	0.0
Median Change from Baseline in Thumb Flexor Muscle Tone on the Ashworth Scale <sup>††c</sup>	-1.0	-1.0
Median Physician Global Assessment of Response to Treatment <sup>††</sup>	2.0*	0.0

<sup>†</sup> Primary endpoint at Week 6

<sup>††</sup> Secondary endpoints at Week 6

\* Significantly different from placebo (p≤0.05)

<sup>a</sup> **BOTOX** injected into both the flexor carpi radialis and ulnaris muscles

<sup>b</sup> **BOTOX** injected into the flexor digitorum profundus and flexor digitorum sublimis muscles

<sup>c</sup> **BOTOX** injected into the adductor pollicis and flexor pollicis longus muscles

Study 2 compared 3 doses of **BOTOX** with placebo and included 91 patients [**BOTOX** 360 Units (N=21), **BOTOX** 180 Units (N=23), **BOTOX** 90 Units (N=21), and placebo (N=26)] with upper limb spasticity (expanded Ashworth score of at least 2 for elbow flexor tone and at least 3 for wrist flexor tone) who were at least 6 weeks post-stroke. **BOTOX** and placebo were injected with EMG guidance into the flexor digitorum profundus, flexor digitorum sublimis, flexor carpi radialis, flexor carpi ulnaris, and biceps brachii (see Table 9).

**Table 9: Study Medication Dose and Injection Sites in Study 2 and Study 3**

Muscles Injected	Total Dose			Volume (mL) per site	Injection Sites (n)
	<b>BOTOX</b> low dose (90 Units)	<b>BOTOX</b> mid dose (180 Units)	<b>BOTOX</b> high dose (360 Units)		
<b>Wrist</b> Flexor Carpi Ulnaris	10 Units	20 Units	40 Units	0.4	1
Flexor Carpi Radialis	15 Units	30 Units	60 Units	0.6	1
<b>Finger</b> Flexor Digitorum Profundus	7.5 Units	15 Units	30 Units	0.3	1
Flexor Digitorum Sublimis	7.5 Units	15 Units	30 Units	0.3	1
<b>Elbow</b> Biceps Brachii	50 Units	100 Units	200 Units	0.5	4

The primary efficacy variable in Study 2 was the wrist flexor tone at Week 6 as measured by the expanded Ashworth Scale. The expanded Ashworth Scale uses the same scoring system as the Ashworth Scale, but allows for half-point increments.

Key secondary endpoints in Study 2 included Physician Global Assessment, finger flexors muscle tone, and elbow flexors muscle tone at Week 6. Study 2 results on the primary endpoint and the key secondary endpoints at Week 6 are shown in Table 10.

**Table 10: Primary and Key Secondary Endpoints by Muscle Group and BOTOX Dose at Week 6 in Study 2**

	<b>BOTOX</b> low dose (90 Units) (N=21)	<b>BOTOX</b> mid dose (180 Units) (N=23)	<b>BOTOX</b> high dose (360 Units) (N=21)	Placebo (N=26)
<b>Median Change from Baseline in Wrist Flexor Muscle Tone on the Ashworth Scale<sup>†b</sup></b>	-1.5*	-1.0*	-1.5*	-1.0
<b>Median Change from Baseline in Finger Flexor Muscle Tone on the Ashworth Scale<sup>††c</sup></b>	-0.5	-0.5	-1.0	-0.5
<b>Median Change from Baseline in Elbow Flexor Muscle Tone on the Ashworth Scale<sup>††d</sup></b>	-0.5	-1.0*	-0.5 <sup>a</sup>	-0.5
<b>Median Physician Global Assessment of Response to Treatment</b>	1.0*	1.0*	1.0*	0.0

<sup>†</sup> Primary endpoint at Week 6

<sup>††</sup> Secondary endpoints at Week 6

\* Significantly different from placebo ( $p \leq 0.05$ )

<sup>a</sup>  $p = 0.053$

<sup>b</sup> Total dose of **BOTOX** injected into both the flexor carpi radialis and ulnaris muscles

<sup>c</sup> Total dose of **BOTOX** injected into the flexor digitorum profundus and flexor digitorum sublimis muscles

<sup>d</sup> Dose of **BOTOX** injected into biceps brachii muscle

Study 3 compared 3 doses of **BOTOX** with placebo and enrolled 88 patients [**BOTOX** 360 Units (N=23), **BOTOX** 180 Units (N=23), **BOTOX** 90 Units (N=23), and placebo (N=19)] with upper limb spasticity (expanded Ashworth score of at least 2 for elbow flexor tone and at least 3 for wrist flexor tone and/or finger flexor tone) who were at least 6 weeks post-stroke. **BOTOX** and placebo were

injected with EMG guidance into the flexor digitorum profundus, flexor digitorum sublimis, flexor carpi radialis, flexor carpi ulnaris, and biceps brachii (see Table 9).

The primary efficacy variable in Study 3 was wrist and elbow flexor tone as measured by the expanded Ashworth score. A key secondary endpoint was assessment of finger flexors muscle tone. Study 3 results on the primary endpoint at Week 4 are shown in Table 11.

**Table 11: Primary and Key Secondary Endpoints by Muscle Group and BOTOX Dose at Week 4 in Study 3**

	<b>BOTOX low dose (90 Units) (N=23)</b>	<b>BOTOX mid dose (180 Units) (N=21)</b>	<b>BOTOX high dose (360 Units) (N=22)</b>	<b>Placebo (N=19)</b>
<b>Median Change from Baseline in Wrist Flexor Muscle Tone on the Ashworth Scale<sup>†b</sup></b>	-1.0	-1.0	-1.5*	-0.5
<b>Median Change from Baseline in Finger Flexor Muscle Tone on the Ashworth Scale<sup>††c</sup></b>	-1.0	-1.0	-1.0*	-0.5
<b>Median Change from Baseline in Elbow Flexor Muscle Tone on the Ashworth Scale<sup>†d</sup></b>	-0.5	-0.5	-1.0*	-0.5

† Primary endpoint at Week 4

†† Secondary endpoints at Week 4

\* Significantly different from placebo ( $p \leq 0.05$ )

<sup>b</sup> Total dose of **BOTOX** injected into both the flexor carpi radialis and ulnaris muscles

<sup>c</sup> Total dose of **BOTOX** injected into the flexor digitorum profundus and flexor digitorum sublimis muscles

<sup>d</sup> Dose of **BOTOX** injected into biceps brachii muscle

### 14.3 Cervical Dystonia

A phase 3 randomized, multi-center, double-blind, placebo-controlled study of the treatment of cervical dystonia was conducted. This study enrolled adult patients with cervical dystonia and a history of having received **BOTOX** in an open label manner with perceived good response and tolerable side effects. Patients were excluded if they had previously received surgical or other denervation treatment for their symptoms or had a known history of neuromuscular disorder. Subjects participated in an open label enrichment period where they received their previously employed dose of **BOTOX**. Only patients who were again perceived as showing a response were advanced to the randomized evaluation period. The muscles in which the blinded study agent injections were to be administered were determined on an individual patient basis.

There were 214 subjects evaluated for the open label period, of which 170 progressed into the randomized, blinded treatment period (88 in the **BOTOX** group, 82 in the placebo group). Patient evaluations continued for at least 10 weeks post-injection. The primary outcome for the study was a dual endpoint, requiring evidence of both a change in the Cervical Dystonia Severity Scale (CDSS) and an increase in the percentage of patients showing any improvement on the Physician Global Assessment Scale at 6 weeks after the injection session. The CDSS quantifies the severity of abnormal head positioning and was newly devised for this study. CDSS allots 1 point for each 5 degrees (or part thereof) of head deviation in each of the three planes of head movement (range of scores up to theoretical maximum of 54). The Physician Global Assessment Scale is a 9 category scale scoring the physician's evaluation of the patients' status compared to baseline, ranging from -4 to +4 (very marked worsening to complete improvement), with 0 indicating no change from baseline and +1 slight improvement. Pain is also an important symptom of cervical dystonia and was evaluated by separate assessments of pain frequency and severity on scales of 0 (no pain) to 4 (constant in frequency or extremely severe in intensity). Study results on the primary endpoints and the pain-related secondary endpoints are shown in Table 12.

**Table 12: Efficacy Outcomes of the Phase 3 Cervical Dystonia Study (Group Means)**

	<b>Placebo (N=82)</b>	<b>BOTOX (N=88)</b>	<b>95% CI on Difference</b>
<b>Baseline CDSS</b>	9.3	9.2	
<b>Change in CDSS at Week 6</b>	-0.3	-1.3	(-2.3, 0.3) <sup>[a,b]</sup>
<b>% Patients with Any Improvement on Physician Global Assessment</b>	31%	51%	(5%, 34%) <sup>[a]</sup>
<b>Pain Intensity Baseline</b>	1.8	1.8	

<b>Change in Pain Intensity at Week 6</b>	-0.1	-0.4	(-0.7, -0.2) <sup>[c]</sup>
<b>Pain Frequency Baseline</b>	1.9	1.8	
<b>Change in Pain Frequency at Week 6</b>	-0.0	-0.3	(-0.5, -0.0) <sup>[c]</sup>

<sup>[a]</sup> Confidence intervals are constructed from the analysis of covariance table with treatment and investigational site as main effects, and baseline CDSS as a covariate.

<sup>[b]</sup> These values represent the prospectively planned method for missing data imputation and statistical test. Sensitivity analyses indicated that the 95% confidence interval excluded the value of no difference between groups and the p-value was less than 0.05. These analyses included several alternative missing data imputation methods and non-parametric statistical tests.

<sup>[c]</sup> Confidence intervals are based on the t-distribution.

Exploratory analyses of this study suggested that the majority of patients who had shown a beneficial response by week 6 had returned to their baseline status by 3 months after treatment. Exploratory analyses of subsets by patient sex and age suggest that both sexes receive benefit, although female patients may receive somewhat greater amounts than male patients. There is a consistent treatment-associated effect between subsets greater than and less than age 65. There were too few non-Caucasian patients enrolled to draw any conclusions regarding relative efficacy in racial subsets.

There were several randomized studies conducted prior to the phase 3 study, which were supportive but not adequately designed to assess or quantitatively estimate the efficacy of **BOTOX**.

In the phase 3 study the median total **BOTOX** dose in patients randomized to receive **BOTOX** (N=88) was 236 Units, with 25th to 75th percentile ranges of 198 Units to 300 Units. Of these 88 patients, most received injections to 3 or 4 muscles; 38 received injections to 3 muscles, 28 to 4 muscles, 5 to 5 muscles, and 5 to 2 muscles. The dose was divided amongst the affected muscles in quantities shown in Table 13. The total dose and muscles selected were tailored to meet individual patient needs.

**Table 13: Number of Patients Treated per Muscle and Fraction of Total Dose Injected into Involved Muscles**

<b>Muscle</b>	<b>Number of Patients Treated in this Muscle (N=88)</b>	<b>Mean % Dose per Muscle</b>	<b>Mid-Range of % Dose per Muscle*</b>
<b>Splenius capitis/cervicis</b>	83	38	25-50
<b>Sternocleidomastoid</b>	77	25	17-31
<b>Levator scapulae</b>	52	20	16-25
<b>Trapezius</b>	49	29	18-33
<b>Semispinalis</b>	16	21	13-25
<b>Scalene</b>	15	15	6-21
<b>Longissimus</b>	8	29	17-41

\*The mid-range of dose is calculated as the 25th to 75th percentiles.

#### **14.4 Primary Axillary Hyperhidrosis**

The efficacy and safety of **BOTOX** for the treatment of primary axillary hyperhidrosis were evaluated in two randomized, multi-center, double-blind, placebo-controlled studies. Study 1 included adult patients with persistent primary axillary hyperhidrosis who scored 3 or 4 on a Hyperhidrosis Disease Severity Scale (HDSS) and who produced at least 50 mg of sweat in each axilla at rest over 5 minutes. HDSS is a 4-point scale with 1 = “underarm sweating is never noticeable and never interferes with my daily activities”; to 4 = “underarm sweating is intolerable and always interferes with my daily activities”. A total of 322 patients were randomized in a 1:1:1 ratio to treatment in both axillae with either 50 Units of **BOTOX**, 75 Units of **BOTOX**, or placebo. Patients were evaluated at 4-week intervals. Patients who responded to the first injection were re-injected when they reported a re-increase in HDSS score to 3 or 4 and produced at least 50 mg sweat in each axilla by gravimetric measurement, but no sooner than 8 weeks after the initial injection.

Study responders were defined as patients who showed at least a 2-grade improvement from baseline value on the HDSS 4 weeks after both of the first two treatment sessions or had a sustained response after their first treatment session and did not receive re-treatment during the study. Spontaneous resting axillary sweat production was assessed by weighing a filter paper held in the axilla over a period of 5 minutes (gravimetric measurement). Sweat production responders were those patients who demonstrated a reduction in axillary sweating from baseline of at least 50% at week 4.

In the three study groups the percentage of patients with baseline HDSS score of 3 ranged from 50% to 54% and from 46% to 50% for a score of 4. The median amount of sweat production (averaged for each axilla) was 102 mg, 123 mg, and 114 mg for the placebo, 50 Units and 75 Units groups respectively.

The percentage of responders based on at least a 2-grade decrease from baseline in HDSS or based on a >50% decrease from baseline in axillary sweat production was greater in both **BOTOX** groups than in the placebo group ( $p<0.001$ ), but was not significantly different between the two **BOTOX** doses (see Table 14).

Duration of response was calculated as the number of days between injection and the date of the first visit at which patients returned to 3 or 4 on the HDSS scale. The median duration of response following the first treatment in **BOTOX** treated patients with either dose was 201 days. Among those who received a second **BOTOX** injection, the median duration of response was similar to that observed after the first treatment.

In study 2, 320 adults with bilateral axillary primary hyperhidrosis were randomized to receive either 50 Units of **BOTOX** ( $n=242$ ) or placebo ( $n=78$ ). Treatment responders were defined as subjects showing at least a 50% reduction from baseline in axillary sweating measured by gravimetric measurement at 4 weeks. At week 4 post-injection, the percentages of responders were 91% (219/242) in the **BOTOX** group and 36% (28/78) in the placebo group,  $p<0.001$ . The difference in percentage of responders between **BOTOX** and placebo was 55% (95% CI=43.3, 65.9).

**Table 14: Study 1 - Study Outcomes**

Treatment Response	<b>BOTOX 50 Units (N=104)</b>	<b>BOTOX 75 Units (N=110)</b>	<b>Placebo (N=108)</b>	<b>BOTOX 50-placebo (95% CI)</b>	<b>BOTOX 75-placebo (95% CI)</b>
<b>HDSS Score change <math>\geq 2</math> (n)<sup>a</sup></b>	55% (57)	49% (54)	6% (6)	49.3% (38.8, 59.7)	43% (33.2, 53.8)
<b>&gt;50% decrease in axillary sweat production % (n)</b>	81% (84)	86% (94)	41% (44)	40% (28.1, 52.0)	45% (33.3, 56.1)

<sup>a</sup> Patients who showed at least a 2-grade improvement from baseline value on the HDSS 4 weeks after both of the first two treatment sessions or had a sustained response after their first treatment session and did not receive re-treatment during the study.

#### 14.5 Blepharospasm

Botulinum toxin has been investigated for use in patients with blepharospasm in several studies. In an open label, historically controlled study, 27 patients with essential blepharospasm were injected with 2 Units of **BOTOX** at each of six sites on each side. Twenty-five of the 27 patients treated with botulinum toxin reported improvement within 48 hours. One patient was controlled with a higher dosage at 13 weeks post initial injection and one patient reported mild improvement but remained functionally impaired.

In another study, 12 patients with blepharospasm were evaluated in a double-blind, placebo-controlled study. Patients receiving botulinum toxin ( $n=8$ ) improved compared with the placebo group ( $n=4$ ). The effects of the treatment lasted a mean of 12 weeks.

One thousand six hundred eighty-four patients with blepharospasm who were evaluated in an open label trial showed clinical improvement as evaluated by measured eyelid force and clinically observed intensity of lid spasm, lasting an average of 12 weeks prior to the need for re-treatment.

#### 14.6 Strabismus

Six hundred seventy-seven patients with strabismus treated with one or more injections of **BOTOX** were evaluated in an open label trial. Fifty-five percent of these patients improved to an alignment of 10 prism diopters or less when evaluated six months or more following injection.

### 16 HOW SUPPLIED/STORAGE AND HANDLING

**BOTOX** is supplied in a single-use vial in the following sizes:

50 Units	NDC 0023-3920-50
100 Units	NDC 0023-1145-01
200 Units	NDC 0023-3921-02

Vials of **BOTOX** have a holographic film on the vial label that contains the name “Allergan” within horizontal lines of rainbow color. In order to see the hologram, rotate the vial back and forth between your fingers under a desk lamp or fluorescent light source. (Note: the holographic film on the label is absent in the date/lot area.) If you do not see the lines of rainbow color or the name “Allergan”, do not use the product and contact Allergan for additional information at 1-800-890-4345 from 7:00 AM to 3:00 PM Pacific Time.

#### Storage

Unopened vials of **BOTOX** should be stored in a refrigerator (2° to 8°C) for up to 36 months for the 50 Units and 100 Units vials or up to 24 months for the 200 Units vial. Do not use after the expiration date on the vial. Administer **BOTOX** within 24 hours of

reconstitution; during this period reconstituted **BOTOX** should be stored in a refrigerator (2° to 8°C). Reconstituted **BOTOX** should be clear, colorless, and free of particulate matter.

All vials, including expired vials, or equipment used with the drug should be disposed of carefully, as is done with all medical waste.

## **Rx Only**

### **17 PATIENT COUNSELING INFORMATION**

Provide a copy of the Medication Guide and review the contents with the patient.

#### **17.1 Swallowing, Speaking or Breathing Difficulties, or Other Unusual Symptoms**

Patients should be advised to inform their doctor or pharmacist if they develop any unusual symptoms (including difficulty with swallowing, speaking, or breathing), or if any existing symptom worsens [*see Boxed Warning and Warnings and Precautions (5.2, 5.4)*].

#### **17.2 Ability to Operate Machinery or Vehicles**

Patients should be counseled that if loss of strength, muscle weakness, blurred vision, or drooping eyelids occur, they should avoid driving a car or engaging in other potentially hazardous activities.

#### **17.3 Medication Guide**

**MEDICATION GUIDE**  
**BOTOX<sup>®</sup>**  
**BOTOX<sup>®</sup> Cosmetic**  
**(Boe-tox)**  
**(onabotulinumtoxinA)**  
**for Injection**

Read the Medication Guide that comes with **BOTOX** or **BOTOX Cosmetic** before you start using it and each time it is given to you. There may be new information. This information does not take the place of talking with your doctor about your medical condition or your treatment. You should share this information with your family members and caregivers.

**What is the most important information I should know about BOTOX and BOTOX Cosmetic?**

**BOTOX and BOTOX Cosmetic may cause serious side effects that can be life threatening. Call your doctor or get medical help right away if you have any of these problems after treatment with BOTOX or BOTOX Cosmetic:**

- **Problems swallowing, speaking, or breathing.** These problems can happen hours to weeks after an injection of **BOTOX** or **BOTOX Cosmetic** usually because the muscles that you use to breathe and swallow can become weak after the injection. Death can happen as a complication if you have severe problems with swallowing or breathing after treatment with **BOTOX** or **BOTOX Cosmetic**.
- People with certain breathing problems may need to use muscles in their neck to help them breathe. These patients may be at greater risk for serious breathing problems with **BOTOX** or **BOTOX Cosmetic**.
- Swallowing problems may last for several months. People who cannot swallow well may need a feeding tube to receive food and water. If swallowing problems are severe, food or liquids may go into your lungs. People who already have swallowing or breathing problems before receiving **BOTOX** or **BOTOX Cosmetic** have the highest risk of getting these problems.
- **Spread of toxin effects.** In some cases, the effect of botulinum toxin may affect areas of the body away from the injection site and cause symptoms of a serious condition called botulism. The symptoms of botulism include:
  - loss of strength and muscle weakness all over the body
  - double vision
  - blurred vision and drooping eyelids
  - hoarseness or change or loss of voice (dysphonia)
  - trouble saying words clearly (dysarthria)
  - loss of bladder control
  - trouble breathing
  - trouble swallowing

These symptoms can happen hours to weeks after you receive an injection of **BOTOX** or **BOTOX Cosmetic**.

These problems could make it unsafe for you to drive a car or do other dangerous activities. See "What should I avoid while receiving **BOTOX** or **BOTOX Cosmetic**?"

There has not been a confirmed serious case of spread of toxin effect away from the injection site when **BOTOX** has been used at the recommended dose to treat chronic migraine, severe underarm sweating, blepharospasm, or strabismus, or when **BOTOX Cosmetic** has been used at the recommended dose to treat frown lines.

**What are BOTOX and BOTOX Cosmetic?**

**BOTOX** is a prescription medicine that is injected into muscles and used:

- to prevent headaches in adults with chronic migraine who have 15 or more days each month with headache lasting 4 or more hours each day.
- to treat increased muscle stiffness in elbow, wrist, and finger muscles in adults with upper limb spasticity.
- to treat the abnormal head position and neck pain that happens with cervical dystonia (CD) in adults.
- to treat certain types of eye muscle problems (strabismus) or abnormal spasm of the eyelids (blepharospasm) in people 12 years and older.

**BOTOX** is also injected into the skin to treat the symptoms of severe underarm sweating (severe primary axillary hyperhidrosis) when medicines used on the skin (topical) do not work well enough.

**BOTOX Cosmetic** is a prescription medicine that is injected into muscles and used to improve the look of moderate to severe frown lines between the eyebrows (glabellar lines) in adults younger than 65 years of age for a short period of time (temporary).

It is not known whether **BOTOX** is safe or effective in patients younger than:

- 18 years of age for treatment of chronic migraine
- 18 years of age for treatment of spasticity
- 16 years of age for treatment of cervical dystonia
- 18 years of age for treatment of hyperhidrosis
- 12 years of age for treatment of strabismus or blepharospasm

**BOTOX Cosmetic** is not recommended for use in children younger than 18 years of age.

It is not known whether **BOTOX** and **BOTOX Cosmetic** are safe or effective to prevent headaches in patients with migraine who have 14 or fewer headache days each month (episodic migraine).

It is not known whether **BOTOX** and **BOTOX Cosmetic** are safe or effective for other types of muscle spasms or for severe sweating anywhere other than your armpits.

### **Who should not take BOTOX or BOTOX Cosmetic?**

Do not take **BOTOX** or **BOTOX Cosmetic** if you:

- are allergic to any of the ingredients in **BOTOX** or **BOTOX Cosmetic**. See the end of this Medication Guide for a list of ingredients in **BOTOX** and **BOTOX Cosmetic**.
- had an allergic reaction to any other botulinum toxin product such as *Myobloc*<sup>®</sup>, *Dysport*<sup>®</sup>, or *Xeomin*<sup>®</sup>
- have a skin infection at the planned injection site

### **What should I tell my doctor before taking BOTOX or BOTOX Cosmetic?**

**Tell your doctor about all your medical conditions, including if you have:**

- a disease that affects your muscles and nerves (such as amyotrophic lateral sclerosis [ALS or Lou Gehrig's disease], myasthenia gravis or Lambert-Eaton syndrome). See "What is the most important information I should know about **BOTOX** and **BOTOX Cosmetic**?"
- allergies to any botulinum toxin product
- had any side effect from any botulinum toxin product in the past
- a breathing problem, such as asthma or emphysema
- swallowing problems
- bleeding problems
- plans to have surgery
- had surgery on your face
- weakness of your forehead muscles, such as trouble raising your eyebrows
- drooping eyelids
- any other change in the way your face normally looks
- are pregnant or plan to become pregnant. It is not known if **BOTOX** or **BOTOX Cosmetic** can harm your unborn baby.
- are breast-feeding or plan to breastfeed. It is not known if **BOTOX** or **BOTOX Cosmetic** passes into breast milk.

**Tell your doctor about all the medicines you take**, including prescription and nonprescription medicines, vitamins and herbal products. Using **BOTOX** or **BOTOX Cosmetic** with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received BOTOX or BOTOX Cosmetic in the past.**

Especially tell your doctor if you:

- have received any other botulinum toxin product in the last four months
- have received injections of botulinum toxin, such as *Myobloc*<sup>®</sup> (rimabotulinumtoxinB), *Dysport*<sup>®</sup> (abobotulinumtoxinA), or *Xeomin*<sup>®</sup> (incobotulinumtoxinA) in the past. Be sure your doctor knows exactly which product you received.
- have recently received an antibiotic by injection
- take muscle relaxants
- take an allergy or cold medicine
- take a sleep medicine

**Ask your doctor if you are not sure if your medicine is one that is listed above.**

Know the medicines you take. Keep a list of your medicines with you to show your doctor and pharmacist each time you get a new medicine.

#### **How should I take BOTOX or BOTOX Cosmetic?**

- **BOTOX** or **BOTOX Cosmetic** is an injection that your doctor will give you.
- **BOTOX** is injected into your affected muscles or skin.
- **BOTOX Cosmetic** is injected into your affected muscles.
- Your doctor may change your dose of **BOTOX** or **BOTOX Cosmetic**, until you and your doctor find the best dose for you.
- Your doctor will tell you how often you will receive your dose of **BOTOX** or **BOTOX Cosmetic** injections.

#### **What should I avoid while taking BOTOX or BOTOX Cosmetic?**

**BOTOX** and **BOTOX Cosmetic** may cause loss of strength or general muscle weakness, or vision problems within hours to weeks of taking **BOTOX** or **BOTOX Cosmetic**. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.** See "What is the most important information I should know about **BOTOX** and **BOTOX Cosmetic**?"

#### **What are the possible side effects of BOTOX and BOTOX Cosmetic?**

**BOTOX** and **BOTOX Cosmetic** can cause serious side effects. See "What is the most important information I should know about **BOTOX** and **BOTOX Cosmetic**?"

#### **Other side effects of BOTOX and BOTOX Cosmetic include:**

- dry mouth
- discomfort or pain at the injection site
- tiredness
- headache
- neck pain
- eye problems: double vision, blurred vision, decreased eyesight, drooping eyelids, swelling of your eyelids, and dry eyes.
- allergic reactions. Symptoms of an allergic reaction to **BOTOX** or **BOTOX Cosmetic** may include: itching, rash, red itchy welts, wheezing, asthma symptoms, or dizziness or feeling faint. Tell your doctor or get medical help right away if you are wheezing or have asthma symptoms, or if you become dizzy or faint.

Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of **BOTOX** and **BOTOX Cosmetic**. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

#### **General information about BOTOX and BOTOX Cosmetic:**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide.

This Medication Guide summarizes the most important information about **BOTOX** and **BOTOX Cosmetic**. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about **BOTOX** and **BOTOX Cosmetic** that is written for healthcare professionals. For more information about **BOTOX** and **BOTOX Cosmetic** call Allergan at 1-800-433-8871 or go to [www.botox.com](http://www.botox.com).

#### **What are the ingredients in BOTOX and BOTOX Cosmetic?**

Active ingredient: botulinum toxin type A

Inactive ingredients: human albumin and sodium chloride

#### **Issued: 10/2010**

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Manufactured by: Allergan Pharmaceuticals Ireland

a subsidiary of: Allergan, Inc.

2525 Dupont Dr.

Irvine, CA 92612

© 2011 Allergan, Inc.

® mark owned by Allergan, Inc.

U.S. Patents 6,974,578; 6,683,049; and 6,896,886

*Myobloc*<sup>®</sup> is a registered trademark of Solstice Neurosciences, Inc.

*Dysport*<sup>®</sup> is a registered trademark of Ipsen Biopharm Limited Company.

*Xeomin*<sup>®</sup> is a registered trademark of Merz Pharma GmbH & Co KGaA.



**BOTOX<sup>®</sup> Cosmetic**  
**(onabotulinumtoxinA)**  
**for injection**

Manufactured by: Allergan Pharmaceuticals Ireland  
a subsidiary of: Allergan, Inc. 2525 Dupont Dr., Irvine, CA 92612

**Distant Spread of Toxin Effect**

Postmarketing reports indicate that the effects of **BOTOX<sup>®</sup> Cosmetic** and all botulinum toxin products may spread from the area of injection to produce symptoms consistent with botulinum toxin effects. These may include asthenia, generalized muscle weakness, diplopia, blurred vision, ptosis, dysphagia, dysphonia, dysarthria, urinary incontinence, and breathing difficulties. These symptoms have been reported hours to weeks after injection. Swallowing and breathing difficulties can be life threatening and there have been reports of death. The risk of symptoms is probably greatest in children treated for spasticity but symptoms can also occur in adults treated for spasticity and other conditions, particularly in those patients who have underlying conditions that would predispose them to these symptoms. In unapproved uses, including spasticity in children and adults, and in approved indications, cases of spread of effect have occurred at doses comparable to those used to treat cervical dystonia and at lower doses.

**DESCRIPTION**

**BOTOX<sup>®</sup> Cosmetic (onabotulinumtoxinA)** for injection, is a sterile, vacuum-dried purified botulinum toxin type A, produced from fermentation of Hall strain *Clostridium botulinum* type A grown in a medium containing casein hydrolysate, glucose, and yeast extract, intended for intramuscular use. It is purified from the culture solution by dialysis and a series of acid precipitations to a complex consisting of the neurotoxin, and several accessory proteins. The complex is dissolved in sterile sodium chloride solution containing Albumin Human and is sterile filtered (0.2 microns) prior to filling and vacuum-drying.

The primary release procedure for **BOTOX<sup>®</sup> Cosmetic** uses a cell-based potency assay to determine the potency relative to a reference standard. The assay is specific to Allergan's products **BOTOX<sup>®</sup>** and **BOTOX<sup>®</sup> Cosmetic**. One Unit of **BOTOX<sup>®</sup> Cosmetic** corresponds to the calculated median intraperitoneal lethal dose (LD<sub>50</sub>) in mice. Due to specific details of this assay such as the vehicle, dilution scheme and laboratory protocols, Units of biological activity of **BOTOX<sup>®</sup> Cosmetic** cannot be compared to nor converted into Units of any other botulinum toxin or any toxin assessed with any other specific assay method. The specific activity of **BOTOX<sup>®</sup> Cosmetic** is approximately 20 Units/nanogram of neurotoxin protein complex.

Each vial of **BOTOX<sup>®</sup> Cosmetic** contains either 100 Units of *Clostridium botulinum* type A neurotoxin complex, 0.5 mg of Albumin Human, and 0.9 mg of sodium chloride or 50 Units of *Clostridium botulinum* type A neurotoxin complex, 0.25 mg of Albumin Human, and 0.45 mg of sodium chloride in a sterile, vacuum-dried form without a preservative.

**CLINICAL PHARMACOLOGY**

**BOTOX<sup>®</sup> Cosmetic** blocks neuromuscular transmission by binding to acceptor sites on motor nerve terminals, entering the nerve terminals, and inhibiting the release of acetylcholine. This inhibition occurs as the neurotoxin cleaves SNAP-25, a protein integral to the successful docking and release of acetylcholine from vesicles situated within nerve endings. When injected intramuscularly at therapeutic doses, **BOTOX<sup>®</sup> Cosmetic** produces partial chemical denervation of the muscle resulting in a localized reduction in muscle activity. In addition, the muscle may atrophy, axonal sprouting may occur, and extrajunctional acetylcholine receptors may develop. There is evidence that reinnervation of the muscle may occur, thus slowly reversing muscle denervation produced by **BOTOX<sup>®</sup> Cosmetic**.

### **Pharmacokinetics**

Using currently available analytical technology, it is not possible to detect **BOTOX<sup>®</sup> Cosmetic** in the peripheral blood following intramuscular injection at the recommended doses.

## **CLINICAL STUDIES**

### **Glabellar Lines**

Two phase 3 randomized, multi-center, double-blind, placebo-controlled studies of identical design were conducted to evaluate **BOTOX<sup>®</sup> Cosmetic** for use in the temporary improvement of the appearance of moderate to severe glabellar facial lines. The studies enrolled healthy adults (ages 18 to 75) with glabellar lines of at least moderate severity at maximum frown. Patients were excluded if they had ptosis, deep dermal scarring, or an inability to substantially lessen glabellar lines even by physically spreading them apart. Subjects received a single treatment with **BOTOX<sup>®</sup> Cosmetic** (N=405, combined studies) or placebo (N=132, combined studies). Injection volume was 0.1 mL/injection site, for a dose/injection site in the active treatment groups of 4 Units. Subjects were injected intramuscularly in five sites, 1 in the procerus muscle and 2 in each corrugator supercilii muscle, for a total dose in the active treatment groups of 20 Units.

The co-primary efficacy endpoints were the investigator's rating of glabellar line severity at maximum frown and the subject's global assessment of change in appearance of glabellar lines, both at Day 30 post-injection. For the investigator rating, using a 4-point grading scale (0=none, 3=severe) a responder was defined as having a severity grade of 0 or 1. For the subject's global assessment of change, the ratings were from +4 (complete improvement) to -4 (very marked worsening). A responder was defined as having a grade of at least +2 (moderate improvement). After completion of the randomized studies, subjects were offered participation in an open label, repeat treatment study to assess the safety of repeated treatment sessions.

The combined results of these two efficacy trials are presented here. The mean age was 46 years, with 32 patients (6%)  $\geq$  65 years of age. Most of the subjects (82%) were women, and Caucasian (84%). At baseline, 210 patients (39%) had glabellar line severity scores at rest of moderate or severe.

In these studies, the severity of glabellar lines was reduced for up to 120 days in the **BOTOX<sup>®</sup> Cosmetic** group compared to the placebo group as measured both by investigator rating of glabellar line severity at maximum frown (Table 1), and by subject's global assessment of change in appearance of glabellar lines (Table 2).

**TABLE 1.****Investigator's Assessment of Glabellar Line Severity at Maximum Frown – Responder Rates (% and Number of Subjects with Severity of None or Mild)**

Day	BOTOX <sup>®</sup> Cosmetic	Placebo	Difference <sup>a</sup>
7	74% 299/405	6% 8/132	68% (62, 74)
30 <sup>b</sup>	80% 325/405	3% 4/132	77% (72, 82)
60	70% 283/403	2% 2/130	69% (64, 74)
90	48% 192/403	2% 3/128	45% (40, 51)
120	25% 102/403	2% 2/128	24% (19, 29)

<sup>a</sup> 95% confidence intervals are shown in parenthesis

<sup>b</sup> Day 30: Co-Primary Efficacy Time point, P<0.001

**TABLE 2.****Subject's Assessment of Change in Appearance of Glabellar Lines – Responder Rates (% and Number of Subjects with at Least Moderate Improvement)**

Day	BOTOX <sup>®</sup> Cosmetic	Placebo	Difference <sup>a</sup>
7	82% 334/405	9% 12/132	73% (68, 80)
30 <sup>b</sup>	89% 362/405	7% 9/132	83% (77, 88)
60	82% 330/403	4% 5/130	78% (73, 83)
90	63% 254/403	3% 4/128	60% (54, 66)
120	39% 157/403	1% 1/128	38% (33, 43)

<sup>a</sup> 95% confidence intervals are shown in parenthesis

<sup>b</sup> Day 30: Co-Primary Efficacy Time point, P<0.001

In the subset of patients with resting severity scores of moderate or severe, the investigator assessment of a resting severity of mild or none at day 30 was also achieved by more **BOTOX<sup>®</sup> Cosmetic** treated patients (74%, 119/161) than placebo treated patients (20%, 10/49).

Analysis of the limited number of patients 65 years or older suggested a lower treatment-associated response compared to patients less than 65 years of age. (Table 3).

**TABLE 3.****Investigator’s and Subject’s Assessment – Responder Rates for Subjects < 65 and ≥ 65 Years of Age at Day 30**

Assessment	Age Group	<b>BOTOX<sup>®</sup> Cosmetic</b> N=405	Placebo N=132	Difference <sup>a</sup>
<b>Investigators (maximal frown)</b>	< 65	83% 316/382	2% 2/123	81% (77, 86)
<b>Subjects</b>	< 65	91% 346/382	7% 8/123	84% (79, 90)
<b>Investigators (maximal frown)</b>	≥ 65	39% 9/23	22% 2/9	17% (-17, 51)
<b>Subjects</b>	≥ 65	70% 16/23	11% 1/9	58% (31, 86)

<sup>a</sup> 95% confidence intervals are shown in parenthesis

Exploratory analyses by gender suggested that responder rates in the **BOTOX<sup>®</sup> Cosmetic** treated group were higher for women than for men for both the investigator assessment (day 30; 85% of 334 women, 59% of 71 men) and the Subject Assessment (day 30; 93% of women, 72% of men). In the limited number of non-Caucasian patients (n=64 in the **BOTOX<sup>®</sup> Cosmetic** treated group) the responder rates were similar to those observed in the Caucasian patients.

**INDICATIONS AND USAGE**

**BOTOX<sup>®</sup> Cosmetic** is indicated for the temporary improvement in the appearance of moderate to severe glabellar lines associated with corrugator and/or procerus muscle activity in adult patients ≤ 65 years of age.

**CONTRAINDICATIONS**

**BOTOX<sup>®</sup> Cosmetic** is contraindicated in the presence of infection at the proposed injection site(s) and in individuals with known hypersensitivity to any botulinum toxin preparation or to any of the components in the formulation.

**WARNINGS**

**BOTOX<sup>®</sup>** and **BOTOX<sup>®</sup> Cosmetic** contain the same active ingredient in the same formulation. Therefore, adverse events observed with the use of **BOTOX<sup>®</sup>** also have the potential to be associated with the use of **BOTOX<sup>®</sup> Cosmetic**.

The recommended dosage and frequency of administration for **BOTOX<sup>®</sup> Cosmetic** should not be exceeded. Risks resulting from administration at higher dosages are not known.

**Lack of Interchangeability between Botulinum Toxin Products**

The potency Units of **BOTOX<sup>®</sup> Cosmetic** are specific to the preparation and assay method utilized. They are not interchangeable with other preparations of botulinum toxin products and, therefore, units of biological activity of **BOTOX<sup>®</sup> Cosmetic** cannot be compared to or converted into units of any other botulinum toxin products assessed with any other specific assay method (see DESCRIPTION).

### **Spread of Toxin Effect**

Postmarketing safety data from **BOTOX**<sup>®</sup> **Cosmetic** and other approved botulinum toxins suggest that botulinum toxin effects may, in some cases, be observed beyond the site of local injection. The symptoms are consistent with the mechanism of action of botulinum toxin and may include asthenia, generalized muscle weakness, diplopia, blurred vision, ptosis, dysphagia, dysphonia, dysarthria, urinary incontinence, and breathing difficulties. These symptoms have been reported hours to weeks after injection. Swallowing and breathing difficulties can be life threatening and there have been reports of death related to spread of toxin effects. The risk of symptoms is probably greatest in children treated for spasticity but symptoms can also occur in adults treated for spasticity and other conditions, and particularly in those patients who have underlying conditions that would predispose them to these symptoms. In unapproved uses, including spasticity in children and adults, and in approved indications, symptoms consistent with spread of toxin effect have been reported at doses comparable to or lower than doses used to treat cervical dystonia.

No definitive serious adverse event reports of distant spread of toxin effect associated with dermatologic use of **BOTOX**<sup>®</sup>/**BOTOX**<sup>®</sup> **Cosmetic** at the labeled dose of 20 Units (for glabellar lines) or 100 Units (for severe primary axillary hyperhidrosis) have been reported.

No definitive serious adverse event reports of distant spread of toxin effect associated with **BOTOX**<sup>®</sup> for blepharospasm at the recommended dose (30 Units and below) or for strabismus at the labeled doses have been reported.

### **Hypersensitivity Reactions**

Serious and/or immediate hypersensitivity reactions have been reported. These reactions include anaphylaxis, urticaria, soft tissue edema, and dyspnea. If such a reaction occurs, further injection of **BOTOX**<sup>®</sup> **Cosmetic** should be discontinued and appropriate medical therapy immediately instituted. One fatal case of anaphylaxis has been reported in which lidocaine was used as the diluent, and consequently the causal agent cannot be reliably determined.

### **Pre-Existing Neuromuscular Disorders**

Individuals with peripheral motor neuropathic diseases, amyotrophic lateral sclerosis, or neuromuscular junctional disorders (e.g., myasthenia gravis or Lambert-Eaton syndrome) should be monitored particularly closely when given botulinum toxin. Patients with neuromuscular disorders may be at increased risk of clinically significant effects including severe dysphagia and respiratory compromise from typical doses of **BOTOX**<sup>®</sup> **Cosmetic** (see **ADVERSE REACTIONS**).

### **Dysphagia and Breathing Difficulties in Treatment of Cervical Dystonia**

Treatment with **BOTOX**<sup>®</sup> and other botulinum toxin products can result in swallowing or breathing difficulties. Patients with pre-existing swallowing or breathing difficulties may be more susceptible to these complications. In most cases, this is a consequence of weakening of muscles in the area of injection that are involved in breathing or swallowing. When distant effects occur, additional respiratory muscles may be involved (see **WARNINGS**).

Deaths as a complication of severe dysphagia have been reported after treatment with botulinum toxin. Dysphagia may persist for several months, and require use of a feeding tube to maintain adequate nutrition and hydration. Aspiration may result from severe dysphagia and is a particular risk when treating patients in whom swallowing or respiratory function is already compromised.

Treatment of cervical dystonia with botulinum toxins may weaken neck muscles that serve as accessory muscles of ventilation. This may result in a critical loss of breathing capacity in patients with respiratory disorders who may have become dependent upon these accessory muscles. There have been postmarketing reports of serious breathing difficulties, including respiratory failure, in cervical dystonia patients.

Patients treated with botulinum toxin may require immediate medical attention should they develop problems with swallowing, speech, or respiratory disorders. These reactions can occur within hours to weeks after injection with botulinum toxin (see **WARNINGS, ADVERSE REACTIONS, CLINICAL PHARMACOLOGY**).

### **Cardiovascular System**

There have been reports following administration of **BOTOX**<sup>®</sup> of adverse events involving the cardiovascular system, including arrhythmia and myocardial infarction, some with fatal outcomes. Some of these patients had risk factors including pre-existing cardiovascular disease.

### **Human Albumin**

This product contains albumin, a derivative of human blood. Based on effective donor screening and product manufacturing processes, it carries an extremely remote risk for transmission of viral diseases. A theoretical risk for transmission of Creutzfeldt-Jakob disease (CJD) also is considered extremely remote. No cases of transmission of viral diseases or CJD have ever been identified for albumin.

### **PRECAUTIONS**

The safe and effective use of **BOTOX**<sup>®</sup> **Cosmetic** depends upon proper storage of the product, selection of the correct dose, and proper reconstitution and administration techniques. Physicians administering **BOTOX**<sup>®</sup> **Cosmetic** must understand the relevant neuromuscular and/or orbital anatomy of the area involved, as well as any alterations to the anatomy due to prior surgical procedures and avoid injection into vulnerable anatomic areas.

Caution should be used when **BOTOX**<sup>®</sup> **Cosmetic** treatment is used in the presence of inflammation at the proposed injection site(s) or when excessive weakness or atrophy is present in the target muscle(s).

Reduced blinking from **BOTOX**<sup>®</sup> **Cosmetic** injection of the orbicularis muscle can lead to corneal exposure, persistent epithelial defect and corneal ulceration, especially in patients with VII nerve disorders. In the use of **BOTOX**<sup>®</sup> for the treatment of blepharospasm, one case of corneal perforation in an aphakic eye requiring corneal grafting has occurred because of this effect. Careful testing of corneal sensation in eyes previously operated upon, avoidance of injection into the lower lid area to avoid ectropion, and vigorous treatment of any epithelial

defect should be employed. This may require protective drops, ointment, therapeutic soft contact lenses, or closure of the eye by patching or other means.

Inducing paralysis in one or more extraocular muscles may produce spatial disorientation, double vision or past pointing. Covering the affected eye may alleviate these symptoms.

Caution should be used when **BOTOX<sup>®</sup> Cosmetic** treatment is used in patients who have an inflammatory skin problem at the injection site, marked facial asymmetry, ptosis, excessive dermatochalasis, deep dermal scarring, thick sebaceous skin or the inability to substantially lessen glabellar lines by physically spreading them apart as these patients were excluded from the Phase 3 safety and efficacy trials.

Needle-related pain and/or anxiety may result in vasovagal responses (including e.g., syncope, hypotension), which may require appropriate medical therapy.

Injection intervals of **BOTOX<sup>®</sup> Cosmetic** should be no more frequent than every three months and should be performed using the lowest effective dose (see **ADVERSE REACTIONS, IMMUNOGENICITY**).

### **Information for Patients**

The physician should provide a copy of the FDA-Approved Patient Medication Guide and review the contents with the patient. Patients should be advised to inform their doctor or pharmacist if they develop any unusual symptoms (including difficulty with swallowing, speaking, or breathing), or if any existing symptom worsens.

Patients should be counseled that if loss of strength, muscle weakness, or impaired vision occur, they should avoid driving a car or engaging in other potentially hazardous activities.

### **Drug Interactions**

Co-administration of **BOTOX<sup>®</sup> Cosmetic** and aminoglycosides<sup>1</sup> or other agents interfering with neuromuscular transmission (e.g., curare-like nondepolarizing blockers, lincosamides, polymyxins, quinidine, magnesium sulfate, anticholinesterases, succinylcholine chloride) should only be performed with caution as the effect of the toxin may be potentiated.

The effect of administering different botulinum neurotoxin serotypes at the same time or within several months of each other is unknown. Excessive neuromuscular weakness may be exacerbated by administration of another botulinum toxin prior to the resolution of the effects of a previously administered botulinum toxin.

### **Pregnancy: Pregnancy Category C**

Administration of **BOTOX<sup>®</sup> Cosmetic** is not recommended during pregnancy. There are no adequate and well-controlled studies of **BOTOX<sup>®</sup> Cosmetic** in pregnant women. When pregnant mice and rats were injected intramuscularly during the period of organogenesis, the developmental NOEL (No Observed Effect Level) of **BOTOX<sup>®</sup> Cosmetic** was 4 Units/kg. Higher doses (8 Units/kg or 16 Units/kg) were associated with reductions in fetal body weights and/or delayed ossification.

In a range finding study in rabbits, daily injection of 0.125 Units/kg/day (days 6 to 18 of gestation) and 2 Units/kg (days 6 and 13 of gestation) produced severe maternal toxicity, abortions and/or fetal malformations. Higher doses resulted in death of the dams. The rabbit appears to be a very sensitive species to **BOTOX<sup>®</sup> Cosmetic**.

If the patient becomes pregnant after the administration of this drug, the patient should be apprised of the potential risks, including abortion or fetal malformations that have been observed in rabbits.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long term studies in animals have not been performed to evaluate carcinogenic potential of **BOTOX<sup>®</sup> Cosmetic**.

The reproductive NOEL following intramuscular injection of 0, 4, 8, and 16 Units/kg was 4 Units/kg in male rats and 8 Units/kg in female rats. Higher doses were associated with dose-dependent reductions in fertility in male rats (where limb weakness resulted in the inability to mate), and testicular atrophy or an altered estrous cycle in female rats. There were no adverse effects on the viability of the embryos.

### **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when **BOTOX<sup>®</sup> Cosmetic** is administered to a nursing woman.

### **Pediatric Use**

Use of **BOTOX<sup>®</sup> Cosmetic** is not recommended in children.

### **Geriatric Use**

The two clinical studies of **BOTOX<sup>®</sup> Cosmetic** did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. However, the responder rates appeared to be higher for patients younger than age 65 than for patients 65 years or older (see **CLINICAL STUDIES**).

There were too few patients (N=3) over the age of 75 to allow any meaningful comparisons.

## **ADVERSE REACTIONS**

### **General**

**BOTOX<sup>®</sup>** and **BOTOX<sup>®</sup> Cosmetic** contain the same active ingredient in the same formulation. Therefore adverse events observed with the use of **BOTOX<sup>®</sup>** also have the potential to be associated with the use of **BOTOX<sup>®</sup> Cosmetic**.

The most serious adverse events reported after treatment with botulinum toxin include spontaneous reports of death, sometimes associated with anaphylaxis, dysphagia, pneumonia, and/or other significant debility.

There have also been reports of adverse events involving the cardiovascular system, including arrhythmia and myocardial infarction, some with fatal outcomes. Some of these patients had risk factors including pre-existing cardiovascular disease (see **WARNINGS**).

New onset or recurrent seizures have also been reported, typically in patients who are predisposed to experiencing these events. The exact relationship of these events to the botulinum toxin injection has not been established. Additionally, a report of acute angle closure glaucoma one day after receiving an injection of botulinum toxin for blepharospasm was received, with recovery four months later after laser iridotomy and trabeculectomy. Focal facial paralysis, syncope and exacerbation of myasthenia gravis have also been reported after treatment of blepharospasm.

In general, adverse events occur within the first week following injection of **BOTOX<sup>®</sup> Cosmetic** and while generally transient may have a duration of several months or longer. Localized pain, infection, inflammation, tenderness, swelling, erythema and/or bleeding/bruising may be associated with the injection. Local weakness of the injected muscle(s) represents the expected pharmacological action of botulinum toxin. However, weakness of adjacent muscles may also occur due to spread of toxin.

#### **Glabellar Lines**

In clinical trials of **BOTOX<sup>®</sup> Cosmetic** the most frequently reported adverse events following injection of **BOTOX<sup>®</sup> Cosmetic** were headache\*, respiratory infection\*, flu syndrome\*, blepharoptosis and nausea.

Less frequently occurring (<3%) adverse reactions included pain in the face, erythema at the injection site\*, paresthesia\* and muscle weakness. While local weakness of the injected muscle(s) is representative of the expected pharmacological action of botulinum toxin, weakness of adjacent muscles may occur as a result of the spread of toxin. These events are thought to be associated with the injection and occurred within the first week. The events were generally transient but may last several months or longer.

(\* incidence not different from Placebo)

The data described in Table 4 reflect exposure to **BOTOX<sup>®</sup> Cosmetic** in 405 subjects aged 18 to 75 who were evaluated in the randomized, placebo-controlled clinical studies to assess the use of **BOTOX<sup>®</sup> Cosmetic** in the improvement of the appearance of glabellar lines (see **CLINICAL STUDIES**). Adverse events of any cause were reported for 44% of the **BOTOX<sup>®</sup> Cosmetic** treated subjects and 42% of the placebo treated subjects. The incidence of blepharoptosis was higher in the **BOTOX<sup>®</sup> Cosmetic** treated arm than in placebo (3% vs. 0).

In the open-label, repeat injection study, blepharoptosis was reported for 2% (8/373) of subjects in the first treatment cycle and 1% (4/343) of subjects in the second treatment cycle. Adverse events of any type were reported for 49% (183/373) of subjects overall. The most frequently reported of these adverse events in the open-label study included respiratory infection, headache, flu syndrome, blepharoptosis, pain and nausea.

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not be predictive of rates observed in practice.

**TABLE 4.**  
**Adverse Events Reported at Higher Frequency (>1%) in the BOTOX<sup>®</sup> Cosmetic Group Compared to the Placebo Group**

Adverse Events by Body System	Percent of Patients Reporting Adverse Events	
	BOTOX <sup>®</sup> Cosmetic (N=405) %	Placebo (N=130) %
Overall	44	42
Body as a Whole		
Pain in Face	2	1
Skin and Appendages		
Skin Tightness	1	0
Digestive System		
Nausea	3	2
Dyspepsia	1	0
Tooth Disorder	1	0
Special Senses		
Blepharoptosis	3	0
Musculoskeletal System		
Muscle Weakness	2	0
Cardiovascular		
Hypertension	1	0

### **Immunogenicity**

Treatment with **BOTOX<sup>®</sup> Cosmetic** may result in the formation of neutralizing antibodies that may reduce the effectiveness of subsequent treatments with **BOTOX<sup>®</sup> Cosmetic** by inactivating the biological activity of the toxin. The rate of formation of neutralizing antibodies in patients receiving **BOTOX<sup>®</sup> Cosmetic** has not been well studied.

The critical factors for neutralizing antibody formation have not been well characterized. The results from some studies suggest that botulinum toxin injections at more frequent intervals or at higher doses may lead to greater incidence of antibody formation. The potential for antibody formation may be minimized by injecting the lowest effective dose given at the longest feasible intervals between injections.

### **Postmarketing Experience**

Transient ptosis, the most frequently reported complication, has been reported in the literature in approximately 5% of patients. There has been a single report of diplopia, which resolved completely in three weeks.

The following other adverse reactions have been identified since the drug has been marketed: abdominal pain; blurred vision; brachial plexopathy; decreased hearing; diarrhea; ear noise; erythema multiforme; fever; focal facial paralysis; glaucoma; localized numbness; loss of appetite; malaise; myalgia; myasthenia gravis; pruritus; psoriasiform eruption; retinal vein occlusion; sweating; syncope; vertigo with nystagmus; and vomiting.

Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to botulinum toxin.

### **Reporting Adverse Events**

Adverse events following use of **BOTOX<sup>®</sup> Cosmetic** should be reported to the Pharmacovigilance Department, Allergan Inc. (1-800-433-8871). Adverse events may also be reported to the U.S. Department of Health and Human Services (DHHS) Adverse Event Reporting System. Report forms and reporting requirement information can be obtained from Adverse Event Reporting System (AERS) through a toll free number 1-800-822-7967.

### **Overdosage**

Excessive doses of **BOTOX<sup>®</sup> Cosmetic** may be expected to produce neuromuscular weakness with a variety of symptoms. Respiratory support may be required where excessive doses cause paralysis of respiratory muscles. In the event of overdose, the patient should be medically monitored for symptoms of excessive muscle weakness or muscle paralysis (see **WARNINGS** and **PRECAUTIONS**). Symptomatic treatment may be necessary.

Symptoms of overdose are likely not to be present immediately following injection. Should accidental injection or oral ingestion occur, the person should be medically supervised for several weeks for signs and symptoms of excessive muscle weakness or muscle paralysis.

In the event of overdose, antitoxin raised against botulinum toxin is available from the Centers for Disease Control and Prevention (CDC) in Atlanta, GA. However, the antitoxin will not reverse any botulinum toxin-induced effects already apparent by the time of antitoxin administration. In the event of suspected or actual cases of botulinum toxin poisoning, please contact your local or state Health Department to process a request for antitoxin through the CDC. If you do not receive a response within 30 minutes, please contact the CDC directly at 1-770-488-7100. More information can be obtained at <http://www.cdc.gov/ncidod/srp/drugs/drug-service.html>.

## **DOSAGE AND ADMINISTRATION**

### **For Intramuscular Injection Only**

**BOTOX<sup>®</sup> Cosmetic** is to be reconstituted only with 0.9% sterile, non-preserved saline prior to intramuscular injection. Per the dilution table below, draw up the required amount of 0.9% sterile non-preserved sodium chloride solution into a syringe to obtain a reconstituted solution at a concentration of 4 Units/0.1 mL and a total treatment dose of 20 Units in 0.5 mL. The duration of activity of **BOTOX<sup>®</sup> Cosmetic** for glabellar lines is approximately 3-4 months. The safety and

effectiveness of more frequent dosing with **BOTOX<sup>®</sup> Cosmetic** has not been clinically evaluated and is not recommended.

#### Dilution Table

Diluent Added to 100 Unit Vial (0.9% Sodium Chloride Injection Only)	Resulting Dose Units per 0.1 mL	Diluent Added to 50 Unit Vial (0.9% Sodium Chloride Injection Only)	Resulting Dose Units per 0.1 mL
2.5 mL	4 Units	1.25 mL	4 Units

Reconstituted **BOTOX<sup>®</sup> Cosmetic** should be clear, colorless, and free of particulate matter.

**BOTOX<sup>®</sup> Cosmetic** is supplied as a single use vial. The product and diluent do not contain a preservative. Once opened and reconstituted it should be stored in a refrigerator (2° to 8°C) and used within 24 hours. Discard any remaining solution. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit. Do not freeze reconstituted **BOTOX<sup>®</sup> Cosmetic**.

#### Dilution Technique

Using a 21-gauge needle and an appropriately sized syringe draw up a total of 2.5 mL/100 Unit vial or 1.25 mL/50 Unit vial of 0.9% sterile saline without a preservative. Insert the needle at a 45° angle and slowly inject into the **BOTOX<sup>®</sup> Cosmetic** vial. Discard the vial if a vacuum does not pull the diluent into the vial. Gently rotate the vial and record the date and time of reconstitution on the space on the label.

Draw at least 0.5 mL of the properly reconstituted toxin into the sterile syringe, preferably a tuberculin syringe and expel any air bubbles in the syringe barrel. Remove the needle used to reconstitute the product and attach a 30-33 gauge needle. Confirm the patency of the needle.

#### Injection Technique

Glabellar facial lines arise from the activity of the corrugator and orbicularis oculi muscles. These muscles move the brow medially, and the procerus and depressor supercillii pull the brow inferiorly. This creates a frown or “furrowed brow”. The location, size, and use of the muscles vary markedly among individuals. Lines induced by facial expression occur perpendicular to the direction of action of contracting facial muscles. An effective dose for facial lines is determined by gross observation of the patient’s ability to activate the superficial muscles injected.

In order to reduce the complication of ptosis the following steps should be taken:

- Avoid injection near the levator palpebrae superioris, particularly in patients with larger brow depressor complexes.
- Lateral corrugator injections should be placed at least 1 cm above the bony supraorbital ridge.
- Ensure the injected volume/dose is accurate and where feasible kept to a minimum.
- Do not inject toxin closer than 1 cm above the central eyebrow.

Using a 30-33 gauge needle, inject a dose of 0.1 mL into each of 5 sites, 2 in each corrugator muscle and 1 in the procerus muscle for a total dose of 20 Units. Typically the initial doses of reconstituted **BOTOX<sup>®</sup> Cosmetic** induce chemical denervation of the injected muscles one to two days after injection, increasing in intensity during the first week.



#### **HOW SUPPLIED**

**BOTOX<sup>®</sup> Cosmetic** is supplied in a single use vial in the following sizes:

50 Units: NDC 0023-3919-50

100 Units: NDC 0023-9232-01

Vials of **BOTOX<sup>®</sup> Cosmetic** have a holographic film on the vial label that contains the name “Allergan” within horizontal lines of rainbow color. In order to see the hologram, rotate the vial back and forth between your fingers under a desk lamp or fluorescent light source. (Note: the holographic film on the label is absent in the date/batch area.) If you do not see the lines of rainbow color or the name “Allergan,” do not use the product and contact Allergan for additional information at 1-800-890-4345 from 7:00 AM to 3:00 PM Pacific Time.

#### **Rx Only**

**Single use vial.**

#### **Storage**

Unopened vials of **BOTOX<sup>®</sup> Cosmetic** should be stored in a refrigerator (2° to 8°C) for up to 36 months for the 50 Units and 100 Units vials.

Administer **BOTOX<sup>®</sup> Cosmetic** within 24 hours of reconstitution; during this period reconstituted **BOTOX<sup>®</sup> Cosmetic** should be stored in a refrigerator (2° to 8°C). Reconstituted **BOTOX<sup>®</sup> Cosmetic** should be clear, colorless and free of particulate matter.

Do not use after the expiration date on the vial. All vials, including expired vials, or equipment used with the drug should be disposed of carefully as is done with all medical waste.

**Revised: XX/2011**

© 2011 Allergan, Inc.

® mark owned by Allergan, Inc.

U.S. Patents 6,974,578; 6,683,049; and 6,896,886

Manufactured by: Allergan Pharmaceuticals Ireland  
a subsidiary of: Allergan, Inc., 2525 Dupont Dr., Irvine, CA 92612

## **REFERENCE**

1. Wang YC, Burr DH, Korthals GJ, Sugiyama H. Acute toxicity of aminoglycoside antibiotics as an aid in detecting botulism. *Appl Environ Microbiol* 1984; 48:951-955.

**MEDICATION GUIDE**  
**BOTOX<sup>®</sup>**  
**BOTOX<sup>®</sup> Cosmetic**  
**(Boe-tox)**  
**(onabotulinumtoxinA)**  
**for Injection**

Read the Medication Guide that comes with **BOTOX** or **BOTOX Cosmetic** before you start using it and each time it is given to you. There may be new information. This information does not take the place of talking with your doctor about your medical condition or your treatment. You should share this information with your family members and caregivers.

**What is the most important information I should know about BOTOX and BOTOX Cosmetic?**

**BOTOX and BOTOX Cosmetic may cause serious side effects that can be life threatening. Call your doctor or get medical help right away if you have any of these problems after treatment with BOTOX or BOTOX Cosmetic:**

- **Problems swallowing, speaking, or breathing.** These problems can happen hours to weeks after an injection of **BOTOX** or **BOTOX Cosmetic** usually because the muscles that you use to breathe and swallow can become weak after the injection. Death can happen as a complication if you have severe problems with swallowing or breathing after treatment with **BOTOX** or **BOTOX Cosmetic**.
- People with certain breathing problems may need to use muscles in their neck to help them breathe. These patients may be at greater risk for serious breathing problems with **BOTOX** or **BOTOX Cosmetic**.
- Swallowing problems may last for several months. People who cannot swallow well may need a feeding tube to receive food and water. If swallowing problems are severe, food or liquids may go into your lungs. People who already have swallowing or breathing problems before receiving **BOTOX** or **BOTOX Cosmetic** have the highest risk of getting these problems.
- **Spread of toxin effects.** In some cases, the effect of botulinum toxin may affect areas of the body away from the injection site and cause symptoms of a serious condition called botulism. The symptoms of botulism include:
  - loss of strength and muscle weakness all over the body
  - double vision
  - blurred vision and drooping eyelids
  - hoarseness or change or loss of voice (dysphonia)
  - trouble saying words clearly (dysarthria)
  - loss of bladder control
  - trouble breathing
  - trouble swallowing

These symptoms can happen hours to weeks after you receive an injection of **BOTOX** or **BOTOX Cosmetic**.

These problems could make it unsafe for you to drive a car or do other dangerous activities. See "What should I avoid while receiving **BOTOX** or **BOTOX Cosmetic**?"

There has not been a confirmed serious case of spread of toxin effect away from the injection site when **BOTOX** has been used at the recommended dose to treat chronic migraine, severe underarm sweating, blepharospasm, or strabismus, or when **BOTOX Cosmetic** has been used at the recommended dose to treat frown lines.

### **What are BOTOX and BOTOX Cosmetic?**

**BOTOX** is a prescription medicine that is injected into muscles and used:

- to prevent headaches in adults with chronic migraine who have 15 or more days each month with headache lasting 4 or more hours each day.
- to treat increased muscle stiffness in elbow, wrist, and finger muscles in adults with upper limb spasticity.
- to treat the abnormal head position and neck pain that happens with cervical dystonia (CD) in adults.
- to treat certain types of eye muscle problems (strabismus) or abnormal spasm of the eyelids (blepharospasm) in people 12 years and older.

**BOTOX** is also injected into the skin to treat the symptoms of severe underarm sweating (severe primary axillary hyperhidrosis) when medicines used on the skin (topical) do not work well enough.

**BOTOX Cosmetic** is a prescription medicine that is injected into muscles and used to improve the look of moderate to severe frown lines between the eyebrows (glabellar lines) in adults younger than 65 years of age for a short period of time (temporary).

It is not known whether **BOTOX** is safe or effective in patients younger than:

- 18 years of age for treatment of chronic migraine
- 18 years of age for treatment of spasticity
- 16 years of age for treatment of cervical dystonia
- 18 years of age for treatment of hyperhidrosis
- 12 years of age for treatment of strabismus or blepharospasm

**BOTOX Cosmetic** is not recommended for use in children younger than 18 years of age.

It is not known whether **BOTOX** and **BOTOX Cosmetic** are safe or effective to prevent headaches in patients with migraine who have 14 or fewer headache days each month (episodic migraine).

It is not known whether **BOTOX** and **BOTOX Cosmetic** are safe or effective for other types of muscle spasms or for severe sweating anywhere other than your armpits.

### **Who should not take BOTOX or BOTOX Cosmetic?**

Do not take **BOTOX** or **BOTOX Cosmetic** if you:

- are allergic to any of the ingredients in **BOTOX** or **BOTOX Cosmetic**. See the end of this Medication Guide for a list of ingredients in **BOTOX** and **BOTOX Cosmetic**.
- had an allergic reaction to any other botulinum toxin product such as *Myobloc*<sup>®</sup>, *Dysport*<sup>®</sup>, or *Xeomin*<sup>®</sup>
- have a skin infection at the planned injection site

### **What should I tell my doctor before taking BOTOX or BOTOX Cosmetic?**

#### **Tell your doctor about all your medical conditions, including if you have:**

- a disease that affects your muscles and nerves (such as amyotrophic lateral sclerosis [ALS or Lou Gehrig's disease], myasthenia gravis or Lambert-Eaton syndrome). See "What is the most important information I should know about **BOTOX** and **BOTOX Cosmetic**?"
- allergies to any botulinum toxin product
- had any side effect from any botulinum toxin product in the past
- a breathing problem, such as asthma or emphysema
- swallowing problems
- bleeding problems
- plans to have surgery
- had surgery on your face
- weakness of your forehead muscles, such as trouble raising your eyebrows
- drooping eyelids
- any other change in the way your face normally looks
- are pregnant or plan to become pregnant. It is not known if **BOTOX** or **BOTOX Cosmetic** can harm your unborn baby.
- are breast-feeding or plan to breastfeed. It is not known if **BOTOX** or **BOTOX Cosmetic** passes into breast milk.

**Tell your doctor about all the medicines you take**, including prescription and nonprescription medicines, vitamins and herbal products. Using **BOTOX** or **BOTOX Cosmetic** with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received BOTOX or BOTOX Cosmetic in the past.**

Especially tell your doctor if you:

- have received any other botulinum toxin product in the last four months
- have received injections of botulinum toxin, such as *Myobloc*<sup>®</sup> (rimabotulinumtoxinB), *Dysport*<sup>®</sup> (abobotulinumtoxinA), or *Xeomin*<sup>®</sup> (incobotulinumtoxinA) in the past. Be sure your doctor knows exactly which product you received.
- have recently received an antibiotic by injection
- take muscle relaxants
- take an allergy or cold medicine
- take a sleep medicine

**Ask your doctor if you are not sure if your medicine is one that is listed above.**

Know the medicines you take. Keep a list of your medicines with you to show your doctor and pharmacist each time you get a new medicine.

### **How should I take BOTOX or BOTOX Cosmetic?**

- **BOTOX** or **BOTOX Cosmetic** is an injection that your doctor will give you.
- **BOTOX** is injected into your affected muscles or skin.
- **BOTOX Cosmetic** is injected into your affected muscles.
- Your doctor may change your dose of **BOTOX** or **BOTOX Cosmetic**, until you and your doctor find the best dose for you.
- Your doctor will tell you how often you will receive your dose of **BOTOX** or **BOTOX Cosmetic** injections.

### **What should I avoid while taking BOTOX or BOTOX Cosmetic?**

**BOTOX** and **BOTOX Cosmetic** may cause loss of strength or general muscle weakness, or vision problems within hours to weeks of taking **BOTOX** or **BOTOX Cosmetic**. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.** See "What is the most important information I should know about **BOTOX** and **BOTOX Cosmetic**?"

### **What are the possible side effects of BOTOX and BOTOX Cosmetic?**

**BOTOX** and **BOTOX Cosmetic** can cause serious side effects. See "What is the most important information I should know about **BOTOX** and **BOTOX Cosmetic**?"

### **Other side effects of BOTOX and BOTOX Cosmetic include:**

- dry mouth
- discomfort or pain at the injection site
- tiredness
- headache
- neck pain
- eye problems: double vision, blurred vision, decreased eyesight, drooping eyelids, swelling of your eyelids, and dry eyes.
- allergic reactions. Symptoms of an allergic reaction to **BOTOX** or **BOTOX Cosmetic** may include: itching, rash, red itchy welts, wheezing, asthma symptoms, or dizziness or feeling faint. Tell your doctor or get medical help right away if you are wheezing or have asthma symptoms, or if you become dizzy or faint.

Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of **BOTOX** and **BOTOX Cosmetic**. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **General information about BOTOX and BOTOX Cosmetic:**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide.

This Medication Guide summarizes the most important information about **BOTOX** and **BOTOX Cosmetic**. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about **BOTOX** and **BOTOX Cosmetic** that is written for healthcare

professionals. For more information about **BOTOX** and **BOTOX Cosmetic** call Allergan at 1-800-433-8871 or go to [www.botox.com](http://www.botox.com).

**What are the ingredients in BOTOX and BOTOX Cosmetic?**

Active ingredient: botulinum toxin type A

Inactive ingredients: human albumin and sodium chloride

**Issued: 10/2010**

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Manufactured by: Allergan Pharmaceuticals Ireland

a subsidiary of: Allergan, Inc.

2525 Dupont Dr.

Irvine, CA 92612

© 2010 Allergan, Inc.

® mark owned by Allergan, Inc.

U.S. Patents 6,974,578; 6,683,049; and 6,896,886

*Myobloc*® is a registered trademark of Solstice Neurosciences, Inc.

*Dysport*® is a registered trademark of Ipsen Biopharm Limited Company.

*Xeomin*® is a registered trademark of Merz Pharma GmbH & Co KGaA.