



50094538

NDC 41167 4310 4

Allegra-D[®]

Allergy & Congestion

fexofenadine HCl 60 mg/antihistamine
 pseudoephedrine HCl 120 mg/nasal decongestant
 Extended Release Tablets

Indoor and Outdoor Allergies

NON-DROWSY
 Original Prescription Strength

12 Hour

KEEP OR: _____

- ✓ Nasal Congestion
- ✓ Sinus Congestion & Pressure
- ✓ Runny Nose
- ✓ Sneezing
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

20 Tablets

Allegra-D[®]
Allergy & Congestion

The makers of Allegra[®] do not make store brand products. The trade dress of this Allegra[®] package is subject to trademark protection.
 Dist. By: Chatham, Inc. (part of the sanofi-aventis Group), Chattanooga, TN 37409-0219 ©2010



50094538-CS



0 41167 43104 7

NON-DROWSY
 Original Prescription Strength

Allegra-D[®]
Allergy & Congestion

12 Hour

Drug Facts (continued)

Warnings

- If you have ever had an allergic reaction to this product or any of its ingredients, do not use.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- If you have difficulty swallowing, ask a doctor before use if you have:
 - heart disease
 - thyroid disease
 - glaucoma
 - high blood pressure
 - kidney disease, your doctor should determine if you need a different dose.
- kidney disease, your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

Stop use and ask a doctor if

- symptoms do not improve within 7 days or are accompanied by a fever.
- an allergic reaction to this product occurs. Seek medical help right away.

Do not get nervous, dizzy, or sleepless

Drug Facts (continued)

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - sneezing
 - runny nose
 - itching of the nose or throat
- temporarily relieves sinus congestion and pressure
- temporarily restores nasal breathing through the nose

Purpose

- Antihistamine
- Nasal decongestant

Active ingredients (in each tablet)

- Fexofenadine HCl 60 mg
- Pseudoephedrine HCl 120 mg

Drug Facts (continued)

Directions

- If pregnant or breast-feeding, ask a health professional before use.
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Other information

- safety sealed; do not use if cap is opened or if individual blister units are torn or opened.
- Store between 20° and 25°C (68° and 77°).

Inactive ingredients

carbamate wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose, polyethylene glycol, pregelatinized starch, stearic acid

Questions or comments?

call toll-free 1-800-633-1610 or www.allegra.com

Drug Facts (continued)

Warnings

- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- If you have difficulty swallowing, ask a doctor before use if you have:
 - heart disease
 - thyroid disease
 - glaucoma
 - high blood pressure
 - kidney disease, your doctor should determine if you need a different dose.
- kidney disease, your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

Stop use and ask a doctor if

- symptoms do not improve within 7 days or are accompanied by a fever.
- an allergic reaction to this product occurs. Seek medical help right away.


Do not get nervous, dizzy, or sleepless






Allegra-D 12hr 5ct Blister Pack







Allegra-D 12hr 5ct Blister Pack

Removed "12 Hour" from the name here.

Positioned "12 Hour" here in alignment with the approved Allegra Allergy blister label.

<p>Allegra-D® Allergy & Congestion pseudoephedrine HCl 120 mg fexofenadine HCl 60 mg 12 Hour Extended Release Tablets Dist. By: Chatterm, Inc. Chattanooga, TN 37409-0219 PUSH TABLET THROUGH FOIL</p> <p>Lot 00000 Exp MM/YYYY</p>	<p>50098529-CS</p>	<p>Allegra-D® Allergy & Congestion pseudoephedrine HCl 120 mg fexofenadine HCl 60 mg 12 Hour Extended Release Tablets Dist. By: Chatterm, Inc. Chattanooga, TN 37409-0219 PUSH TABLET THROUGH FOIL</p> <p>Lot 00000 Exp MM/YYYY</p>
<p>Allegra-D® Allergy & Congestion pseudoephedrine HCl 120 mg fexofenadine HCl 60 mg 12 Hour Extended Release Tablets Dist. By: Chatterm, Inc. Chattanooga, TN 37409-0219 PUSH TABLET THROUGH FOIL</p> <p>Lot 00000 Exp MM/YYYY</p>	<p>Allegra-D® Allergy & Congestion pseudoephedrine HCl 120 mg fexofenadine HCl 60 mg 12 Hour Extended Release Tablets Dist. By: Chatterm, Inc. Chattanooga, TN 37409-0219 PUSH TABLET THROUGH FOIL</p> <p>Lot 00000 Exp MM/YYYY</p>	<p>Allegra-D® Allergy & Congestion pseudoephedrine HCl 120 mg fexofenadine HCl 60 mg 12 Hour Extended Release Tablets Dist. By: Chatterm, Inc. Chattanooga, TN 37409-0219 PUSH TABLET THROUGH FOIL</p> <p>Lot 00000 Exp MM/YYYY</p>

Removed "12 Hour"

Title for Continued Panel:
 8 point Helvetica Bold Italic

8 point Helvetica Regular

Headings:
 8 point Helvetica Bold italic, left justified

Bullet:
 5 point solid square

Subheadings:
 6 point Helvetica Bold left justified

0.5 point hairline

2.5 point barline

table format for 3 or more dosages

Body Text:
 6 point Helvetica Regular left justified, 6.5 leading

Drug Facts (continued)

Warnings
Do not use
 if you have ever had an allergic reaction to this product or any of its ingredients if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 if you have difficulty swallowing

Ask a doctor before use if you have
 heart disease thyroid disease glaucoma
 high blood pressure diabetes
 trouble urinating due to an enlarged prostate gland
 kidney disease. Your doctor should determine if you need a different dose.

When using this product
do not take more than directed
 do not take at the same time as aluminum or magnesium antacids
 the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

Stop use and ask a doctor if
 an allergic reaction to this product occurs. Seek medical help right away, symptoms do not improve within 7 days or are accompanied by a fever you get nervous, dizzy, or sleepless

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 do not divide, crush, chew or dissolve the tablet; swallow tablet whole
 adults and children 12 years of age and over take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
 children under 12 years of age do not use
 adults 65 years of age and older ask a doctor
 consumers with kidney disease ask a doctor

Other information
 safety sealed: do not use if carton is opened or if individual blister units are torn or opened
 store between 20° and 25°C (68° and 77°F)

Inactive ingredients
 carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, stearic acid

Questions or comments?
 call toll-free 1-800-633-1610 or www.allegra.com

Removed "12 Hour"

Removed "12 Hour"

Removed "12 Hour"

30 Tablets

12 Hour

Original Prescription Strength

NON-DROWSY

Indoor and Outdoor Allergies

Extended Release Tablets
 pseudoephedrine HCl 120 mg/nasal decongestant
 fexofenadine HCl 60 mg/antihistamine

Relief of:
 ✓ Nasal Congestion & Pressure
 ✓ Sinus Congestion
 ✓ Sneezing
 ✓ Itchy, Watery Eyes
 ✓ Itchy Nose or Throat

Drug Facts

Active ingredients (in each tablet)

Fexofenadine HCl 60 mg	Antihistamine
Pseudoephedrine HCl 120 mg	Nasal decongestant

Purpose

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: runny nose, sneezing, nasal congestion, itchy, watery eyes, itching of the nose or throat, reduces swelling of nasal passages, temporarily relieves sinus congestion and pressure, temporarily restores freer breathing through the nose

Drug Facts (continued)

Uses
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: runny nose, sneezing, nasal congestion, itchy, watery eyes, itching of the nose or throat, reduces swelling of nasal passages, temporarily relieves sinus congestion and pressure, temporarily restores freer breathing through the nose

Title:
 14 point Helvetica Bold italic, left justified

Right Justified

Graphic leading to next panel

2.5 point box barline