

Reference ID: 3175052

(b) (4)

(b) (4)

Exp. Date xx/xx
Lot No. xxxxx
Cat. No. 260479 NDC 054365-400-15
ChloraPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1 ml APPLICATOR Single Use • Latex Free
WARNING. FLAMMABLE.
KEEP AWAY FROM FIRE OR FLAME.
do not drape or use ignition source (e.g., cautery, laser)
until solution is completely dry (minimum of 3 minutes
on hairless skin; up to 1 hour in hair)
APPLICATOR STERILE UNLESS SEAL IS BROKEN
CareFusion, Leawood, KS 66211
Artwork No. 6 001479 • C0112
Made in the USA of US and imported parts
Bend and Peel
Trim Area

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

ChloraPrep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

1 ml APPLICATOR

Single Use | Do Not Reuse | Latex Free

External Use Only | Professional Use Only

Cat. No. 260479 NDC 054365-400-11

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. ▶

ChloraPrep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm^2)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic.
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- USP purified water

Questions?

- www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork 6-775479• A0611 • Made in the USA of US and Imported parts

ChloraPrep. one-step

1.5ml FREPP®

ChloraPrep One-Step

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients — **Purposes**

Chlorhexidine gluconate 2% w/v ..... Antiseptic

Isopropyl alcohol 70% v/v ..... Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator horizontally with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F)
■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloraprep.com
■ call 1-800-523-0502 (M- F 8 a.m.–5 p.m. CST)

(b) (4)

(b) (4)

(b) (4)

8.500

3.469

3.500

0.625

8.500

Artwork No. 6-824299 0311

NDC 0538-4400-02

1.5 ml
FREPP®
APPLICATORS

Clear

Single Use

Latex Free

Applicator is STERILE if package is intact

20 applicators
0.05 fl. oz (1.5 ml) each

ChloraPrep®
One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
- solution contains alcohol and gives off flammable vapors
- avoid pooling solution if it is fully dries. Hair may take up to 1 hour to dry. Pat hair to flammable.
- do not drape or use ignition source (e.g., cautery, lasers) until solution is completely dry (minimum of 3 minutes on hairless skin, up to 1 hour in hair)
- do not allow solution to pool
- remove excess material from prep area

Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat above 40 °C (104 °F)

ChloraPrep®
One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
1.5 ml
FREPP® APPLICATORS
Clear

LOT XXXXXX EXP DATE: XX/XX
(printed in-line)

No Varnish

Cat. No. 260299

No Varnish / No printing area

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active Ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:
- solution contains alcohol and gives off flammable vapors
- avoid pooling solution if it is fully dries. Hair may take up to 1 hour to dry. Pat hair to flammable.
- do not drape or use ignition source (e.g., cautery, lasers) until solution is completely dry (minimum of 3 minutes on hairless skin, up to 1 hour in hair)
- do not allow solutions to pool
- remove excess material from prep area

Do not use
- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact in the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- intended to prepare skin for one applicator is approximately 2.5 in. x 2.5 in. (12 cm²)
- remove applicator from package, do not touch sponge
- hold the applicator horizontally with the sponge down. Pinch wings gently to activate the ampule and release the solution
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites (e.g., inguinal folds): use gentle repeated back-and-forth strokes for 2 minutes
- allow the solution to completely dry (minimum of 2 minutes on hairless skin, up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other Information ■ store between 15–30 °C (59–86 °F)
■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive Ingredients ■ USP purified water

Questions? ■ www.chloraprep.com
■ call 1-800-519-9053 (8:F a.m.–5 p.m. CST)

(b) (4)

2.484

24.594

ChloraPrep®
One-Step

1.5 ml
FREPP®
APPLICATORS

Clear

20 applicators

ChloraPrep®
One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
1.5 ml
FREPP® APPLICATORS
Clear

CareFusion
Leawood, KS 66211
1.800.523.0502
www.chloraprep.com
Made in the USA of US and imported parts

10.422

2.484

3.500

Reference ID: 3175052

(b) (4)

(b) (4)

To review remaining history – see Rev. G

(b) (4)

(b) (4)

2.5”

7.5”

ChloraPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1.5ml FREPP® APPLICATOR

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to surgery: Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable: Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is lammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator horizontally with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F)
■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloraprep.com
■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

See other side for additional information

Front

Back

2.5”

ChloraPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1.5ml FREPP® APPLICATOR

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

See Drug Facts for Full Labeling Information

Do Not Reuse Latex Free

External Use Only Professional Use Only

Single Use

CareFusion, Leawood, KS 66211
Artwork 6-775403F - L0411
Made in the USA of US and imported parts

Reference ID: 3175052


1 15/32
3 11/16
11/1
7/32
5 5/8
12 27/32
5 19/32
37 19/32
3 1/4
6 1/2
2 51/64
17 1/4
ChloraPrep One-Step
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
3 ml
WARNING
FLAMMABLE

6-786315 ChloraPrep 3ml Applicator lidding

[redacted]

To view remaining history - see Rev. E



[redacted]

Lot/Exp Placeholder - for reference only:

Lot # and Exp. Date are embossed on package.

Reference ID: 3175052

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

ChloraPrep® One-Step
ChloraPrep® With Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
3 ml APPLICATOR

Single Use (2) Do Not Reuse (Latex) Latex Free
External Use Only Professional Use Only
Cat. No. 260415 / 260415NS / 260415NSB NDC 054365-400-11
Cat. No. 260400 / 260400NS / 260500NS NDC 054365-400-01

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection. (11)

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use
- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. ▶

ChloraPrep® One-Step / ChloraPrep® With Tint
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm^2)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic.
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information
- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)
- The tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired. (9)

Inactive ingredients
- FD&C green #3 dye (Scrub-Teal applicators only)
- FD&C yellow #6 dye (Hi-Lite Orange applicators only)
- USP purified water

Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork 6-775400-E0611 • Made in the USA of US and imported parts


1 15/32
3 11/16
3¼
6½
2 51/64
17¼
1 1/16
5⅝
12 27/32
5 19/32
37 19/32
ChloraPrep
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
3 ml
WARNING
FLAMMABLE

(b) (4)

(b) (4) 3-mL ChloraPrep With Tint (Teal) Applicator Lidding

(b) (4)



ChloraPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
3ml APPLICATOR *Scrub Teal®*

Use

For the preparation of the patient's skin prior to surgery: Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is lammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Keep out of reach of children. f swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Important: See insert for complete Drug Facts information

Cat. No. 260425 NDC 054365-400-10
Single Use • Applicator is STERILE if package is intact

CareFusion, Leawood, KS 66211
call **1-800-523-0502** (M-F 8 a.m. - 5 p.m. CST) • www.chloraprep.com
Made in the USA of US and imported parts

Artwork No. 6-786425 • E0411

(b) (4)

Lot/Exp Placeholder - for reference only:

Lot # and Exp. Date are embossed on package.


1 15/32
3 11/16
3 1/4
6 1/2
2 51/64
17 1/4
1
1
7/32
5 5/8
12 27/32
5 19/32
37 19/32
ChloraPrep
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% w/v isopropyl alcohol (IPA)
WARNING
FLAMMABLE
3 ml
3 ml
3 ml
3 ml

(b) (4)

[redacted]

To view remaining history - see Rev. F



(b) (4)

[redacted]

Lot/Exp Placeholder - for reference only:

Lot # and Exp. Date are embossed on package.


1 15/32
3 11/16
11/1
12 27/32
12 27/32
5 19/32
37 19/32
3 1/4
6 1/2
2 51/64
17 1/4
ChloraPrep
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
10.5ml
PP
10.5ml
PP
10 5ml
PP
ChloraPrep
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
P P v P
10.5ml
PP
WARNING
FLAMMABLE
ChloraPrep
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
10.5ml
PP
10.5ml
PP
ChloraPrep
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

(b) (4)



(b) (4)

(b) (4)

Reference ID: 3175052

(b) (4)

(b) (4)

(b) (4)

Lot # and Exp. Date are embossed on the blister film at this end of the package.

ChloraPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
10.5ml APPLICATOR

Clear

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Use
For the preparation of the patient's skin prior to surgery: Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only. Flammable. Keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 8.4 in. x 8.4 in. (457 cm^2)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Important: See insert for complete Drug Facts information

Cat. No. 260700 NDC 054365-400-04
Single Use • Applicator is STERILE if package is intact

CareFusion, Leawood, KS 66211
call **1-800-523-0502** (M-F 8 a.m. - 5 p.m. CST) • www.chloraprep.com
Made in the USA of US and imported parts

Artwork No. 6-786408 • K0611

Lot/Exp Placeholder - for reference only:

Lot # and Exp. Date are embossed on package.

R 20

183

171

40

27°

9

12

54.25

10.5

57.25

9

75.25

R 8

Reference ID: 3175052

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

ChloraPrep® One-Step
ChloraPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation

10.5ml APPLICATOR

Do Not Reuse | Latex Free

External Use Only | Professional Use Only

Cat. No. 260700 / 260700NS / 260700NSB	NDC 054365-400-04
Cat. No. 260715 / 260715NS / 260715NSB	NDC 054365-400-12
Cat. No. 260725 / 260725NS / 260725NSB	NDC 054365-400-09

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection. (11)

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire: PREP CAREFULLY

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser ▶

ChloraPrep® One-Step / ChloraPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 8.4 in. x 8.4 in. (457 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necesary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F)

- avoid freezing and excessive heat above 40 °C (104 °F)
- The tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired. (9)

Inactive ingredients ■ FD&C green #3 dye (Scrub-Teal applicators only) ■ FD&C yellow #6 dye (High-Lite Orange applicators only) ■ USP purified water

Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork No. 6-640248 • 10411 • Made in the USA of US and imported parts

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients — **Purposes**

Chlorhexidine gluconate 2% w/v ..... Antiseptic

Isopropyl alcohol 70% v/v ..... Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection. (11)

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 8.4 in. x 8.4 in. (457 cm^2)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- FD&C green #3 dye
- USP purified water

Questions?

- www.chloraprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)



Reference ID: 3175052

(b) (4)



(b) (4)

(b) (4)

(b) (4)

Reference ID: 3175052

(b) (4)

ith Tint Applicator Tyvek lidding

(b) (4)

(b) (4)



Lot/Exp Placeholder - for reference only:

Lot # and Exp. Date are embossed on package.

Reference ID: 3175052


1 15/32
3 11/16
3 1/4
6 1/2
2 51/64
17 1/4
1
2
5 5/8
12 27/32
5 19/32
37 19/32
ChloraPrep
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
10.5ml
PP
WARNING
FLAMMABLE
With Tint

(b) (4)



(b) (4)

(b) (4)

Reference ID: 3175052

(b) (4) 10.5-mL ChloraPrep With Tint (Orange) Applicator Tyvek lidding

(b) (4)

(b) (4)



Lot/Exp Placeholder - for reference only:

Lot # and Exp. Date are embossed on package.

Reference ID: 3175052


ChloraPrep® One-Step
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
26 ml APPLICATOR
Clear
⚠ WARNING
FLAMMABLE
Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
• do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
• solution contains alcohol and gives off flammable vapors
• avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
• do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
• do not allow solution to pool
• remove wet materials from prep area
1
Made in the USA of US and imported parts
CareFusion, Leawood, KS 66211
Artwork No. 6-773800-L - F0811

ChloraPrep® With Tint
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
26 ml APPLICATOR
Hi-Lite Orange®
⚠ WARNING
FLAMMABLE
Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
• do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
• solution contains alcohol and gives off flammable vapors
• avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
• do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
• do not allow solution to pool
• remove wet materials from prep area
1
Made in the USA of US and imported parts
CareFusion, Leawood, KS 66211
Artwork No. 6-773815-L - F0611

ChloraPrep® With Tint
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
26 ml APPLICATOR
Scrub Teal®
⚠ WARNING
FLAMMABLE
Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
• do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
• solution contains alcohol and gives off flammable vapors
• avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
• do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
• do not allow solution to pool
• remove wet materials from prep area
1
Made in the USA of US and imported parts
CareFusion, Leawood, KS 66211
Artwork No. 6-773825-L - F0811

ChloraPrep® One-Step
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
26 ml APPLICATOR
Clear
⚠ WARNING
FLAMMABLE
Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
• do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
• solution contains alcohol and gives off flammable vapors
• avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
• do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
• do not allow solution to pool
• remove wet materials from prep area
Made in the USA of US and imported parts
CareFusion, Leawood, KS 66211
Artwork No. 6-773800-R - F0611


ChloraPrep® With Tint
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
26 ml APPLICATOR
Hi-Lite Orange®
⚠ WARNING
FLAMMABLE
Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
• do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
• solution contains alcohol and gives off flammable vapors
• avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
• do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
• do not allow solution to pool
• remove wet materials from prep area
Made in the USA of US and imported parts
CareFusion, Leawood, KS 66211
Artwork No. 6-773815-R - F0611

ChloraPrep® With Tint
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
26 ml APPLICATOR
Scrub Teal®
⚠ WARNING
FLAMMABLE
Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
• do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
• solution contains alcohol and gives off flammable vapors
• avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
• do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
• do not allow solution to pool
• remove wet materials from prep area
Made in the USA of US and imported parts
CareFusion, Leawood, KS 66211
Artwork No. 6-773825-R - F0811

Cat. No. 260800 NDC 054365-400-14
Single Use • Applicator is STERILE if package is intact

ChloraPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

26ml APPLICATOR

Clear

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire; PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hr in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm^2). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- do not use 26-ml applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloraprep.com
■ call **1-800-523-0502** (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork No. 6-787800 • H0611
Made in the USA of US and imported parts

Reference ID: 3175052

Cat. No. 260825 NDC 054365-400-05

Single Use • Applicator is STERILE if package is intact

ChloraPrep® With Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

26ml APPLICATOR

Scrub Teal®

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire; PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm^2). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- do not use 26-ml applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- FD&C green #3 dye
- USP purified water

Questions?

- www.chloraprep.com
- call **1-800-523-0502** (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork No. 6-786825 • K0411
Made in the USA of US and imported parts

Reference ID: 3175052

Cat. No. 260815 NDC 054365-400-13

Single Use • Applicator is STERILE if package is intact

ChloraPrep® With Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

26ml APPLICATOR

Hi-Lite Orange®

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire;
PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm^2). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- do not use 26-ml applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- FD&C yellow #6 dye
- USP purified water

Questions?

- www.chloraprep.com
- call **1-800-523-0502** (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 Artwork No. 6-786815 • J0411
Made in the USA of US and imported parts

Reference ID: 3175052

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in. x 8.4 in. Use a small applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

(3)

ChloraPrep® One-Step
ChloraPrep® With Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

26 ml APPLICATOR

Latex Free — Do Not Reuse

External Use Only — Professional Use Only

Cat. No. 260800 / 260800NS / 260800NSB NDC 054365-400-14
Cat. No. 260815 / 260815NS / 260815NSB NDC 054365-400-13
Cat. No. 260825 / 260825NS / 260825NSB NDC 054365-400-05

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

(11)

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

▶

ChloraPrep® One-Step/ChloraPrep With Tint
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears,and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm^2).
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) ■ The tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.

(9)

Inactive ingredients ■ FD&C green #3 dye (Scrub-Teal applicators only) ■ FD&C yellow #6 dye (Hi-Lite Orange applicators only) ■ USP purified water

Questions? ■ www.chloraprep.com ■ call **1-800-523-0502** (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork No. 6-775815 • H0411 • Made in the USA of US and imported parts

[redacted] - Intermediate Carton Dispenser - ChloraPrep Sepp Box artwork

[redacted]

[redacted]

[redacted]

[redacted]

[redacted]

29/32

ChloraPrep® One-Step®
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

0.67ml SEPP® APPLICATORS
Clear
200 applicators

Artwork No. 6-824449 00411

2

16 33/64

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery or injection: Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For **external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:**
- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not **use**
- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury. If permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop **use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions
- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch applicator tip
- hold the sponge tip down and pinch the barrel of the applicator only once to activate the ampule and release the antiseptic
- wet the sponge tip by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information
- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

ChloraPrep® One-Step®
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
0.67ml SEPP® APPLICATORS
Clear

ChloraPrep® One-Step®
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
0.67ml SEPP® APPLICATORS
Clear

CareFusion
Leawood, KS 66211
1.800.523.0502
www.chloraprep.com
Made in the USA of US and imported parts

NDC 054365-400-03
0.67ml SEPP® APPLICATORS
Clear

Single Use

Latex Free

Applicator is STERILE if package is intact

200 applicators
0.02 fl. oz. (0.67 ml) each

ChloraPrep® One-Step®
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation



⚠ WARNING
FLAMMABLE
Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hr in hair)
- do not allow solution to pool
- remove wet material from prep area

Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat above 40 °C (104 °F)

ChloraPrep® One-Step®
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
0.67ml SEPP® APPLICATORS
Clear

LOT XXXXXX (printed in-line) EXP DATE: XX/XXXX
No Varnish

* +H98526044921.*

Cat. No. 260449 ©2009 CareFusion. All rights reserved. ChloraPrep is a registered trademark of CareFusion or one of its subsidiaries.

No Varnish / No printing area

5 1/16

1 39/64

2 23/64

8 3/16

8 7/32

3 1/4

1 1/16

9/32

Reference ID: 3175052

(b) (4)

(b) (4)

(b) (4)

(b) (4)


Cat No. 260449
ChloraPrep®
SEPP®
2% w/v chlorhexidine gluconate (CHG) and 70% w/v isopropyl alcohol (IPA)
0.67 ml
APPLICATOR
WARNING.
FLAMMABLE.
KEEP AWAY FROM FIRE OR FLAME.
do not use ignition source (cautery, laser) unless solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
Applicator sterile unless seal is broken
CareFusion
Leawood, KS 66211
Made in the USA of US and imported parts
www.chloraprep.com
Lot No: xxxxx
Exp.: xx/xxxx
M0611

(b) (4)

(b) (4) ChloraPrep 2% Sepp Applicator, Non-Sterile - Folded Drug Facts Insert

(b) (4)

2.5"

7.5"

ChloraPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
0.67ml SEPP® APPLICATOR

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v.	Antiseptic
Isopropyl alcohol 70% v/v.	Antiseptic

Use for the preparation of the patient's skin prior to surgery or injection: Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is lammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch applicator tip
- hold the sponge tip down and pinch the barrel of the applicator only once to activate the ampule and release the antiseptic
- wet the sponge tip by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloraprep.com ■ call **1-800-523-0502** (M-F 8 a.m.–5 p.m. CST)

See other side for additional information

Front

Back

ChloraPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
0.67ml SEPP® APPLICATOR

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

See Drug Facts for Full Labeling Information

Do Not Reuse Latex Free

External Use Only Professional Use Only

Single Use

CareFusion Leawood, KS 66211
Artwork 6-775449F • L0411
Made in the USA of US and imported parts

7.5"

(b) (4)

Reference ID: 3175052



WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients — **Purposes**

Chlorhexidine gluconate 2% w/v ..... Antiseptic

Isopropyl alcohol 70% v/v ..... Antiseptic

Use for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove all wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm^2)
- tear pouch at side notch to reveal applicator handle. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

USP purified water

Questions?

- www.chloraprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

SING
SWABST
APPLICAT

Clear

LOT XX

Cat. No. 260100

©2009 CareFusion, All righ

(b) (4)

(b) (4)

(b) (4)

(b) (4)

(b) (4)

(b) (4)

1

1½

2 3 1/64

5 5/8

1 3 21/64

1 6 3/64

3/4

6 4 6

3 3 1/32

20 2 3/32

48 applicators

1.75ml SINGLE SWABSTICK APPLICATORS

Clear

ChloraPrep one-Step

2% w/v chlorhexidine gluconate (CHG) and 70% w/v isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

Artwork No. 8-824100-00111

ChloraPrep One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

1.75ml SINGLE SWABSTICK APPLICATORS

Clear

NDC 054365-400-07

1.75ml SINGLE SWABSTICK APPLICATORS

Clear

Single Use

Latex Free

48 applicators

0.06 fl. oz (1.75 ml) each

ChloraPrep One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

⚠ WARNING

FLAMMABLE

Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas that may take up to 1 hour to dry, but hair is flammable
- do not drape or use gel/liquid (e.g., surgery lotion) until solution is completely dry (minimum of 3 minutes on hairless skin, up to 1 hour in hair)
- do not allow solution to pool
- remove excess material from prep area

Store between 15-30 °C (59-86 °F)

Avoid freezing and excessive heat above 40 °C (104 °F)

ChloraPrep One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

1.75ml SINGLE SWABSTICK APPLICATORS

Clear

CareFusion

Lenexa, KS 66211

1.800.523.0502

www.chloraprep.com

Made in the USA of US and imported parts

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active Ingredients — **Purposes**

Chlorhexidine gluconate 2% w/v ..... Antiseptic

Isopropyl alcohol 70% v/v ..... Antiseptic

Use for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infections.

Warnings

For external use only. Flammable, keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas that may take up to 1 hour to dry, but hair is flammable
- do not drape or use gel/liquid (e.g., surgery lotion) until solution is completely dry (minimum of 3 minutes on hairless skin, up to 1 hour in hair)
- do not allow solution to pool
- remove excess material from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for tracheal puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious eye damage if permitted to enter and remain. If contact occurs, rinse with cool water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well-ventilated area
- clean test treatment area for one application in approximately a 2½ in. x 2½ in. (42 cm²)
- tear pouch at slits notch to reveal applicator handle. Do not touch foam applicator tip. Place foam tip side down on the treatment area, completely wet the treatment area in the direction
- dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- allow the solution to completely dry (minimum of 3 minutes on hairless skin, up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15-30 °C (59-86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)

LOT XXXXXX EXP DATE: XX/XX (printed by firm)

+H98526010025

CSL No. 260100

©2014 CareFusion. All rights reserved. ChloraPrep is a registered trademark of CareFusion or one of its subsidiaries.

Reference ID: 3175052

(b) (4) Single Swabstick Foil Pouch

(b) (4)

(b) (4)

(b) (4)

5.5"

4.5"

ChloraPrep® Single Swabstick

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1.75ml APPLICATOR

Single Use
Latex Free

Cat. No. 260100
NDC 054365-400-07

See Drug Facts for full labeling information
Made in the USA of US and imported parts

1.25" 1.625"

Manufactured by:
CareFusion
Leawood, KS 66211
1-800-523-0502
(M-F 8 a.m. - 5 p.m. CST)
www.chloraprep.com

Artwork No. 6-800101
H061

Tear Here

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
  Wet hair is flammable
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

Lot # and Exp. Date are embossed on package.

(b) (4)

Lot/Exp Placeholder - for reference only:

Lot # and Exp. Date are embossed on package.

Reference ID: 3175052

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

ChloraPrep® Single Swabstick
ChloraPrep® Triple Swabstick

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1.75ml APPLICATOR
5.25ml APPLICATOR

Latex Free | Do Not Reuse
Single Use | External Use Only | Professional Use Only

Cat. No. 260100 / 260100B NDC 054365-400-07
Cat. No. 260103 / 260103B NDC 054365-400-08

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery: Helps to reduce bacteria that potentially can cause skin infection. (14)

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser ▶

ChloraPrep® Single Swabstick/ChloraPrep® Triple Swabstick
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

When using this product keep out of eyes, ears,and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area using
  - one single swabstick applicator is approximately 2.5 in. x 2.5 in. (42 cm^2)
  - three swabsticks sequentially is approximately 5 in. x 5 in. (160 cm^2)
- tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork No. 6-656900 • L0411 • Made in the USA of US and imported parts



WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients — **Purposes**

Chlorhexidine gluconate 2% w/v ..... Antiseptic

Isopropyl alcohol 70% v/v ..... Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove all wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area using three swabsticks sequentially is approximately 5 in x 5 in (160 cm^2)
- tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicators after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- USP purified water

Questions?

- www.chloraprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

TRIP
SWABST

Clear

LOT



Reference ID: 3175052

(b) (4)

(b) (4)

5.5"

5"

ChloraPrep® Triple Swabstick

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

5.25ml APPLICATOR

Single Use
Latex Free
Cat. No. 260103 NDC 054365-400-08

See Drug Facts for full labeling information
Made in the USA of US and imported parts

2"

2.375"

Single use only

Manufactured by:
CareFusion
Leawood, KS 66211
1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST)
www.chloraprep.com

Artwork No. 6-800104 G0611

Tear Here

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

Directions:

- Tear pouch at notch.
- Pull across package to expose the ends of the swabsticks.
- Do not touch foam when removing swabsticks from pouch.
- Place foam flat side down on treatment area.

Lot # and Exp. Date are embossed on package film

.95" From left edge of live area to point of arrow for tear perforation

Lot/Exp Date placeholder - for reference only:

Lot # and Exp. Date are embossed on package film

Reference ID: 3175052

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER
08/16/2012