

NDC 054365-400-15

1 ml APPLICATORS

Clear

Single Use
Latex Free


Applicator is STERILE
if package is intact

20 applicators
0.03 fl. oz. (1 ml) each

**CloraPrep[®]
One-Step**

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

Patient Preoperative
Skin Preparation



**WARNING
FLAMMABLE**

Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas.
- Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hr in hair)
- do not allow solution to pool
- remove wet material from prep area

Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat
above 40 °C (104 °F)

**CloraPrep[®]
One-Step**

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

1 ml APPLICATORS

Clear

LOT XXXXXX EXP DATE: XXXX
(printed in-line)



Call No. 250479
©2009 CareFusion. All rights reserved. CloraPrep is a registered trademark of CareFusion or one of its subsidiaries.

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients
Chlorhexidine gluconate 2% w/v..... Antiseptic
Isopropyl alcohol 70% v/v..... Antiseptic

Use for the preparation of the patient's skin prior to surgery: Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:
• solution contains alcohol and gives off flammable vapors
• avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
• do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
• do not allow solution to pool
• remove wet materials from prep area

Do not use
• on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
• for lumbar puncture or in contact with the meninges
• on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Purposes

Antiseptic
Antiseptic

Directions
• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
• use in a well ventilated area
• maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²).
• remove applicator from package; do not touch sponge
• hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
• wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
• completely wet the treatment area with antiseptic
• dry surgical sites (e.g., abdomen or arm); use gentle repeated back-and-forth strokes for 30 seconds
• moist surgical sites (e.g., inguinal fold); use gentle repeated back-and-forth strokes for 2 minutes
• allow the solution to **completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
• discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

• store between 15-30 °C (59-86 °F)
• avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients • USP purified water

Questions? • www.chloraprep.com
• call 1-800-523-0602 (M-F 8 a.m.-5 p.m. CST)

Artwork No. 6-824479 E0411

No Varnish / No printing area

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

**CloraPrep[®]
One-Step**

Patient Preoperative Skin Preparation

**CloraPrep[®]
One-Step**

1 ml APPLICATORS

Clear

20 applicators

**CloraPrep[®]
One-Step**

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

1 ml APPLICATORS

Clear

CareFusion
Leewood, KS 66211
1.800.523.0502
www.chloraprep.com
Made in the USA of US and imported parts

0.625

3.500

2.484

24.594

2.484

8.500

10.422

(b) (4)

Exp. Date xx/xx
Lot No. xxxxx

Cat. No. 260479 NDC 054365-400-15

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1 ml APPLICATOR Single Use • Latex Free

 **WARNING. FLAMMABLE.**
KEEP AWAY FROM FIRE OR FLAME.
do not drape or use ignition source (e.g., cautery, laser)
until solution is completely dry (minimum of 3 minutes
on hairless skin; up to 1 hour in hair)

APPLICATOR STERILE UNLESS SEAL IS BROKEN
CareFusion, Leawood, KS 66211
Artwork No. 6 001479 - C012
Made in the USA of US and imported parts

Bend and Peel

Trim Area

⚠ WARNING
FLAMMABLE

Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

1 ml APPLICATOR

Single Use Do Not Reuse Latex Free

External Use Only Professional Use Only

Cat. No. 260479 NDC 054365-400-11

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v.....	Purposes
Isopropyl alcohol 70% v/v.....	Antiseptic
	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection. (11)

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. ▶

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic.
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- USP purified water

Questions?

- www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork 6-775479• A0611 • Made in the USA of US and Imported parts

Chloraprep[®] One-Step[®]

1.5 ml
FREPP[®]

Chloraprep[®]
One-Step[®]

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v Antiseptic
Isopropyl alcohol 70% v/v Antiseptic

Purposes

..... Antiseptic
..... Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator horizontally with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.


Other information ■ store between 15–30 °C (59–86 °F)
■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloraprep.com
■ call 1-800-523-0502 (M- F 8 a.m.–5 p.m. CST)

(1)

(5)
(4)



NDC 054365-400-02
1.5ml
FREPP[®]
 APPLICATORS
 Clear

Single Use
 Latex Free
 Applicator is STERILE
 if package is intact

20 applicators
 0.05 fl. oz (1.5 ml) each

ChloraPrep[®] One-Step[®]
 2% w/v chlorhexidine gluconate (CHX)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative
 Skin Preparation

WARNING
FLAMMABLE

Keep away from fire or flame.
 To reduce risk of fire, PREP CAREFULLY:
 • solution contains alcohol and gives off flammable vapors
 • avoid getting solution on hairy areas. Hair may take up to 1 hour to dry. **Hair is flammable.**
 • do not drape or use (preparation) on hairy areas until solution is completely dry (minimum of 3 minutes on hairy areas) up to 1 hour in hair
 • do not allow solution to pool
 • remove all material from prep area

Store between 15-30 °C (59-86 °F)
 Avoid freezing and excessive heat
 above 40 °C (104 °F)

Article No. 6-624299-0511

ChloraPrep[®] One-Step[®]
 2% w/v chlorhexidine gluconate (CHX)
 and 70% v/v isopropyl alcohol (IPA)
1.5ml
FREPP[®] APPLICATORS
 Clear

LOT XXXXX (printed in line)
 Exp. Date: XXXX
 No Variants

1 9 8 5 2 6 0 2 9 9 2 0

Art. No. 250299

No Variants/No printing area

WARNING, FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts	Purpose(s)
Active Ingredients Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	Antiseptic Antiseptic
Use: For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infections.	
Warnings: For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY: • solution contains alcohol and gives off flammable vapors • avoid getting solution on hairy areas. Hair may take up to 1 hour to dry. Hair is flammable. • do not drape or use (preparation) on hairy areas until solution is completely dry (minimum of 3 minutes on hairy areas) up to 1 hour in hair • do not allow solution to pool • remove all material from prep area	
DO NOT USE: • on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol • for lacerations or on sores in the prep area • on broken skin or on areas of general skin abrasion • when using this product, keep out of eyes, ears, and mouth. May cause stinging or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. Stop use and seek a doctor if: Irritation, sensitization, or allergic reaction occurs. There may be signs of a serious condition.	
Directions: • use in this area in preparing infants or infants under 2 months of age. These products may cause irritation or chemical burns. • clean skin with water first • minimal prep area for one applicator is approximately 2.5 in. x 2.5 in. (6.3 cm) • remove applicator from package. Do not touch sponge. • hold the applicator horizontally with the sponge side facing right. Pinch the edge of the sponge and release the applicator. • wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin. • completely wet the treatment area with antiseptic. • dry surgical area (e.g., abdomen or arm). Use gentle repeated back-and-forth strokes for 30 seconds. • avoid surgical sites (e.g., inguinal fold). Use gentle repeated back-and-forth strokes for 2 minutes. • allow the solution to completely dry (minimum of 3 minutes on hairy skin, up to 1 hour in hair). Do not blot or wipe away. • discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.	
Other Information: • store between 15-30 °C (59-86 °F) • avoid freezing and excessive heat above 40 °C (104 °F)	
Inactive Ingredients: USP purified water	
Questions? • www.chloraprep.com • call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)	

ChloraPrep[®] One-Step[®]
 2% w/v chlorhexidine gluconate (CHX)
 and 70% v/v isopropyl alcohol (IPA)
1.5ml
FREPP[®] APPLICATORS
 Clear

Carefusion
 Leawood, KS 66211
 1-800-523-0502
 www.chloraprep.com
 Made in the USA of US and imported parts

2.484

3.500

0.625

8.500

3.500

8.500

3.469

24.594

2.484

10.422

Exp. Date xx/xx
Lot No. xxxxx

72 mm

Cat. No. 260299 NDC 054365-400-02

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1.5ml FREPP® APPLICATOR
Single Use • Latex Free

 **WARNING. FLAMMABLE.**
KEEP AWAY FROM FIRE OR FLAME.

**do not drape or use ignition source (e.g., cautery, laser)
until solution is completely dry (minimum of 3 minutes
on hairless skin; up to 1 hour in hair)**

APPLICATOR STERILE UNLESS SEAL IS BROKEN

CareFusion, Leawood, KS 66211
Artwork No. 6-000299 • K0611
Made in the USA of US and imported parts

Bend and Peel

40 mm

(b) (4)

2.5"

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1.5ml FREPP® APPLICATOR

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v.....	Antiseptic
Isopropyl alcohol 70% v/v.....	Antiseptic

Use For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable: Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator horizontally with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F)
■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloroprep.com
■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

Reference ID: 3175064
See other side for additional information

Front

Back

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1.5ml FREPP® APPLICATOR

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

⚠ WARNING FLAMMABLE



Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

See Drug Facts for Full Labeling Information



Do Not Reuse




Latex Free

External Use Only

Professional Use Only

Single Use

CareFusion, Leawood, KS 66211
Artwork 6-775403F - L0411
Made in the USA of US and imported parts



⚠ WARNING
FLAMMABLE

Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

ChloroPrep® One-Step ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
3 ml APPLICATOR

Single Use  Do Not Reuse  Latex Free

External Use Only Professional Use Only
Cat. No. 260415 / 260415NS / 260415NSB NDC 054365-400-11
Cat. No. 260400 / 260400NS / 260500NS NDC 054365-400-01

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v.....	Antiseptic
Isopropyl alcohol 70% v/v.....	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection. (11)

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. ▶

ChloroPrep® One-Step / ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic.
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)
- The tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired. (9)

Inactive ingredients


- FD&C green #3 dye (Scrub-Teal applicators only)
- FD&C yellow #6 dye (Hi-Lite Orange applicators only)
- USP purified water

Questions? ■ www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

P W A d

1 15/32

3 1 1/16



A G A B P WAY O O A

3 1/4

1 7/4

6 1/2

2 5 1/6 4

1 1/4



2

5 5/8

1 2 7/32



5 1 9/32

3 7 1 9/32



$1\frac{15}{32}$

ChloroPrep[®]
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)




3 ml
PP
g*

3 ml
P
g*


3 ml
PP
g*

3 ml
PP
g*

ChloroPrep[®]
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)



ChloroPrep[®]
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)



$3\frac{1}{16}$

1

1

$\frac{7}{32}$

$5\frac{5}{8}$

$12\frac{7}{32}$

$5\frac{9}{32}$

$37\frac{9}{32}$


$3\frac{1}{4}$

$6\frac{1}{2}$

$2\frac{5}{16}$

$17\frac{1}{4}$

P W g A d




Lot # and Exp. Date are embossed on the blister film at this end of the package.

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.



Lot/Exp Placeholder - for reference only:
Lot # and Exp. Date are embossed on package.

$1\frac{15}{32}$



$3\frac{1}{4}$

$1\ 7\frac{1}{4}$

$6\frac{1}{2}$

$2\ 5\frac{1}{64}$

$3\ 1\frac{1}{16}$

$1\frac{1}{16}$

$12\ 2\frac{7}{32}$

$12\ 2\frac{7}{32}$

$5\ 1\frac{9}{32}$


$37\ 1\frac{9}{32}$

(b) (4)

10.5-ml ChloroPrep One-Step In-Line Mold Labels




(b) (4)



(b) (4)

(b) (4)



**WARNING
FLAMMABLE**



- Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:
- solution contains alcohol and gives off **flammable vapors**
 - avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
 - do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
 - do not allow solution to pool
 - remove wet material from prep area

**ChloraPrep® One-Step
ChloraPrep® With Tint**

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
10.5ml APPLICATOR

Do Not Reuse

Latex Free

External Use Only Professional Use Only

Cat. No. 260700 / 260700NS / 260700NSB NDC 054365-400-04
Cat. No. 260715 / 260715NS / 260715NSB NDC 054365-400-12
Cat. No. 260725 / 260725NS / 260725NSB NDC 054365-400-09

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v.....Antiseptic
Isopropyl alcohol 70% v/v.....Antiseptic

Purposes
.....Antiseptic
.....Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection. (11)

Warnings

- For external use only. Flammable, keep away from fire or flame. To reduce risk of fire: PREP CAREFULLY
- solution contains alcohol and gives off **flammable vapors**
 - avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
 - do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
 - do not allow solution to pool
 - remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

ChloraPrep® One-Step / ChloraPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 8.4 in. x 8.4 in. (457 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)
- The tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired. (9)

Inactive ingredients

■ FD&C green #3 dye (Scrub-Teal applicators only) ■ FD&C yellow #6 dye (High-Lite Orange applicators only) ■ USP purified water

Questions?

■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork No. 6-640248 • 10411 • Made In the USA of US and imported parts

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts	
Active ingredients Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	Purposes Antiseptic Antiseptic
Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection. (11)	
Warnings For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY: <ul style="list-style-type: none"> solution contains alcohol and gives off flammable vapors. avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) do not allow solution to pool remove wet materials from prep area Do not use <ul style="list-style-type: none"> on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol for lumbar puncture or in contact with the meninges on open skin wounds or as a general skin cleanser When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions <ul style="list-style-type: none"> use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. use in a well ventilated area maximal treatment area for one applicator is approximately 8.4 in. x 8.4 in. (457 cm²) remove applicator from package; do not touch sponge hold the applicator with the sponge down. Pinch wings <i>only once</i> to activate the ampule and release the antiseptic. wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin completely wet the treatment area with antiseptic dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes do not allow solution to pool; tuck prep towels to absorb solution, and then remove allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. Other information <ul style="list-style-type: none"> store between 15–30 °C (59–86 °F) avoid freezing and excessive heat above 40 °C (104 °F) Inactive ingredients <ul style="list-style-type: none"> FD&C green #3 dye USP purified water Questions? <ul style="list-style-type: none"> www.chloraprep.com call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST) 	

Chloraprep
W
2% w/v chlorhexidine
and 70% v/v isopropyl
Patient Preparation


10
APPLICATORS

Scrub Teal

25 applicators


10.5-ml ChloroPrep With Tint In-Line Mold Labels

(b) (4)



(b) (4)


(b) (4)



Lot/Exp Placeholder - for reference only:
 Lot # and Exp. Date are embossed on package.

$1\frac{15}{32}$

Chloraprep
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)




10.5ml
PP
H L t O *


Chloraprep
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
PP
10.5ml
H L t O *

Chloraprep
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
PP
10.5ml
H L t O *


10.5ml
PP
i t O *



Chloraprep
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
PP



Chloraprep
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
PP
10.5ml
i t O *



$3\frac{1}{16}$

11

$5\frac{5}{8}$

$12\frac{27}{32}$

$5\frac{19}{32}$

$37\frac{19}{32}$

$2\frac{51}{64}$

$6\frac{1}{2}$


$3\frac{1}{4}$

$17\frac{1}{4}$

P g A d

(b) (4)

10.5-ml ChloroPrep With Tint In-Line Mold Labels



(b) (4)

(b) (4)

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

26 ml APPLICATOR

Clear

**⚠ WARNING
FLAMMABLE**



Keep away
from fire
or flame.

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area



Made in the USA of US and imported parts
CareFusion, Leawood, KS 66211
Artwork No. 6-773800-L-F0811

ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

26 ml APPLICATOR

Hi-Lite Orange®

**⚠ WARNING
FLAMMABLE**



Keep away
from fire
or flame.

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area



Made in the USA of US and imported parts
CareFusion, Leawood, KS 66211
Artwork No. 6-773815-L-F0811

ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

26 ml APPLICATOR

Scrub Teal®

**⚠ WARNING
FLAMMABLE**



Keep away
from fire
or flame.

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area



Made in the USA of US and imported parts
CareFusion, Leawood, KS 66211
Artwork No. 6-773825-L-F0811


Made in the USA of US and imported parts
CareFusion, Leewood, KS 66211
Artwork No. 6-773800-R - F0611

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

26ml APPLICATOR *Clear*

**⚠ WARNING
FLAMMABLE**

 Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

①


Made in the USA of US and imported parts
CareFusion, Leewood, KS 66211
Artwork No. 6-773815-R - F0611

ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

26ml APPLICATOR *Hi-Lite Orange®*

**⚠ WARNING
FLAMMABLE**

 Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

①


Made in the USA of US and imported parts
CareFusion, Leewood, KS 66211
Artwork No. 6-773825-R - F0811

ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

26ml APPLICATOR *Scrub Teal®*

**⚠ WARNING
FLAMMABLE**

 Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

①

Cat. No. 260800 NDC 054365-400-14
Single Use • Applicator is STERILE if package is intact

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
26 ml APPLICATOR

Clear

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire; PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hr in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm²). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- do not use 26-ml applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloroprep.com
■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork No. 6-787800 • H0671
Made in the USA of US and imported parts

Cat. No. 260825 NDC 054365-400-05

Single Use • Applicator is STERILE if package is intact

ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG)

and 70% v/v isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

26ml APPLICATOR

Scrub Teal®

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients Purposes

Chlorhexidine gluconate 2% w/v Antiseptic
Isopropyl alcohol 70% v/v Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire;

PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm²). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- do not use 26-ml applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- FD&C green #3 dye
- USP purified water

Questions? ■ www.chloroprep.com
■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork No. 6-786825 • K0411
Made in the USA of US and imported parts

2

11

Cat. No. 260815 NDC 054365-400-13
Single Use • Applicator is STERILE if package is intact

ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

26ml APPLICATOR

Hi-Lite Orange®

WARNING: FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire;

PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm²). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- do not use 26-ml applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- FD&C yellow #6 dye
- USP purified water

Questions?

- www.chloroprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 Artwork No. 6-786815 • J0411
Made in the USA of US and imported parts

WARNING
FLAMMABLE



- Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:
- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in. x 8.4 in. Use a small applicator instead.
 - solution contains alcohol and gives off **flammable vapors**
 - avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
 - do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
 - do not allow solution to pool
 - remove wet material from prep area

ChloroPrep® One-Step
ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

26ml APPLICATOR

Latex Free

Do Not Reuse

External Use Only

Professional Use Only

Cat. No. 260800 / 260800NS / 260800NSB NDC 054365-400-14
Cat. No. 260815 / 260815NS / 260815NSB NDC 054365-400-13
Cat. No. 260825 / 260825NS / 260825NSB NDC 054365-400-05

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v.....

Isopropyl alcohol 70% v/v.....

Purposes

Antiseptic

Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

ChloroPrep® One-Step/ChloroPrep With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions


- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm²).
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.


Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) ■ The tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.


Inactive ingredients ■ FD&C green #3 dye (Scrub-Teal applicators only) ■ FD&C yellow #6 dye (Hi-Lite Orange applicators only) ■ USP purified water

Questions? ■ www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

Carefusion, Leawood, KS 66211 • Artwork No. 6-775815 • H0411 • Made in the USA of US and imported parts







(b) (4)

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/vAntiseptic
Isopropyl alcohol 70% v/vAntiseptic

Purposes

Use for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove all wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or Isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- tear pouch at side notch to reveal applicator handle. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15-30 °C (59-86 °F)
■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions?

- www.chloraprep.com
- call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)

SING
SWABST
APPLICATOR

Clear


LOT XX



Cat. No. 260100
©2009 CareFusion, All rights reserved.

Single Swabstick Foil Pouch

(b) (4)

ChloroPrep® Single Swabstick
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
 1.75ml APPLICATOR

Single Use
 Latex Free
 Cat. No. 260100
 NDC 054365-400-07

See Drug Facts for full labeling information
 Made in the USA of US and imported parts



Manufactured by:
 CareFusion
 Leawood, KS 66211
 1-800-523-0502
 (M-F 8 a.m. - 5 p.m. CST)
 www.chloroprep.com

Artwork No. 6-800101
 H0611

Tear Here

WARNING
FLAMMABLE


Keep away from fire or flame.
 To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.

Wet hair is flammable

- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area


Lot # and Exp. Date are embossed on package.



Lot/Exp Placeholder - for reference only:

Lot # and Exp. Date are embossed on package.

**⚠ WARNING
FLAMMABLE**

 **Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:**

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

**ChloroPrep® Single Swabstick
ChloroPrep® Triple Swabstick**

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1.75ml APPLICATOR
5.25ml APPLICATOR

 Latex Free  Do Not Reuse
Single Use External Use Only Professional Use Only
Cat. No. 260100 / 260100B NDC 054365-400-07
Cat. No. 260103 / 260103B NDC 054365-400-08

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery: Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

14

ChloroPrep® Single Swabstick/ChloroPrep® Triple Swabstick
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area using
 - one single swabstick applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
 - three swabsticks sequentially is approximately 5 in. x 5 in. (160 cm²)
- tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork No. 6-656900 • L0411 • Made in the USA of US and imported parts

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v Antiseptic
Isopropyl alcohol 70% w/v Antiseptic

Purposes

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove all wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area using three swabsticks sequentially is approximately 5 in x 5 in (160 cm²)
- tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicators after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- USP purified water

Questions?

- www.chloraprep.com
- call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)

TRIP
SWABST

Clear

LOT

(b) (4)

ChloraPrep® Triple Swabstick
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
 5.25ml APPLICATOR

See Drug Facts for full labeling information
 Made in the USA of US and imported parts

Single Use
 Latex Free
 Cat. No. 260103 NDC 054365-400-08

Single use only
 Manufactured by:
 CareFusion
 Leawood, KS 66211
 1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST)
 www.chloraprep.com

Artwork No. 6-800104 60611

Tear Here

WARNING
FLAMMABLE

Keep away from fire or flame.
 To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

Directions:

- Tear pouch at notch.
- Pull across package to expose the ends of the swabsticks.
- Do not touch foam when removing swabsticks from pouch.
- Place foam flat side down on treatment area.

Lot # and Exp. Date are embossed on package film

↑

.95" From left edge of live area to point of arrow for tear perforation

Lot/Exp Date placeholder - for reference only:

Lot # and Exp. Date are embossed on package film

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER
08/16/2012