

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

SAVE \$XXX
on your next purchase

24HR
PREVACID
Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- Clinically Proven To Treat Frequent Heartburn

14 CAPSULES
ONE 14-DAY COURSE OF TREATMENT

Sodium Free

1 BOTTLE INSIDE
14 CAPSULES TOTAL



Actual Size
ONE 14-DAY COURSE OF TREATMENT

PREVACID 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer
CAPSULES
ONE 14-DAY COURSE OF TREATMENT

Drug Facts

Active ingredient (in each capsule)	Purpose
Lansoprazole 15 mg	Acid reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)

Drug Facts (continued)

- theophylline (asthma medicine)
- tacrolimus (immune system medicine)
- atazanavir (medicine for HIV infection)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

• children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

MADE IN: _____

EXP: _____

LOT: _____

0 00 0000 0000 0
FPO
0 000000 000000 0

Pharmaceuticals North America, Inc. and is used under license.
PREVACID® is a registered trademark of Takeda
Pharmapny, NJ 07054-0622 ©2011 XXXXXX
Distributed by: Novartis Consumer Health, Inc.
NOVARTIS

MANUFACTURER'S COUPON • EXPIRES XX/XX/XX



SAVE \$X.XX

Use as directed.

on any Prevacid®24HR XX ct. product

Use as directed for 14 days to treat frequent heartburn (occurring 2 or more days a week). May take 1-4 days for full effect.

SAVE \$XXX
on your next purchase

COUPON NOT VALID FOR PURCHASES REIMBURSED BY MEDICARE, MEDICAID, OR SIMILAR FEDERAL AND STATE PROGRAMS (INCLUDING STATE PHARMACEUTICAL ASSISTANCE PROGRAMS). COUPON NOT VALID FOR RESIDENTS OF MASSACHUSETTS IF PURCHASE IS REIMBURSED BY A HEALTHCARE INSURER. COUPON IS NOT VALID WHERE PROHIBITED OR RESTRICTED BY LAW. **RETAILER:** We will reimburse you for the lesser of the face value of this coupon or the retail price of this product, plus 8¢, if submitted in compliance with our redemption policy, available upon request. Cash value 1/20¢. Mail to: Novartis Consumer Health, CMS Dept, 00067, 1 Fawcett Dr., Del Rio, TX 78840. Limit one coupon per purchase. Coupon cannot be combined with any other offer.

©2011 Novartis Consumer Health, Inc. Prevacid® is a registered trademark of Takeda Pharmaceuticals North America, Inc. and is used under license.

000000-000000



FPO

GS1 Interim Barcode

DRAFT - FOR FDA REVIEW

DRUG FACTS TEXT	
	TYPE SIZE/STYLE
DRUG FACTS TITLE	9.0 pt. / Helvetica Condensed Bold Obl.
HEADINGS	8.0 pt. / Helvetica Condensed Bold Obl.
SUBHEADS	6.0 pt. / Helvetica Condensed Bold
BODY TEXT	6.0 pt. / Helvetica Condensed
BULLETS	5.0 pt. / Helvetica Condensed
TRACKING: 0	HORIZONTAL SCALE: 85% LEADING: 6.5 pt.
CHARACTERS/INCH: <39	
BARLINES, HAIRLINES: 1.5 pt., 0.5 pt.	

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

NDC 0067-6286-28

SAVE \$XXX
on your next purchase

PREVACID[®] 24HR

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- Clinically Proven To Treat Frequent Heartburn

28 CAPSULES
TWO 14-DAY COURSES OF TREATMENT

Sodium Free

2 BOTTLES INSIDE
28 CAPSULES TOTAL



Actual Size
ONE 14-DAY COURSE OF TREATMENT

Drug Facts

Active ingredient (in each capsule)	Purpose
Lansoprazole 15 mg	Acid reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)

Drug Facts (continued)

- theophylline (asthma medicine)
- tacrolimus (immune system medicine)
- atazanavir (medicine for HIV infection)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

0 00 0000 0000 0

FPO

MADE IN: _____

EXP: _____

LOT: _____

Distributed by: **Novartis Consumer Health, Inc.**
Pharmaceuticals North America, Inc. and is used under license.
PREVACID[®] is a registered trademark of Takeda
Farapany, NJ 07054-0622 ©2011 XXXXXX

MANUFACTURER'S COUPON • EXPIRES XX/XX/XX



SAVE \$X.XX

Use as directed.

on any Prevacid[®]24HR XX ct. product

Use as directed for 14 days to treat frequent heartburn (occurring 2 or more days a week). May take 1-4 days for full effect.

SAVE \$XXX
on your next purchase

COUPON NOT VALID FOR PURCHASES REIMBURSED BY MEDICARE, MEDICAID, OR SIMILAR FEDERAL AND STATE PROGRAMS (INCLUDING STATE PHARMACEUTICAL ASSISTANCE PROGRAMS). COUPON NOT VALID FOR RESIDENTS OF MASSACHUSETTS IF PURCHASE IS REIMBURSED BY A HEALTHCARE INSURER. COUPON IS NOT VALID WHERE PROHIBITED OR RESTRICTED BY LAW. **RETAILER:** We will reimburse you for the lesser of the face value of this coupon or the retail price of this product, plus 8¢, if submitted in compliance with our redemption policy, available upon request. Cash value 1/20¢. Mail to: Novartis Consumer Health, CMS Dept. 00067, 1 Fawcett Dr., Del Rio, TX 78840. Limit one coupon per purchase. Coupon cannot be combined with any other offer.

©2011 Novartis Consumer Health, Inc. Prevacid[®] is a registered trademark of Takeda Pharmaceuticals North America, Inc. and is used under license.

000000-000000

FPO

GS1 Interim Barcode

0 00000 00000 0

DRAFT - FOR FDA REVIEW

DRUG FACTS TEXT	
	TYPE SIZE/STYLE
DRUG FACTS TITLE	9.0 pt. / Helvetica Condensed Bold Obl.
HEADINGS	8.0 pt. / Helvetica Condensed Bold Obl.
SUBHEADS	6.0 pt. / Helvetica Condensed Bold
BODY TEXT	6.0 pt. / Helvetica Condensed
BULLETS	5.0 pt. / Helvetica Condensed
TRACKING: 0	HORIZONTAL SCALE: 85% LEADING: 6.5 pt.
CHARACTERS/INCH: <39	
BARLINES, HAIRLINES: 1.5 pt., 0.5 pt.	



PREVACID[®] 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer

3 BOTTLES INSIDE
CAPSULES
THREE 14-DAY COURSES OF TREATMENT

NDC 0067-6286-42

3 BOTTLES INSIDE
42 CAPSULES TOTAL



Actual Size
ONE 14-DAY COURSE OF TREATMENT

Drug Facts

Active ingredient (in each capsule) Lansoprazole 15 mg	Purpose Acid reducer
Use • treats frequent heartburn (occurs 2 or more days a week) • not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect	
Warnings Allergy alert: Do not use if you are allergic to lansoprazole Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have • liver disease • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain Ask a doctor or pharmacist before use if you are taking • warfarin (blood-thinning medicine) • prescription antifungal or anti-yeast medicines • digoxin (heart medicine)	
Drug Facts (continued) • theophylline (asthma medicine) • tacrolimus (immune system medicine) • atazanavir (medicine for HIV infection) Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days • you need to take more than 1 course of treatment every 4 months if pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions • adults 18 years of age and older • this product is to be used once a day (every 24 hours), every day for 14 days • it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours 14-Day Course of Treatment • swallow 1 capsule with a glass of water before eating in the morning • take every day for 14 days • do not take more than 1 capsule a day • swallow whole. Do not crush or chew capsules. • do not use for more than 14 days unless directed by your doctor Repeated 14-Day Courses (if needed) • you may repeat a 14-day course every 4 months • do not take for more than 14 days or more often than every 4 months unless directed by a doctor • children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.	

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert.
- They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

SAVE \$XXX
on your next purchase

PREVACID[®] 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- Clinically Proven To Treat Frequent Heartburn

42 CAPSULES
THREE 14-DAY COURSES OF TREATMENT

Sodium Free

0 00 0000 0000 0

FPO

0 000000 000000 0

MADE IN: _____
EXP: _____
LOT: _____

Distributed by: Novartis Consumer Health, Inc.
Parapany, NJ 07654-0622 ©2011 XXXXXX
PREVACID[®] is a registered trademark of Takeda Pharmaceuticals North America, Inc. and is used under license.

MANUFACTURER'S COUPON • EXPIRES XX/XX/XX



SAVE \$X.XX

SAVE \$XXX

on your next purchase

Use as directed.

on any Prevacid[®]24HR XX ct. product

Use as directed for 14 days to treat frequent heartburn (occurring 2 or more days a week). May take 1-4 days for full effect.

COUPON NOT VALID FOR PURCHASES REIMBURSED BY MEDICARE, MEDICAID, OR SIMILAR FEDERAL AND STATE PROGRAMS (INCLUDING STATE PHARMACEUTICAL ASSISTANCE PROGRAMS). COUPON NOT VALID FOR RESIDENTS OF MASSACHUSETTS IF PURCHASE IS REIMBURSED BY A HEALTHCARE INSURER. COUPON IS NOT VALID WHERE PROHIBITED OR RESTRICTED BY LAW. **RETAILER:** We will reimburse you for the lesser of the face value of this coupon or the retail price of this product, plus 8¢, if submitted in compliance with our redemption policy, available upon request. Cash value 1/20¢. Mail to: Novartis Consumer Health, CMS Dept. 00067, 1 Fawcett Dr., Del Rio, TX 78840. Limit one coupon per purchase. Coupon cannot be combined with any other offer.
©2011 Novartis Consumer Health, Inc. Prevacid[®] is a registered trademark of Takeda Pharmaceuticals North America, Inc. and is used under license.

000000-000000

FPO

0 000000 000000 0

GS1
Interim
Barcode

DRUG FACTS TEXT	
	TYPE SIZE/STYLE
DRUG FACTS TITLE	9.0 pt. / Helvetica Condensed Bold Obl.
HEADINGS	8.0 pt. / Helvetica Condensed Bold Obl.
SUBHEADS	6.0 pt. / Helvetica Condensed Bold
BODY TEXT	6.0 pt. / Helvetica Condensed
BULLETS	5.0 pt. / Helvetica Condensed
TRACKING: 0	HORIZONTAL SCALE: 85% LEADING: 6.5 pt.
CHARACTERS/INCH: <39	
BARLINES, HAIRLINES: 1.5 pt., 0.5 pt.	

DRAFT - FOR FDA REVIEW

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER
04/05/2012