

1125

1126

## MEDICATION GUIDE

1127

### CellCept® [SEL-sept]

1128

(mycophenolate mofetil capsules)

1129

(mycophenolate mofetil tablets)

1130

### CellCept® Oral Suspension

1131

(mycophenolate mofetil for oral suspension)

1132

### CellCept® Intravenous

1133

(mycophenolate mofetil hydrochloride for injection)

1134

Read the Medication Guide that comes with CellCept before you start taking it and each time you refill your prescription. There may be new information. This Medication Guide does not take the place of talking with your healthcare provider about your medical condition or your treatment.

1135

1136

1137

1138

### What is the most important information I should know about CellCept?

1139

#### CellCept can cause serious side effects:

1140

- **Possible loss of a pregnancy and higher risk of birth defects.** Women who take CellCept during pregnancy have a higher risk of losing a pregnancy (miscarriage) during the first 3 months (first trimester), and a higher risk that their baby will be born with birth defects

1141

1142

1143

1144

If you are a female and are able to become pregnant

1145

- your healthcare provider must talk with you about effective birth control methods (contraceptive counseling)

1146

1147

- you should have a negative pregnancy test within 1 week before you start to take CellCept

1148

1149

- you must use 2 different types of effective birth control at the same time, for 4 weeks before you start taking CellCept, during your entire CellCept therapy and for 6 weeks after stopping CellCept, unless you choose to avoid sexual intercourse completely (abstinence). CellCept decreases blood levels of the hormones in birth control pills that you take by mouth. Birth control pills may not work as well while you take CellCept, and you could become pregnant

1150

1151

1152

1153

1154

1155

1156

If you plan to become pregnant, talk with your healthcare provider. Your healthcare provider will decide if other medicines to prevent rejection may be right for you. In certain situations, you and your healthcare provider may decide that taking CellCept is more important to your health than the possible risks to your unborn baby.

1157

1158

1159

1160

1161

- **If you get pregnant while taking CellCept, do not stop taking CellCept. Call your healthcare provider right away.** You and your healthcare provider should report any cases of pregnancies to

1162

1163

1164

1165  
1166  
1167  
1168  
1169  
  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207

- FDA MedWatch at 1-800-FDA-1088
- Genentech at 1-888-835-2555

Talk to your healthcare provider about joining the National Transplantation Pregnancy Registry at 1-877-955-6877.

- **Increased risk of getting serious infections.** CellCept weakens the body’s immune system and affects your ability to fight infections. Serious infections can happen with CellCept and can lead to death. Types of infections can include:
  - **Viral infections.** Certain viruses can live in your body and cause active infections when your immune system is weak. Viral infections that can happen with CellCept include:
    - Shingles, other herpes infections, and cytomegalovirus (CMV). CMV can cause serious tissue and blood infections.
    - BK virus. BK virus can affect how your kidney works and cause your transplanted kidney to fail.
  - **A brain infection called Progressive Multifocal Leukoencephalopathy (PML).** In some patients, CellCept may cause an infection of the brain that may cause death. You are at risk for this brain infection because you have a weakened immune system. You should tell your healthcare provider right away if you have any of the following symptoms:
    - Weakness on one side of the body
    - You do not care about things that you usually care about (apathy)
    - You are confused or have problems thinking
    - You can not control your muscles
  - **Fungal infections.** Yeasts and other types of fungal infections can happen with CellCept and can cause serious tissue and blood infections (see “What are the possible side effects of CellCept?”)

**Call your healthcare provider right away if you have any of the following signs and symptoms of infection:**

- Temperature of 100.5°F or greater
- Cold symptoms, such as a runny nose or sore throat
- Flu symptoms, such as an upset stomach, stomach pain, vomiting or diarrhea
- Earache or headache
- Pain during urination
- White patches in the mouth or throat
- Unexpected bruising or bleeding
- Cuts, scrapes or incisions that are red, warm and oozing pus

- 1208
- 1209
- 1210
- 1211
- 1212
- 1213
- 1214
- 1215
- 1216
- 1217
- 1218
- 1219
- **Increased risk of getting certain cancers.** People who take CellCept have a higher risk of getting lymphoma, and other cancers, especially skin cancer. Tell your healthcare provider if you have:
    - unexplained fever, prolonged tiredness, weight loss or lymph node swelling
    - a brown or black skin lesion with uneven borders, or one part of the lesion does not look like the other
    - a change in the size and color of a mole
    - a new skin lesion or bump
    - any other changes to your health

**See the section “What are the possible side effects of CellCept?” for information about other serious side effects.**

### 1220 **What is CellCept?**

1221 CellCept is a prescription medicine to prevent rejection (antirejection medicine) in people  
1222 who have received a kidney, heart or liver transplant. Rejection is when the body’s  
1223 immune system perceives the new organ as a “foreign” threat and attacks it.

1224 CellCept is used with other medicines called cyclosporine (Sandimmune<sup>®</sup>, Gengraf<sup>®</sup>,  
1225 Neoral<sup>®</sup>) and corticosteroids. These medicines work together to prevent rejection to your  
1226 transplanted organ.

1227 CellCept has been used safely and works in children who received a kidney transplant as  
1228 it does in adults. It is not known if CellCept is safe and works in children who receive a  
1229 heart or liver transplant.

### 1230 **Who should not take CellCept?**

1231 **Do not take CellCept if you are allergic to mycophenolate mofetil or any of the**  
1232 **ingredients in CellCept.** See the end of this Medication Guide for a complete list of  
1233 ingredients in CellCept.

### 1234 **What should I tell my healthcare provider before taking CellCept?**

1235 Tell your healthcare provider about all of your medical conditions, if you:

- 1236
- 1237
- 1238
- 1239
- 1240
- 1241
- 1242
- 1243
- 1244
- 1245
- **have any digestive problems**, such as ulcers
  - **have Phenylketonuria (PKU).** CellCept oral suspension contains aspartame (a source of phenylalanine)
  - **have Lesch-Nyhan or Kelley-Seegmiller syndrome or another rare inherited deficiency hypoxanthine-guanine phosphoribosyl-transferase (HGPRT).** You should not take CellCept if you have one of these disorders
  - **plan to receive any vaccines. People taking CellCept should not take live vaccines.** Some vaccines may not work as well during treatment with CellCept
  - **are pregnant or are planning to become pregnant.** See “What is the most important information I should know about CellCept?”

- 1246 • **are breastfeeding.** It is not known if CellCept passes into breast milk. You and  
1247 your healthcare provider will decide if you will take CellCept or breastfeed. You  
1248 should not do both without first talking with your healthcare provider  
1249

1250 **Tell your healthcare provider about all of the medicines you are taking including**  
1251 **prescription and nonprescription medicines, vitamins and herbal supplements.**

1252 Some medicines may affect the way CellCept works, and CellCept may affect how some  
1253 medicines work. Especially tell your healthcare provider if you take:

- 1254 • birth control pills (oral contraceptives). See **“What is the most important**  
1255 **information I should know about CellCept?”**
- 1256 • sevelamer (Renagel<sup>®</sup>, Renvela<sup>™</sup>). These products should be taken 2 hours after  
1257 taking CellCept
- 1258 • acyclovir (Zovirax<sup>®</sup>), valacyclovir (Valtrex<sup>®</sup>), ganciclovir (CYTOVENE<sup>®</sup>-IV,  
1259 Vitrasert<sup>®</sup>), valganciclovir (VALCYTE<sup>®</sup>)
- 1260 • rifampin (Rifater<sup>®</sup>, Rifamate<sup>®</sup>, Rimactane<sup>®</sup>, Rifadin<sup>®</sup>)
- 1261 • antacids that contain magnesium and aluminum (CellCept and the antacid should  
1262 not be taken at the same time)
- 1263 • proton pump inhibitors (PPIs) (Prevacid<sup>®</sup>, Protonix<sup>®</sup>)
- 1264 • sulfamethoxazole/trimethoprim (BACTRIM<sup>™</sup>, BACTRIM DS<sup>™</sup>)
- 1265 • norfloxacin (Noroxin<sup>®</sup>) and metronidazole (Flagyl<sup>®</sup>, Flagyl<sup>®</sup> ER, Flagyl<sup>®</sup> IV,  
1266 Metro IV, Helidac<sup>®</sup>, Pylera<sup>™</sup>)
- 1267 • ciprofloxacin (Cipro<sup>®</sup>, Cipro<sup>®</sup> XR, Ciloxan<sup>®</sup>, Proquin<sup>®</sup> XR) and amoxicillin plus  
1268 clavulanic acid (Augmentin<sup>®</sup>, Augmentin XR<sup>™</sup>)
- 1269 • azathioprine (Azasan<sup>®</sup>, Imuran<sup>®</sup>)
- 1270 • cholestyramine (Questran Light<sup>®</sup>, Questran<sup>®</sup>, Locholest Light, Locholest,  
1271 Prevalite<sup>®</sup>)

1272 Know the medicines you take. Keep a list of them to show to your healthcare provider  
1273 and pharmacist when you get a new medicine. Do not take any new medicine without  
1274 talking with your healthcare provider.

1275 **How should I take CellCept?**

- 1276 • Take CellCept exactly as prescribed  
1277
- 1278 • Do not stop taking CellCept or change the dose unless your healthcare provider  
1279 tells you to  
1280
- 1281 • If you miss a dose of CellCept, or are not sure when you took your last dose, take  
1282 the regular amount of CellCept prescribed as soon as you remember. If it is time  
1283 for your next dose, skip the missed dose. Do not take 2 doses at the same time.  
1284 Call your healthcare provider if you are not sure what to do  
1285
- 1286 • Take CellCept capsules, tablets and oral suspension on an empty stomach, either 1  
1287 hour before or 2 hours after a meal, unless your healthcare provider tells you

1288 otherwise. With the approval of your healthcare provider, in stable kidney  
1289 transplant patients, CellCept can be taken with food if necessary  
1290  
1291 • Most people take CellCept by mouth either as blue and brown capsules or  
1292 lavender tablets. Some people may get CellCept soon after their transplant surgery  
1293 as an infusion into a vein  
1294  
1295 • Do not crush CellCept tablets. Do not open or crush CellCept capsules  
1296  
1297 • If you are not able to swallow CellCept tablets or capsules, your healthcare  
1298 provider may prescribe CellCept Oral Suspension. This is a liquid form of  
1299 CellCept. Your pharmacist will mix the medicine before giving it to you  
1300  
1301 • Do not mix CellCept Oral Suspension with any other medicine  
1302  
1303 • If you take too much CellCept, call your healthcare provider or the poison control  
1304 center right away  
1305

1306 **What should I avoid while taking CellCept?**

- 1307 • **Avoid pregnancy.** See “**What is the most important information I should know about CellCept?**”
- **Limit the amount of time you spend in sunlight.** Avoid using tanning beds or sunlamps. People who take CellCept have a higher risk of getting skin cancer. (See “**What is the most important information I should know about CellCept?**”) Wear protective clothing when you are in the sun and use a sunscreen with a high protection factor (SPF 30 and above). This is especially important if your skin is very fair or if you have a family history of skin cancer

1308

1309 **What are the possible side effects of CellCept?**

**CellCept can cause serious side effects:**

- 1310 • See “**What is the most important information I should know about CellCept?**”
- **Low blood cell counts.** People taking high doses of CellCept each day may have a decrease in blood counts, including
  - 1311 • **white blood cells, especially neutrophils.** Neutrophils fight against bacterial
1312 infections. You have a higher chance of getting an infection when your white
1313 blood cell count is low. This is most common from 3 months to 6 months after1314 your transplant- 1315 • **red blood cells.** Red blood cells carry oxygen to your body tissues. You have a
1316 higher chance of getting severe anemia when your red blood cell count is low- 1317 • **platelets.** Platelets help with blood clotting
1318

1319 Your healthcare provider will do blood tests before you start taking CellCept and  
1320 during treatment with CellCept to check your blood cell counts.

1321 Tell your healthcare provider right away if you have any signs of infection (see  
1322 **“What is the most important information I should know about CellCept?”**), or  
1323 any unexpected bruising or bleeding. Also, tell your healthcare provider if you have  
1324 unusual tiredness, lack of energy, dizziness or fainting.

- **Stomach problems.** Stomach and intestinal bleeding can happen in people who take high doses of CellCept. Bleeding can be severe and you may have to be hospitalized for treatment

1325

1326 **Common side effects include:**

- diarrhea. Call your healthcare provider right away if you have diarrhea. Do not stop taking CellCept without first talking with your healthcare provider
- vomiting
- pain
- stomach area pain
- swelling of the lower legs, ankles and feet
- high blood pressure

1333

1334

1335

Side effects that happen more often in children than in adults taking CellCept include:

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

- stomach area pain
- fever
- infection
- pain
- blood infection (sepsis)
- diarrhea
- vomiting
- sore throat
- colds (respiratory tract infections)
- high blood pressure
- low white blood cell count
- low red blood cell count

These are not all of the possible side effects of CellCept. Tell your healthcare provider about any side effect that bothers you or that does not go away.

1346

1347

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088 or to Genentech at 1-888-835-2555.

1348

**How should I store CellCept?**

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

- Store CellCept capsules and tablets at room temperature, between 59°F to 86°F (15°C to 30°C). Keep the container closed tightly
- Store the prepared CellCept Oral Suspension at room temperature, between 59°F to 86°F (15°C to 30°C), for up to 60 days. You can also store CellCept Oral Suspension in the refrigerator at 36°F to 46°F (2°C to 8°C). **Do not freeze CellCept Oral Suspension**
- **Keep CellCept and all medicines out of the reach of children**

1359 **General Information about CellCept**

1360 Medicines are sometimes prescribed for purposes other than those listed in a Medication  
1361 Guide. Do not use CellCept for a condition for which it was not prescribed. Do not give  
1362 CellCept to other people, even if they have the same symptoms that you have. It may  
1363 harm them.

1364 This Medication Guide summarizes the most important information about CellCept. If  
1365 you would like more information, talk with your doctor. You can ask your doctor or  
1366 pharmacist for information about CellCept that is written for healthcare professionals. For  
1367 more information, call 1-888-835-2555 or visit  
1368 [www.gene.com/gene/products/information/cellept](http://www.gene.com/gene/products/information/cellept).

1369 **What are the ingredients in CellCept?**

1370 **Active Ingredient:** mycophenolate mofetil

1371 **Inactive Ingredients:**

1372 CellCept 250 mg capsules: croscarmellose sodium, magnesium stearate, povidone (K-90)  
1373 and pregelatinized starch. The capsule shells contain black iron oxide, FD&C blue #2,  
1374 gelatin, red iron oxide, silicon dioxide, sodium lauryl sulfate, titanium dioxide, and  
1375 yellow iron oxide.

1376 CellCept 500 mg tablets: black iron oxide, croscarmellose sodium, FD&C blue #2  
1377 aluminum lake, hydroxypropyl cellulose, hydroxypropyl methylcellulose, magnesium  
1378 stearate, microcrystalline cellulose, polyethylene glycol 400, povidone (K-90), red iron  
1379 oxide, talc, and titanium dioxide; may also contain ammonium hydroxide, ethyl alcohol,  
1380 methyl alcohol, n-butyl alcohol, propylene glycol, and shellac.

1381 CellCept Oral Suspension: aspartame, citric acid anhydrous, colloidal silicon dioxide,  
1382 methylparaben, mixed fruit flavor, sodium citrate dihydrate, sorbitol, soybean lecithin,  
1383 and xanthan gum.

1384 CellCept Intravenous: polysorbate 80, and citric acid. Sodium hydroxide may have been  
1385 used in the manufacture of CellCept Intravenous to adjust the pH.

1386

1387 This Medication Guide has been approved by the US Food and Drug Administration.

1388 CELLCEPT, CYTOVENE-IV, and VALCYTE are registered trademarks of Hoffmann-La Roche Inc.  
1389 BACTRIM and BACTRIM DS are trademarks of Hoffmann-La Roche Inc.

1390

1391

Distributed by:

**Genentech USA, Inc.**

A Member of the Roche Group

1 DNA Way

1392 South San Francisco, CA 94080-4990

MG Revised: June 2012

1393 For additional copies of this Medication Guide, please call 1-800-617-8191 or visit  
1394 [www.gene.com/gene/products/informaiton/cellcept](http://www.gene.com/gene/products/informaiton/cellcept).

1395

1396 CTCTIO\_1061443\_PI/MG\_2011\_02\_K

1397

1398 © 20xx Genentech, Inc. All rights reserved.