

**MEDICATION GUIDE**  
**DYSPO<sup>®</sup> (DIS-port)**  
**(abobotulinumtoxin A)**

**Injection**

Read the Medication Guide that comes with DYSPO<sup>®</sup> before you start using it and each time DYSPO<sup>®</sup> is given to you. There may be new information. This information does not take the place of talking with your doctor about your medical condition or your treatment. You should share this information with your family members and caregivers.

**What is the most important information I should know about DYSPO<sup>®</sup>?**

**DYSPO<sup>®</sup> may cause serious side effects that can be life threatening including:**

- **Problems breathing or swallowing**
- **Spread of toxin effects**

**These problems can happen within hours, or days to weeks after an injection of DYSPO<sup>®</sup>. Call your doctor or get medical help right away if you have any of these problems after treatment with DYSPO<sup>®</sup>:**

- 1. Problems swallowing, speaking, or breathing.** These problems can happen within hours, or days to weeks after an injection of DYSPO<sup>®</sup>, usually because the muscles that you use to breathe and swallow can become weak after the injection. Death can happen as a complication if you have severe problems with swallowing or breathing after treatment with DYSPO<sup>®</sup>.
  - People with certain breathing problems may need to use muscles in their neck to help them breathe. These patients may be at greater risk for serious breathing problems with DYSPO<sup>®</sup>.
  - Swallowing problems may last for several weeks. People who can not swallow well may need a feeding tube to receive food and water. If swallowing problems are severe, food or liquids may go into your lungs. People who already have swallowing or breathing problems before receiving DYSPO<sup>®</sup> have the highest risk of getting these problems.
  
- 2. Spread of toxin effects.** In some cases, the effect of botulinum toxin may affect areas of the body away from the injection site and cause symptoms of a serious condition called botulism. The symptoms of botulism include:
  - loss of strength and muscle weakness all over the body

- double vision
- blurred vision and drooping eyelids
- hoarseness or change or loss of voice (dysphonia)
- trouble saying words clearly (dysarthria)
- loss of bladder control
- trouble breathing
- trouble swallowing

**These symptoms can happen within hours, or days to weeks after you receive an injection of DYSPORT®.**

These problems could make it unsafe for you to drive a car or do other dangerous activities. See “What should I avoid while receiving DYSPORT®?”.

### **What is DYSPORT®?**

DYSPORT® is a prescription medicine that is injected into muscles and used:

- to treat the abnormal head position and neck pain that happens with cervical dystonia (CD) in adults
- to improve the look of moderate to severe frown lines between the eyebrows (glabellar lines) in adults younger than 65 years of age for a short period of time (temporary)

CD is caused by muscle spasms in the neck. These spasms cause abnormal position of the head and often neck pain. After DYSPORT® is injected into muscles, those muscles are weakened for up to 12 to 16 weeks or longer. This may help lessen your symptoms.

Frown lines (wrinkles) happen because the muscles that control facial expression are used often (muscle tightening over and over). After DYSPORT® is injected into the muscles that control facial expression, the medicine stops the tightening of these muscles for up to 4 months.

It is not known whether DYSPORT® is safe or effective in children under 18 years of age.

It is not known whether DYSPORT® is safe or effective for the treatment of other types of muscle spasms. It is not known whether DYSPORT® is safe or effective for the treatment of other wrinkles.

### **Who should not take DYSPORT®?**

#### **Do not take DYSPORT® if you:**

- are allergic to DYSPORT® or any of the ingredients in DYSPORT®. See the end of this Medication Guide for a list of ingredients in DYSPORT®
- are allergic to cow's milk protein

- had an allergic reaction to any other botulinum toxin product such as Myobloc<sup>®</sup> (rimabotulinumtoxinB)<sup>\*</sup>, Botox<sup>®</sup> (onabotulinumtoxinA)<sup>\*</sup>, or Xeomin<sup>®</sup> (incobotulinumtoxinA)<sup>\*</sup>.
- have a skin infection at the planned injection site

### **What should I tell my doctor before taking DYSPORT<sup>®</sup>?**

#### **Tell your doctor about all your medical conditions, including if you:**

- have a disease that affects your muscles and nerves (such as amyotrophic lateral sclerosis [ALS or Lou Gehrig's disease], myasthenia gravis or Lambert-Eaton syndrome). See "What is the most important information I should know about DYSPORT<sup>®</sup>?"
- have allergies to any botulinum toxin product
- had any side effect from any botulinum toxin product in the past
- have or have had a breathing problem, such as asthma or emphysema
- have or have had swallowing problems
- have or have had bleeding problems
- have diabetes
- have or have had a slow heart beat or other problem with your heart rate or rhythm
- have plans to have surgery
- had surgery on your face
- have weakness of your forehead muscles (such as trouble raising your eyebrows)
- have drooping eyelids
- have any other change in the way your face normally looks
- are pregnant or plan to become pregnant. It is not known if DYSPORT<sup>®</sup> can harm your unborn baby
- are breast-feeding or planning to breast-feed. It is not known if DYSPORT<sup>®</sup> passes into breast milk

**Tell your doctor about all the medicines you take**, including prescription and nonprescription medicines, vitamins and herbal products. Using DYSPORT<sup>®</sup> with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received DYSPORT<sup>®</sup> in the past.**

---

<sup>\*</sup> All trademarks are the property of their respective owners

### **Especially tell your doctor if you:**

- have received any other botulinum toxin product in the last four months
- have received injections of botulinum toxin, such as Myobloc<sup>®</sup> (rimabotulinumtoxinB)\*, Botox<sup>®</sup> (onabotulinumtoxinA)\* or Xeomin<sup>®</sup> (incobotulinumtoxinA)\* in the past; be sure your doctor knows exactly which product you received
- have recently received an antibiotic by injection
- take muscle relaxants
- take an allergy or cold medicine
- take a sleep medicine

### **Ask your doctor if you are not sure if your medicine is one that is listed above.**

Know the medicines you take. Keep a list of your medicines with you to show your doctor and pharmacist each time you get a new medicine.

### **How should I take DYSPORT<sup>®</sup>?**

- DYSPORT<sup>®</sup> is an injection that your doctor will give you
- DYSPORT<sup>®</sup> is injected into the affected muscles
- Your doctor may give you another dose of DYSPORT<sup>®</sup> after 12 weeks or longer, if it is needed
- If you are being treated for CD, your doctor may change your dose of DYSPORT<sup>®</sup>, until you and your doctor find the best dose for you
- The dose of DYSPORT<sup>®</sup> is not the same as the dose of any other botulinum toxin product

### **What should I avoid while taking DYSPORT<sup>®</sup>?**

DYSPORT<sup>®</sup> may cause loss of strength or general muscle weakness, blurred vision, or drooping eyelids within hours to weeks of taking DYSPORT<sup>®</sup>. **If this happens, do not drive a car, operate machinery, or do other dangerous activities. See “What is the most important information I should know about DYSPORT<sup>®</sup>?”**

### **What are the possible side effects of DYSPORT<sup>®</sup>?**

---

\* All trademarks are the property of their respective owners

**DYSPO<sup>®</sup> can cause serious side effects. See “What is the most important information I should know about DYSPO<sup>®</sup>?”**

**Other side effects of DYSPO<sup>®</sup> include:**

- dry mouth
- injection site discomfort or pain
- tiredness
- headache
- neck pain
- muscle pain
- eye problems: double vision, blurred vision, decreased eyesight, problems with focusing the eyes (accommodation), drooping eyelids, swelling of the eyelids
- allergic reactions. Symptoms of an allergic reaction to DYSPO<sup>®</sup> may include: itching, rash, red itchy welts, wheezing, asthma symptoms, or dizziness or feeling faint. Tell your doctor or get medical help right away if you get wheezing or asthma symptoms, or if you get dizzy or faint

**Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DYSPO<sup>®</sup>. For more information, ask your doctor or pharmacist.**

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1–800–FDA–1088. General information about DYSPO<sup>®</sup>:

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide.

This Medication Guide summarizes the most important information about DYSPO<sup>®</sup>. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about DYSPO<sup>®</sup> that is written for healthcare professionals. For more information about DYSPO<sup>®</sup> call 877-397-7671 or go to [www.dysport.com](http://www.dysport.com) or [www.DysportUSA.com](http://www.DysportUSA.com).

**What are the ingredients in DYSPO<sup>®</sup>?**

Active ingredient: (botulinum toxin Type A)

Inactive ingredients: human albumin, and lactose. DYSPO<sup>®</sup> may contain cow's milk protein.

Revised: MAY 2012

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Distributed by:

Tercica, Inc.

a subsidiary of the Ipsen Group

Brisbane, CA 94005

and

Medicis Aesthetics Inc.

a wholly owned subsidiary of Medicis Pharmaceutical Corporation

Scottsdale, AZ 85256