

ACTUAL SIZE

3/16" X 3/16" DATA MATRIX CODE (ECC200),
 PRINTED IN BLACK OR DARK BLUE ON
 UNPRINTED WHITE BACKGROUND WITH A
 MINIMUM OF 1/8" QUIET ZONE ALL 4 SIDES.
 DATA MATRIX CODE = **025196**

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult a doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by a doctor.
Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever; right before or after heart surgery
Ask a doctor before use if stomach bleeding warning applies to the child; child has a history of stomach problems, such as heartburn; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic.
Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition; taking any other drug
When using this product take with food or milk if upset stomach occurs; the risk of heart attack or stroke may increase if you use more than directed or for longer than directed.
Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomits blood, has bloody or black stools, or has

Do Not Use if breakable ring on bottle cap is separated or missing.

Children's Advil[®] Suspension
Fever Reducer / Pain Reliever (NSAID)

Active ingredient (in each 5 mL) ... Ibuprofen 100 mg
Use: temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Sugar-Free Dye-Free Berry-Flavored Liquid
4 FL OZ (120 mL)

stomach pain that does not get better; the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
 Each teaspoon contains sodium 10 mg. Store at 20-25°C (68-77°F).
Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. Find right dose on chart below. If possible, use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Weight (lb)	Age (yr)	Dose (tsp)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	1 tsp
36-47 lb	4-5 yr	1 1/2 tsp
48-59 lb	6-8 yr	2 tsp
60-71 lb	9-10 yr	2 1/2 tsp
72-95 lb	11 yr	3 tsp

Pfizer, Madison, NJ 07940 USA © 2013 Pfizer Inc.
 For most recent product information, visit www.AdvIL.com.
Questions or comments? call 1-800-88-ADVIL
 PAA025196.FDA02

UNVARNISHED
 UNPRINTED AREA
 FOR LOT & EXP.

090177e1846f0dfb\Final\Final On: 05-Aug-2013 14:24

PRODUCTION (NOT COLOR ACCURATE)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	5 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt

Children's Advil[®] Suspension

Packaged with Tamper-Evident bottle cap. Do Not Use if breakable ring is separated or missing.

SUGAR-FREE

Children's Advil[®] Suspension

Ibuprofen 100 mg Fever Reducer/Pain Reliever (NSAID) per 5 mL

Fever

Aches & Pains Lasts Up To 8 Hours

BERRY
Berry-Flavored Liquid for ages 2-11 years

4 FL OZ (120 mL) Alcohol-Free

Drug Facts (continued)
Ask a doctor or pharmacist before use if the child is
 ■ under a doctor's care for any serious condition
 ■ taking any other drug
When using this product
 ■ take with food or milk if stomach upset occurs
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
Stop use and ask a doctor if
 ■ child experiences any of the following signs of stomach bleeding:
 ■ feels faint
 ■ vomits blood
 ■ has bloody or black stools
 ■ has stomach pain that does not get better
 ■ the child does not get any relief within first day (24 hours) of treatment
 ■ fever or pain gets worse or lasts more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ this product does not contain directions or complete warnings for adult use
 ■ do not give more than directed
 ■ shake well before using
 ■ find right dose on chart below. If possible, use weight to dose; otherwise use age.
 ■ repeat dose every 6-8 hours, if needed
 ■ do not use more than 4 times a day
 ■ measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (tsp)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	1 tsp
36-47 lb	4-5 yr	1 1/2 tsp
48-59 lb	6-8 yr	2 tsp
60-71 lb	9-10 yr	2 1/2 tsp
72-95 lb	11 yr	3 tsp

Other information
 ■ each teaspoon contains: sodium 10 mg
 ■ one dose lasts 6-8 hours
 ■ store at 20-25°C (68-77°F)
 ■ see bottom of box for lot number and expiration date

Inactive ingredients
 artificial flavor, carbonylmethylcellulose sodium, citric acid monohydrate, edetate disodium, FD&C blue no. 1, FD&C red no. 40, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?
 call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional

Long Lasting: Up to 8 hours of fever reduction.
Great Convenience: Long duration means fewer doses needed per day.

For most recent product information, visit www.Advii.com
 Pfizer, Madison, NJ 07940 USA © 2013 Pfizer Inc.

PAA025199.FDA02

3 0573 0231 30 5

LOT:
EXP:


GLUE AREA UNVARNISHED

PRE-PRINTED TEXT. NOTE: LOT/EXP (OR EQUIVALENT) TO BE PRINTED PER MARKET REQUIREMENTS AT A MINIMUM OF 8 pt. DATE OF MANUFACTURE CAN BE ADDED AS A THIRD LINE IF NEEDED.

UNPRINTED AREA FOR DATE & LOT NUMBER UNVARNISHED

3/16" X 3/16" ARRAY ALPHA NUMERIC DATA MATRIX CODE (ECC 200) PRINTED IN BLACK OR DARK BLUE ON WHITE FIELD WITH WHITE 1/8" QUIET ZONE ON ALL 4 SIDES.
DATA MATRIX CODE = 025199

PRODUCTION (NOT COLOR ACCURATE)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Children's Advil[®] Suspension

Packaged with Tamper-Evident bottle cap. Do Not Use if breakable ring is separated or missing.

SUGAR-FREE

Children's Advil[®] Suspension

Ibuprofen 100 mg Fever Reducer/Pain Reliever (NSAID) per 5 mL

Fever

Aches & Pains Lasts Up To 8 Hours

Dye-Free BERRY
Berry-Flavored Liquid for ages 2-11 years

4 FL OZ (120 mL) Alcohol-Free

Drug Facts (continued)

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- repeat dose every 6-8 hours, if needed
- do not use more than 4 times a day
- measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (tsp)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	1 tsp
36-47 lb	4-5 yr	1 1/2 tsp
48-59 lb	6-8 yr	2 tsp
60-71 lb	9-10 yr	2 1/2 tsp
72-85 lb	11 yr	3 tsp

Other information

- each teaspoon contains: sodium 10 mg
- one dose lasts 6-8 hours
- store at 20-25°C (68-77°F)
- see bottom of box for lot number and expiration date

Inactive ingredients
artificial flavor, carboxymethylcellulose sodium, citric acid monohydrate, edetate disodium, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?
call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional

Long Lasting: Up to 8 hours of fever reduction
Great Convenience: Long duration means fewer doses needed per day.

For most recent product information, visit www.Advil.com.
Pfizer, Madison, NJ 07940 USA © 2013 Pfizer Inc.

Drug Facts

Active ingredient (in each 5 mL)

Ibuprofen 100 mg (NSAID)* ...Fever reducer/Pain reliever
*nonsteroidal anti-inflammatory drug

Purpose

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to the child
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has a history of stomach problems, such as heartburn
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child is taking a diuretic

3 0573 0232 30 2

PAA025197.FDA02

LOT:
EXP:

3/16" X 3/16" ARRAY ALPHA NUMERIC DATA MATRIX CODE (ECC 200) PRINTED IN BLACK OR DARK BLUE ON WHITE FIELD WITH WHITE 1/8" QUIET ZONE ON ALL 4 SIDES.
DATA MATRIX CODE = 025197

PRE-PRINTED TEXT. NOTE: LOT/EXP (OR EQUIVALENT) TO BE PRINTED PER MARKET REQUIREMENTS AT A MINIMUM OF 8 pt. DATE OF MANUFACTURE CAN BE ADDED AS A THIRD LINE IF NEEDED.

UNPRINTED AREA FOR DATE & LOT NUMBER UNVARNISHED

PRODUCTION (NOT COLOR ACCURATE) (b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
12/20/2013