


Cat. No. 260100 / 260100B NDC 054365-400-07
Cat. No. 260103 / 260103B NDC 054365-400-08

ChloroPrep® Single Swabstick/ChloroPrep® Triple Swabstick
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1.75ml APPLICATOR / 5.25ml APPLICATOR



Professional Use Only
External Use Only

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v
Isopropyl alcohol 70% v/v

Purposes

Antiseptic
Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame.

- do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Swabstick Insert – Back

ChloroPrep® Single Swabstick/ChloroPrep® Triple Swabstick
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area using
 - one single swabstick applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
 - three swabsticks sequentially is approximately 5 in. x 5 in. (160 cm²)
- tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds (when using the triple swabstick applicators, use each swabstick sequentially within the 30 seconds). Allow the area to air dry for approximately 1 minute. Do not blot or wipe away.
- moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes (when using the triple swabstick applicators, use each swabstick sequentially within the 2 minutes). Allow the area to air dry for approximately 1 minute. Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients USP purified water

Questions? ■ www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, El Paso, TX 79912 • Artwork No. 6-656900 • MXXXX • Made in the USA of US and imported parts

Back Single

Swabstick Foil



ChloroPrep® Single Swabstick
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1.75ml APPLICATOR

Do Not Reuse

See Drug Facts for full labeling information
Made in the USA of US and imported parts

Cat. No. 260100 NDC 054365-400-07

Manufactured by:
Carefusion
El Paso, TX 79972
1-800-523-0602
(M-F 8 a.m. - 5 p.m. CST)
www.chloroprep.com

Artwork No. 6-800101
10403

Tear Here


 **WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.**

Directions:

- Tear pouch at notch.
- Pull across package to expose the ends of the swabsticks.
- Do not touch foam when removing swabsticks from pouch.
- Place foam flat side down on treatment area.

Lot # and Exp. Date are displayed on package.

Single Swabstick Carton - Top Panel



ChloroPrep[®] One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v Antiseptic
Isopropyl alcohol 70% v/v Antiseptic

Purposes

..... Antiseptic
..... Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame.

- do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- tear pouch at side notch to reveal applicator handle. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds. Allow the area to air dry for approximately 1 minute. Do not blot or wipe away.
- moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes. Allow the area to air dry for approximately 1 minute. Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

USP purified water

Questions?

- www.chloroprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

Single Swabstick Carton – Front Panel

NDC 054365-400-07

1.75ml
**SINGLE
SWABSTICK**
APPLICATORS

Clear

 Do Not Reuse

 Not made with natural
rubber latex

48 applicators
0.06 fl. oz. (1.75 ml) each

ChloroPrep[®]
One-Step


2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation



Professional Use Only

Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat
above 40 °C (104 °F)

Single Swabstick Carton – Side Panel A



Single Swabstick Carton – Side Panel B

ChloroPrep[®]
One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

1.75 ml
SINGLE
SWABSTICK
APPLICATORS

Clear

CareFusion
El Paso, TX 79912
1.800.523.0502
www.chloroprep.com
Made in the USA of US and imported parts

Triple Swabstick Foil



ChloroPrep® Triple Swabstick

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

5.25ml APPLICATOR

Cat. No. 260103 NDC 054945-400-06



Do Not Reuse



Do Not Reuse

See Drug Facts for full labeling information
Made in the USA of US and imported parts

Manufactured by:
CareFusion
El Paso, TX 79912
1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST)
www.chloroprep.com
Artwork No. 6-800104 608302



Tear Here


 **WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.**

Directions:

- Tear pouch at notch.
- Pull across package to expose the ends of the swabsticks.
- Do not touch foam when removing swabsticks from pouch.
- Place foam flat side down on treatment area.

Lot # and Exp. Date are embossed on package film

Triple Swabstick Carton – Top Panel



ChloroPrep[®]

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate, 2% w/v Antiseptic
Isopropyl alcohol 70% v/v Antiseptic

Purposes

For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame.

- do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product

keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area using three swabsticks sequentially is approximately 5 in x 5 in (160 cm²)
- tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area. completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): using each swabstick sequentially, use gentle repeated back-and-forth strokes for a total of 30 seconds. Allow the area to air dry for approximately 30 seconds. Do not blot or wipe away.
- **moist surgical sites** (e.g., inguinal fold): using each swabstick sequentially, use gentle repeated back-and-forth strokes for a total of 2 minutes. Allow the area to air dry for approximately 1 minute. Do not blot or wipe away.
- discard the applicators after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

■ store between 15-30 °C (59-86 °F)

■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

USP purified water

Questions?

■ www.chloroprep.com

■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

Triple Swabstick Carton – Front Panel

NDC 054365-400-08

5.25ml
**TRIPLE
SWABSTICK**
APPLICATORS

Clear

 Do Not Reuse

 ~~LATEX~~ Not made with natural rubber latex

40 packages
0.18 fl. oz. (5.25 ml) each

ChloroPrep[®]
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation



Professional Use Only

Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat
above 40 °C (104 °F)

Triple Swabstick Carton – Side Panel A

ChloroPrep[®]
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

5.25ml
**TRIPLE
SWABSTICK**
APPLICATORS

Clear

LOT XXXXX EXP DATE: XX/XX
(printed in-line)

NO VARNISH AREA



(01) 20354365400086

Cat. No. 260103
©2009 CareFusion, All rights reserved. ChloroPrep is a registered trademark of CareFusion or one of its subsidiaries.

Triple Swabstick Carton – Side Panel B

ChloroPrep[®]

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

5.25ml
TRIPLE
SWABSTICK
APPLICATORS

Clear

CareFusion
El Paso, TX 79912
1.800.523.0502
www.chloroprep.com
Made in the USA of US and imported parts

Sepp Insert

ChloroPrep® One-Step
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
 0.67ml SEPP® APPLICATOR
WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts	
Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v.....	Antiseptic
Isopropyl alcohol 70% v/v.....	Antiseptic
Use for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infection.	
Warnings	
For external use only	
Flammable, keep away from fire or flame	
■ do not use with electrocautery procedures	
Do not use	
■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol	
■ for lumbar puncture or in contact with the meninges	
■ on open skin wounds or as a general skin cleanser	
When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions	
■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.	
■ maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm ²)	
■ remove applicator from package; do not touch applicator tip	
■ hold the sponge tip down and pinch the barrel of the applicator <u>only once</u> to activate the ampule and release the antiseptic	
■ wet the sponge tip by pressing and releasing the sponge against the treatment area until liquid is visible on the skin	
■ completely wet the treatment area with antiseptic	
■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds.	
■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute.	
■ do not blot or wipe away	
■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.	
Other information ■ store between 15–30 °C (50–86 °F)	
■ avoid freezing and excessive heat above 40 °C (104 °F)	
Inactive ingredients USP purified water	
Questions? ■ www.chloroprep.com	
■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)	

See other side for additional information

Front

ChloroPrep® One-Step
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
 0.67ml SEPP® APPLICATOR

Cat. No. 260441NSB / 260449NSB
 NDC 054365-400-03

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

See Drug Facts for Full Labeling Information

 Do Not Reuse  Not made with natural rubber latex

External Use Only Professional Use Only

CareFusion El Paso, TX 79912

Artwork 6-775449F • MXXXX
 Made in the USA of US and imported parts

Back

Sepp Cyrel Lidding


top area 16x9mm

ChloroPrep[®]
SEPP[®]
2% w/v chlorhexidine
gluconate (CHG)
and 70% v/v isopropyl
alcohol (IPA)
0.67 ml
APPLICATOR


**WARNING.
FLAMMABLE.
KEEP AWAY FROM
FIRE OR FLAME.**

 
Do Not Reuse
Applicator is sterile if
package is intact
CareFusion
El Paso, TX 79912
1-800-523-0502
(MF 8 a.m.- 5 p.m. CST)
Lot No: XXXXX
Exp.: XX/XXXX
OXXXX
Cat No. 260449
NDC 054365-400-03
Made in the USA of
US and imported parts

Sepp Carton – Top Panel



ChloroPrep[®] One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v Antiseptic
Isopropyl alcohol 70% v/v Antiseptic

Use for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame.
 ■ do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbur puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch applicator tip
- hold the sponge tip down and pinch the barrel of the applicator only once to activate the ampule and release the antiseptic
- wet the sponge tip by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm); use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds.
- moist surgical sites (e.g., inguinal fold); use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately one 1 minute.
- do not blot or wipe away
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

USP purified water

Questions?

- www.chloroprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

Sepp Carton – Front Panel

NDC 054365-400-03

**0.67ml
SEPP[®]
APPLICATORS**

Clear

 Do Not Reuse

 Not made with natural rubber latex

STERILE EO

Applicator is sterile if package is intact

200 applicators
0.02 fl. oz. (0.67 ml) each

**ChloroPrep[®]
One-Step**

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative
Skin Preparation



Professional Use Only

Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat
above 40 °C (104 °F)

Sepp Carton – Side Panel A



Sepp Carton - Side Panel B

ChloroPrep[®]
One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

0.67ml
SEPP[®]
APPLICATORS

Clear

LOT XXXXXX EXP DATE: XX/XX
(printed in-line)

No Varnish



(01) 20354365400031

Cat. No. 260449 ©2009 CareFusion. All rights reserved. ChloroPrep is a registered trademark of CareFusion or one of its subsidiaries.

Frepp Insert

ChloroPrep® One-Step
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
 1.5ml FREPP® APPLICATOR

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts	
Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v.....Antiseptic	
Isopropyl alcohol 70% v/v.....Antiseptic	
Use For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.	
Warnings	
For external use only	
Flammable, keep away from fire or flame	
<ul style="list-style-type: none"> do not use with electrocautery procedures 	
Do not use	
<ul style="list-style-type: none"> on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol for lumbar puncture or in contact with the meninges on open skin wounds or as a general cleanser 	
When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions	
<ul style="list-style-type: none"> use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²) remove applicator from package; do not touch sponge hold the applicator horizontally with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic. wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin completely wet the treatment area with antiseptic dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds. moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute. do not blot or wipe away discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. 	
Other information <ul style="list-style-type: none"> store between 15–30 °C (59–86 °F) avoid freezing and excessive heat above 40 °C (104 °F) 	
Inactive ingredients USP purified water	
Questions? <ul style="list-style-type: none"> www.chloroprep.com call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST) 	

See other side for additional information

Front

ChloroPrep® One-Step
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
 1.5ml FREPP® APPLICATOR

Cat. No. 260299NSB / 260599NSB
 NDC 054365-400-02

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

See Drug Facts for Full Labeling Information



Do Not Reuse



Not made with natural rubber latex

External Use Only

Professional Use Only

CareFusion, El Paso, TX 79912

Artwork 6-775403F · MXXXX
 Made in the USA of US and imported parts

Back

Frepp Cyrel Lidding

Exp. Date xx/xx
Lot No. xxxxx

Cat. No. 260299 NDC 054365-400-02

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

1.5ml FREPP® APPLICATOR



WARNING. FLAMMABLE.

KEEP AWAY FROM FIRE OR FLAME.




Applicator is sterile if package is intact

CareFusion, El Paso, TX 79912
1-800-523-0502 (M-F 8 a.m.- 5 p.m. CST)

Artwork No. 6-000299 • LXXXX
Made in the USA of US and imported parts

Bend and
Peel

Frepp Carton – Top Panel



ChloroPrep[®] One-Step

WARNING: FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/vAntiseptic
Isopropyl alcohol 70% v/vAntiseptic

Purposes

.....Antiseptic
.....Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame.
 ■ do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator horizontally with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds. Allow the area to air dry for approximately 30 seconds.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes. Allow the area to air dry for approximately 1 minute.
- do not blot or wipe away
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients USP purified water

Questions?

- www.chloroprep.com
- call 1-800-523-0502 (M–F 8 a.m.–5 p.m. CST)

NDC 054365-400-02

1.5ml
FREPP®
APPLICATORS

Clear

Do Not Reuse

~~Latex~~ Not made with natural rubber latex

STERILE EO

Applicator is sterile if package is intact


20 applicators

0.05 fl. oz (1.5 ml) each

ChlorPrep®

One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)


Patient Preoperative Skin Preparation



Professional Use Only

Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat
above 40 °C (104 °F)

Frepp Carton – Side Panel A



ChloroPrep[®]
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

1.5ml
FREPP[®] APPLICATORS

Clear

CareFusion
El Paso, TX 79912
1.800.523.0502
www.chloroprep.com
Made in the USA of US and imported parts

Frepp Carton – Side Panel B

ChloroPrep[®]
One-Step[®]
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
1.5ml
FREPP[®] APPLICATORS
Clear

LOT XXXXXX EXP DATE: XX/XX
(printed in-line)

No Varnish



(01) 20354365400024

Cat. No. 260299
©2009 CareFusion. All rights reserved. ChloroPrep is a registered trademark of CareFusion or one of its subsidiaries.

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1ml APPLICATOR

 Not made with natural rubber latex

 Do Not Reuse

Professional Use Only
External Use Only

Cat. No. 260479 / 260479NS / 260479NSB NDC 054365-400-15

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v.....Antiseptic
Isopropyl alcohol 70% v/v.....Antiseptic

Purposes

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame

- do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic.
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute.
- do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients USP purified water

Questions?

- www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)


CareFusion, El Paso, TX 79912 • Artwork 6-775479 • CXXXX • Made in the USA of US and imported parts

Back

1ml Cyrel Lidding



1ml Carton – Top Panel



ChloroPrep[®] One-Step

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active Ingredients

Chlorhexidine gluconate 2% w/v.....Antiseptic
Isopropyl alcohol 70% v/v.....Antiseptic

Purposes

.....Antiseptic
.....Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame
do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²).
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings *only once* to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds. Allow the area to air dry for approximately 30 seconds.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes. Allow the area to air dry for approximately 1 minute.
- do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)


Inactive ingredients

USP purified water

Questions?

- www.chloroprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

ChloroPrep®
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
**Patient Preoperative
Skin Preparation**



Professional Use Only
Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat
above 40 °C (104 °F)

NDC 054365-400-15
1 ml APPLICATORS
Clear

 Do Not Reuse
 Not made with natural rubber latex
STERILE


Applicator is sterile if package is intact
20 applicators
0.03 fl. oz. (1 ml) each

1ml Carton – Side Panel A

Chloraprep[®]
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
1 ml APPLICATORS
Clear

CareFusion
El Paso, TX 79912
1.800.523.0502
www.chloraprep.com
Made in the USA of US and imported parts

1ml Carton – Side Panel B



<div style="background-color: #f00; color: white; text-align: center; padding: 2px;">⚠ WARNING</div> <div style="background-color: #f00; color: white; text-align: center; padding: 2px;">FLAMMABLE</div> <div style="display: flex; align-items: center; margin-top: 5px;"> <p>Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet materials from prep area </div>	<p>Cat. No. 260415 / 260415NS / 260415NSB NDC 054365-400-11 Cat. No. 260400 / 260400NS / 260500NS NDC 054365-400-01</p> <p>ChloroPrep® One-Step ChloroPrep® With Tint 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation</p> <p style="text-align: center;">3 ml APPLICATOR </p> <p>Professional Use Only Do Not Reuse Not made with natural rubber latex</p>						
<div style="border: 2px solid black; padding: 5px;"> <h3 style="margin: 0;">Drug Facts</h3> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 60%;">Active ingredients</td> <td style="width: 40%; text-align: right;">Purposes</td> </tr> <tr> <td>Chlorhexidine gluconate 2% w/v.....</td> <td style="text-align: right;">Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% w/v.....</td> <td style="text-align: right;">Antiseptic</td> </tr> </table> <p>Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.</p> <p>Warnings</p> <p>For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area <p>Do not use</p> <ul style="list-style-type: none"> ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser </div>		Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v.....	Antiseptic	Isopropyl alcohol 70% w/v.....	Antiseptic
Active ingredients	Purposes						
Chlorhexidine gluconate 2% w/v.....	Antiseptic						
Isopropyl alcohol 70% w/v.....	Antiseptic						

Chloraprep® One-Step / Chloraprep® With Tint
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.
- **if using an ignition source, allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) ■ the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.

Inactive ingredients FD&C green #3 dye (Scrub-Teal applicators only), FD&C yellow #6 dye (Hi-Lite Orange applicators only), USP purified water

Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, El Paso, TX 79912 • Artwork 6-775400• FXXXX • Made in the USA of US and imported parts

Back

3ml Clear Lidding

at this end or pac

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

3ml APPLICATOR Clear

Use
For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- tuck prep towels to absorb solution, and then remove
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²). Discard the applicator after a single use.
- hold the applicator with sponge down and pinch the wings on the applicator once to break the ampule and release the antiseptic. Do not touch the sponge. Wet the sponge by repeatedly pressing and releasing the sponge against the treatment area until liquid is visible on the skin.
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Completely wet the treatment area with antiseptic. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Completely wet the treatment area with antiseptic. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.
- **if using an ignition source, allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.


Important: See insert for complete Drug Facts information

  **STERILE** Professional Use Only

Do Not Reuse
Applicator is sterile if package is intact.
Cat. No. 260400 NDC 054365-400-01
CareFusion, El Paso, TX 79912
call 1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST) • www.chloroprep.com
Made in the USA of US and imported parts
Artwork No. 6-786315 • KXXXX

3ml Clear Carton – Top Panel

ChloroPrep[®]
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative
Skin Preparation



3 ml
APPLICATORS

Clear

25 applicators

3ml Clear Carton – Back Panel

ChloroPrep®
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% w/v isopropyl alcohol (IPA)
Patient-Preoperative Skin Preparation

3 ml

APPLICATORS

Clear

25 applicators

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active Ingredients
Chlorhexidine gluconate 2% w/v Antiseptic
Isopropyl alcohol 70% w/v Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not clamp or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not use on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lacerations, punctures or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product, keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if: irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area.
- use the applicator. The applicator is approximately 4 in. x 5 in. (100 cm²).
- remove applicator from package, do not touch sponge.
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin.
- completely wet the treatment area with antiseptic.
- dry surgical sites (e.g., abdomen or arm); use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or moist surgical sites.
- moist surgical sites (e.g., inguinal fold); use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.
- if using an ignition source, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store at 20° to 25° C (68° to 77° F)
- avoid freezing and excessive heat above 40° C (104° F)

Inactive Ingredients USP purified water

Questions?

- visit us at www.usp.com
- call 1-800-822-5922 (M-F 8 a.m.-5 p.m. CST)

3ml Clear Carton Drug Facts

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v
Isopropyl alcohol 70% v/v

Purposes

Antiseptic
Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbur puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area.
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm); use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away.
- moist surgical sites (e.g., inguinal fold); use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.
- if using an ignition source, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

USP purified water

Questions?

- www.chlorprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

3ml Clear Carton – Front Panel

NDC 054365-400-01

3 ml

APPLICATORS

Clear

 Do Not Reuse

 Not made with natural rubber latex

STERILE EO

Applicator is sterile if package is intact

Professional Use Only

Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat above 40 °C (104 °F)

25 applicators
0.10 fl. oz. (3 ml) each

ChloroPrep[®]

One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)


Patient Preoperative Skin Preparation

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hair or eyes.
- Hair may take up to 1 hour to dry. Wet hair is flammable.
- do not drink or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area




3ml Clear Carton – Side Panel A

ChloroPrep[®]
One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

3 ml
APPLICATORS

Clear



CareFusion
El Paso, TX 79912
1.800.523.0502
www.chloraprep.com
Made in the USA of US and imported parts


3ml Clear Carton – Side Panel B

ChloroPrep[®]
One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

3 ml
APPLICATORS

Clear



LOT XXXXXX

(printed in-line)
No Varnish

EXP DATE XX/XX



(01) 20351365400017

Cat. No. 260400

©2009 CareFusion, All rights reserved. ChloroPrep is a registered trademark of CareFusion or one of its subsidiaries.

3ml Orange Lidding

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

3ml APPLICATOR **Hi-Lite Orange***

Use

For the preparation of the patient's skin prior to surgery.
Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- tuck prep towels to absorb solution, and then remove
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²). Discard the applicator after a single use.
- hold the applicator with sponge down and pinch the wings on the applicator once to break the ampule and release the antiseptic. Do not touch the sponge. Wet the sponge by repeatedly pressing and releasing the sponge against the treatment area until liquid is visible on the skin.
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Completely wet the treatment area with antiseptic. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Completely wet the treatment area with antiseptic. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.
- **if using an ignition source, allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.

Important: See insert for complete Drug Facts information



Do Not Reuse



Professional Use Only

Applicator is sterile if package is intact.

Cat. No. 260400 NDC 054365-400-01


CareFusion, El Paso, TX 79912

call 1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST) • www.chloroprep.com

Made in the USA of US and imported parts

Artwork No. 6-786415 • JXXXX

3ml Orange Carton – Top Panel



ChloroPrep®
With Tint
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation

3 ml
APPLICATORS

Hi-Lite Orange®

25 applicators

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active Ingredients

Chlorhexidine gluconate 2% w/v
 Isopropyl alcohol 70% v/v

Purposes

Antiseptic
 Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

- For external use only. Flammable. Keep away from fire or flame. To reduce the risk of fire, **PREP CAREFULLY**:
- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lacerations or in contact with the mucous membranes
- on open skin wounds or on a general skin cleanser

When using this product, keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- wipe applicator sites (e.g., abdomen or arm); use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or rub.
- moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.
- **If using an ignition source, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.**
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive Ingredients FDCC yellow #6 dye, USP purified water

Questions?

- www.chloroprep.com
- call 1-800-593-5692 (M-F 8 a.m. - 5 p.m. CST)

3ml Orange Carton – Drug Facts

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v
Isopropyl alcohol 70% v/v

Purposes
..... Antiseptic
..... Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

- Do not use
- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
 - for lumbal puncture or in contact with the meninges
 - on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away.
- moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.
- **if using an ignition source, allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients FD&C yellow #6 dye, USP purified water

Questions?

- www.chlorprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

3ml Orange Carton – Front Panel

NDC 054365-400-11

3 ml

APPLICATORS


Hi-Lite Orange®

 Do Not Reuse
 Not made with natural rubber latex
 **STERILE EO**
 Applicator is sterile if package is intact

Professional Use Only

Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat above 40 °C (104 °F)

25 applicators
0.10 fl. oz. (3 ml) each



ChloroPrep®

with Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

⚠ WARNING

FLAMMABLE

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off flammable vapors
- avoid getting solution into hairy areas.
- hair may take up to 1 hour to dry. **Hair is flammable.**
- do not drape or use ignition source (e.g. cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

3ml Orange Carton – Side Panel A

ChloroPrep[®]
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

3 ml
APPLICATORS

Hi-Lite Orange[®]



CareFusion
El Paso, TX 79912
1.800.523.0502
www.chloroprep.com
Made in the USA of US and imported parts


3ml Orange Carton – Side Panel B

ChloroPrep[®]
With Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

3 ml
APPLICATORS

Hi-Lite Orange[®]



LOT XXXXXX

(printed in-line)
No Varnish

EXP DATE XXXXX



(01) 20354365400116

Cat. No. 260415
©2009 CareFusion, All rights reserved. ChloroPrep is a registered trademark of CareFusion or one of its subsidiaries.

3ml Teal Lidding

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Chloraprep® With Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

3ml APPLICATOR

Scrub Teal®

Use

For the preparation of the patient's skin prior to surgery.
Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- tuck prep towels to absorb solution, and then remove
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²). Discard the applicator after a single use.
- hold the applicator with sponge down and pinch the wings on the applicator once to break the ampule and release the antiseptic. Do not touch the sponge. Wet the sponge by repeatedly pressing and releasing the sponge against the treatment area until liquid is visible on the skin.
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Completely wet the treatment area with antiseptic. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Completely wet the treatment area with antiseptic. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.
- **if using an ignition source, allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.

Important: See insert for complete Drug Facts information



Do Not Reuse



STERILE

Professional Use Only

Applicator is sterile if package is intact.

Cat. No. 260400 NDC 054365-400-01


CareFusion, El Paso, TX 79912

call 1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST) • www.chloraprep.com

Made in the USA of US and imported parts

Artwork No. 6-786425 • FXXXX

3ml Teal Carton – Top Panel



WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients
 Chlorhexidine gluconate 2% w/v
 Isopropyl alcohol 70% v/v

Purposes
 Antiseptic
 Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, **PREP CAREFULLY**:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbopuncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away.
- moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away.
- **if using an ignition source, allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients FD&C green #3 dye, USP purified water

Questions?

- www.chloraprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

NDC 054365-400-10

3 ml

APPLICATORS

Scrub Teal®

Do Not Reuse
 Not made with natural rubber latex

STERILE EO

 Applicator is sterile if package is intact
 Professional Use Only
 Store between 15-30 °C (59-86 °F)
 Avoid freezing and excessive heat above 40 °C (104 °F)

25 applicators
0.10 fl. oz. (3 ml) each



ChloraPrep®

with Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas.
Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g. cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

3ml Teal Carton – Side Panel A

ChloroPrep[®]
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

3 ml
APPLICATORS

Scrub Teal[®]



CareFusion
El Paso, TX 79912
1.800.523.0502
www.chloraprep.com
Made in the USA of US and imported parts


3ml Teal Carton – Side Panel B

ChloroPrep[®]
With Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

3 ml
APPLICATORS

Scrub Teal[®]



LOT XXXXXX

(printed in-line)
No Varnish

EXP DATE XXXXX

BARCODE PLACEHOLDER

Cat. No. 260425
©2009 CareFusion, All rights reserved. ChloroPrep is a registered trademark of CareFusion or one of its subsidiaries.