

Approved for FDA Submission  
L Kindwald 10/05/12

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use dexmedetomidine hydrochloride safely and effectively. See full prescribing information for Precedex.

Precedex (dexmedetomidine hydrochloride) injection  
For intravenous infusion

Initial U.S. Approval: 1999

INDICATIONS AND USAGE

Precedex is a relatively selective alpha<sub>2</sub>-adrenergic agonist indicated for:

- Sedation of initially intubated and mechanically ventilated patients during treatment in an intensive care setting. Administer Precedex by continuous infusion not to exceed 24 hours. (1.1)
- Sedation of non-intubated patients prior to and/or during surgical and other procedures. (1.2)

DOSAGE AND ADMINISTRATION

- Individualize and titrate Precedex dosing to desired clinical effect. (2.1)
- Administer Precedex using a controlled infusion device. (2.1)
- Dilute vial contents in 0.9% sodium chloride solution to achieve required concentration (4 mcg/mL) prior to administration. (2.4)

**For Intensive Care Unit Sedation:** Generally initiate at 1 mcg/kg over 10 minutes, followed by a maintenance infusion of 0.2 to 0.7 mcg/kg/hr. (2.2)

**For Procedural Sedation:** Generally initiate at 1 mcg/kg over 10 minutes, followed by a maintenance infusion initiated at 0.6 mcg/kg/hr and titrated to achieve desired clinical effect with doses ranging from 0.2 to 1 mcg/kg/hr.

**Alternative Doses:** Recommended for patients over 65 years of age and awake fiberoptic intubation patients. (2.2)

DOSAGE FORMS AND STRENGTHS

Precedex Injection, Concentrate, 200 mcg/2 mL (100 mcg/mL) in a glass vial. (3)

Precedex Injection 200 mcg/50 mL (4 mcg/mL) in a 50 mL glass bottle. (3)

Precedex Injection 400 mcg/100 mL (4 mcg/mL) in a 100 mL glass bottle. (3)

CONTRAINDICATIONS

None. (4)

WARNINGS AND PRECAUTIONS

- Monitoring: Continuously monitor patients while receiving Precedex. (5.1)
- Bradycardia and Sinus Arrest: Have occurred in young healthy volunteers with high vagal tone or with different routes of administration, e.g., rapid intravenous or bolus administration. (5.2)
- Hypotension and Bradycardia: May necessitate medical intervention. May be more pronounced in patients with hypovolemia, diabetes mellitus, or chronic hypotension, and in the elderly. Use with caution in patients with advanced heart block or severe ventricular dysfunction. (5.2)
- Co-administration with Other Vasodilators or Negative Chronotropic Agents: Use with caution due to additive pharmacodynamic effects. (5.2)
- Transient Hypertension: Observed primarily during the loading dose. Consider reduction in loading infusion rate. (5.3)
- Arousability: Patients can become aroused/alert with stimulation; this alone should not be considered as lack of efficacy. (5.4)
- Prolonged exposure to dexmedetomidine beyond 24 hours may be associated with tolerance and tachyphylaxis and a dose-related increase in adverse events. (5.6)

ADVERSE REACTIONS

- The most common adverse reactions (incidence greater than 2%) are hypotension, bradycardia, and dry mouth. (6.1)
- Adverse reactions associated with infusions greater than 24 hours in duration include ARDS, respiratory failure, and agitation. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Hospira, Inc. at 1-800-441-4100 or electronically at [ProductComplaintsPP@hospira.com](mailto:ProductComplaintsPP@hospira.com), or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

DRUG INTERACTIONS

Anesthetics, Sedatives, Hypnotics, Opioids: Enhancement of pharmacodynamic effects. Reduction in dosage of Precedex or the concomitant medication may be required. (7.1)

USE IN SPECIFIC POPULATIONS

- Geriatric Patients: Dose reduction should be considered. (2.2, 2.3, 5.1, 8.5)
- Hepatic Impairment: Dose reduction should be considered. (2.1, 2.2, 2.3, 5.7, 8.6)
- Pregnancy: Based on animal data, may cause fetal harm. (8.1)
- Nursing Mothers: Caution should be exercised when administered to a nursing woman. (8.3)

See 17 for PATIENT COUNSELING INFORMATION

Revised: 09/2012

FULL PRESCRIBING INFORMATION: CONTENTS\*

1	INDICATIONS AND USAGE	8	USE IN SPECIFIC POPULATIONS
1.1	Intensive Care Unit Sedation	8.1	Pregnancy
1.2	Procedural Sedation	8.2	Labor and Delivery
2	DOSAGE AND ADMINISTRATION	8.3	Nursing Mothers
2.1	Dosing Guidelines	8.4	Pediatric Use
2.2	Dosage Information	8.5	Geriatric Use
2.3	Dosage Adjustment	8.6	Hepatic Impairment
2.4	Preparation of Solution	9	DRUG ABUSE AND DEPENDENCE
2.5	Administration with Other Fluids	9.1	Controlled Substance
2.6	Compatibility with Natural Rubber	10	OVERDOSAGE
3	DOSAGE FORMS AND STRENGTHS	11	DESCRIPTION
4	CONTRAINDICATIONS	12	CLINICAL PHARMACOLOGY
5	WARNINGS AND PRECAUTIONS	12.1	Mechanism of Action
5.1	Drug Administration	12.2	Pharmacodynamics
5.2	Hypotension, Bradycardia, and Sinus Arrest	12.3	Pharmacokinetics
5.3	Transient Hypertension	13	NONCLINICAL TOXICOLOGY
5.4	Arousability	13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
5.5	Withdrawal	13.2	Animal Pharmacology and/or Toxicology
5.6	Tolerance and Tachyphylaxis	14	CLINICAL STUDIES
5.7	Hepatic Impairment	14.1	Intensive Care Unit Sedation
6	ADVERSE REACTIONS	14.2	Procedural Sedation
6.1	Clinical Studies Experience	16	HOW SUPPLIED/STORAGE AND HANDLING
6.2	Postmarketing Experience	17	PATIENT COUNSELING INFORMATION
7	DRUG INTERACTIONS	17	PATIENT COUNSELING INFORMATION
7.1	Anesthetics, Sedatives, Hypnotics, Opioids	*	Sections or subsections omitted from the full prescribing information are not listed
7.2	Neuromuscular Blockers		

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Intensive Care Unit Sedation

Precedex™ is indicated for sedation of initially intubated and mechanically ventilated patients during treatment in an intensive care setting. Precedex should be administered by continuous infusion not to exceed 24 hours.

Precedex has been continuously infused in mechanically ventilated patients prior to extubation, during extubation, and post-extubation. It is not necessary to discontinue Precedex prior to extubation.

1.2 Procedural Sedation

Precedex is indicated for sedation of non-intubated patients prior to and/or during surgical and other procedures.

2 DOSAGE AND ADMINISTRATION

2.1 Dosing Guidelines

- Precedex dosing should be individualized and titrated to desired clinical response.
- Precedex is not indicated for infusions lasting longer than 24 hours.
- Precedex should be administered using a controlled infusion device.

2.2 Dosage Information

Table 1: Dosage Information	
INDICATION	DOSAGE AND ADMINISTRATION
Initiation of Intensive Care Unit Sedation	<b>For adult patients:</b> a loading infusion of 1 mcg/kg over 10 minutes.
	<b>For patients being converted from alternate sedative therapy:</b> a loading dose may not be required [see <i>Dosage and Administration: Maintenance of Intensive Care Unit Sedation</i> (2.2)].
	<b>For patients over 65 years of age:</b> a dose reduction should be considered [see <i>Use in Specific Populations</i> (8.5)].
	<b>For patients with impaired hepatic function:</b> a dose reduction should be considered [see <i>Use in Specific Populations</i> (8.6), <i>Clinical Pharmacology</i> (12.3)].
Maintenance of Intensive Care Unit Sedation	<b>For adult patients:</b> a maintenance infusion of 0.2 to 0.7 mcg/kg/hr. The rate of the maintenance infusion should be adjusted to achieve the desired level of sedation.
	<b>For patients over 65 years of age:</b> a dose reduction should be considered [see <i>Use in Specific Populations</i> (8.5)].
	<b>For patients with impaired hepatic function:</b> a dose reduction should be considered [see <i>Use in Specific Populations</i> (8.6), <i>Clinical Pharmacology</i> (12.3)].
	<b>For awake fiberoptic intubation patients:</b> a loading infusion of 1 mcg/kg over 10 minutes.
Initiation of Procedural Sedation	<b>For adult patients:</b> a loading infusion of 1 mcg/kg over 10 minutes. For less invasive procedures such as ophthalmic surgery, a loading infusion of 0.5 mcg/kg given over 10 minutes may be suitable.
	<b>For awake fiberoptic intubation patients:</b> a loading infusion of 1 mcg/kg over 10 minutes.
	<b>For patients over 65 years of age:</b> a loading infusion of 0.5 mcg/kg over 10 minutes [see <i>Use in Specific Populations</i> (8.5)].
	<b>For patients with impaired hepatic function:</b> a dose reduction should be considered [see <i>Use in Specific Populations</i> (8.6), <i>Clinical Pharmacology</i> (12.3)].
Maintenance of Procedural Sedation	<b>For adult patients:</b> the maintenance infusion is generally initiated at 0.6 mcg/kg/hr and titrated to achieve desired clinical effect with doses ranging from 0.2 to 1 mcg/kg/hr. The rate of the maintenance infusion should be adjusted to achieve the targeted level of sedation.
	<b>For awake fiberoptic intubation patients:</b> a maintenance infusion of 0.7 mcg/kg/hr is recommended until the endotracheal tube is secured.
	<b>For patients over 65 years of age:</b> a dose reduction should be considered [see <i>Use in Specific Populations</i> (8.5)].
	<b>For patients with impaired hepatic function:</b> a dose reduction should be considered [see <i>Use in Specific Populations</i> (8.6), <i>Clinical Pharmacology</i> (12.3)].

2.3 Dosage Adjustment

Due to possible pharmacodynamic interactions, a reduction in dosage of Precedex or other concomitant anesthetics, sedatives, hypnotics or opioids may be required when co-administered [see *Drug Interactions* (7.1)].

Dosage reductions may need to be considered for patients with hepatic impairment, and geriatric patients [see *Warnings and Precautions* (5.7), *Use in Specific Populations* (8.6), *Clinical Pharmacology* (12.3)].

2.4 Preparation of Solution

Strict aseptic technique must always be maintained during handling of Precedex. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

Precedex Injection, Concentrate, 200 mcg/2 mL (100 mcg/mL)

Precedex must be diluted in 0.9% sodium chloride solution to achieve required concentration (4 mcg/mL) prior to administration. Preparation of solutions is the same, whether for the loading dose or maintenance infusion.

To prepare the infusion, withdraw 2 mL of Precedex and add to 48 mL of 0.9% sodium chloride injection to a total of 50 mL. Shake gently to mix well.

Precedex Injection, 200 mcg/50 mL (4 mcg/mL) and 400 mcg/100 mL (4 mcg/mL)

Precedex Injection is supplied in glass containers containing a premixed, ready to use dexmedetomidine hydrochloride solution in 0.9% sodium chloride in water. No further dilution of these preparations are necessary.

2.5 Administration with Other Fluids

Precedex infusion should not be co-administered through the same intravenous catheter with blood or plasma because physical compatibility has not been established.

Precedex has been shown to be incompatible when administered with the following drugs: amphotericin B, diazepam.

Precedex has been shown to be compatible when administered with the following intravenous fluids:

- 0.9% sodium chloride in water
- 5% dextrose in water
- 20% mannitol
- Lactated Ringer's solution
- 100 mg/mL magnesium sulfate solution
- 0.3% potassium chloride solution

2.6 Compatibility with Natural Rubber

Compatibility studies have demonstrated the potential for absorption of Precedex to some types of natural rubber. Although Precedex is dosed to effect, it is advisable to use administration components made with synthetic or coated natural rubber gaskets.

3 DOSAGE FORMS AND STRENGTHS

Precedex Injection, Concentrate

Precedex Injection, 200 mcg/2 mL (100 mcg/mL) in a glass vial

Precedex Injection

Precedex Injection, 200 mcg/50 mL (4 mcg/mL) in a 50 mL glass bottle

Precedex Injection, 400 mcg/100 mL (4 mcg/mL) in a 100 mL glass bottle

4 CONTRAINDICATIONS

None

5 WARNINGS AND PRECAUTIONS

5.1 Drug Administration

Precedex should be administered only by persons skilled in the management of patients in the intensive care or operating room setting. Due to the known pharmacological effects of Precedex, patients should be continuously monitored while receiving Precedex.

5.2 Hypotension, Bradycardia, and Sinus Arrest

Clinically significant episodes of bradycardia and sinus arrest have been reported with Precedex administration in young, healthy volunteers with high vagal tone or with different routes of administration including rapid intravenous or bolus administration.

Reports of hypotension and bradycardia have been associated with Precedex infusion. If medical intervention is required, treatment may include decreasing or stopping the infusion of Precedex, increasing the rate of intravenous fluid administration, elevation of the lower extremities, and use of pressor agents. Because Precedex has the potential to augment bradycardia induced by vagal stimuli, clinicians should be prepared to intervene. The intravenous administration of anticholinergic agents (e.g., glycopyrrolate, atropine) should be considered to modify vagal tone. In clinical trials, glycopyrrolate or atropine were effective in the treatment of most episodes of Precedex-induced bradycardia. However, in some patients with significant cardiovascular dysfunction, more advanced resuscitative measures were required.

Caution should be exercised when administering Precedex to patients with advanced heart block and/or severe ventricular dysfunction. Because Precedex decreases sympathetic nervous system activity, hypotension and/or bradycardia may be expected to be more pronounced in patients with hypovolemia, diabetes mellitus, or chronic hypotension and in elderly patients.

In clinical trials where other vasodilators or negative chronotropic agents were co-administered with Precedex an additive pharmacodynamic effect was not observed. Nonetheless, caution should be used when such agents are administered concomitantly with Precedex.

5.3 Transient Hypertension

Transient hypertension has been observed primarily during the loading dose in association with the initial peripheral vasoconstrictive effects of Precedex. Treatment of the transient hypertension has generally not been necessary, although reduction of the loading infusion rate may be desirable.

5.4 Arousability

Some patients receiving Precedex have been observed to be arousable and alert when stimulated. This alone should not be considered as evidence of lack of efficacy in the absence of other clinical signs and symptoms.

5.5 Withdrawal

**Intensive Care Unit Sedation**  
With administration up to 7 days, regardless of dose, 12 (5%) Precedex subjects experienced at least 1 event related to withdrawal within the first 24 hours after discontinuing study drug and 7 (3%) Precedex subjects experienced at least 1 event 24 to 48 hours after end of study drug. The most common events were nausea, vomiting, and agitation.

Tachycardia and hypertension requiring intervention in the 48 hours following study drug discontinuation occurred at frequencies of <5%. If tachycardia and/or hypertension occurs after discontinuation of Precedex supportive therapy is indicated.

Procedural Sedation

Withdrawal symptoms were not seen after discontinuation of short term infusions of Precedex (<6 hours).

5.6 Tolerance and Tachyphylaxis

Use of dexmedetomidine beyond 24 hours has been associated with tolerance and tachyphylaxis and a dose-related increase in adverse reactions [see *Adverse Reactions* (6.1)].

5.7 Hepatic Impairment

Since Precedex clearance decreases with severity of hepatic impairment, dose reduction should be considered in patients with impaired hepatic function [see *Dosage and Administration* (2.2)].

6 ADVERSE REACTIONS

6.1 Clinical Studies Experience

Because clinical trials are conducted under widely varying conditions, adverse reactions rates observed in the clinical trials of a drug cannot be directly compared to rates in clinical trials of another drug and may not reflect the rates observed in practice.

Use of Precedex has been associated with the following serious adverse reactions:

- Hypotension, bradycardia and sinus arrest [see *Warnings and Precautions* (5.2)]
- Transient hypertension [see *Warnings and Precautions* (5.3)]

Most common treatment-emergent adverse reactions, occurring in greater than 2% of patients in both Intensive Care Unit and procedural sedation studies include hypotension, bradycardia and dry mouth.

Intensive Care Unit Sedation

Adverse reaction information is derived from the continuous infusion trials of Precedex for sedation in the Intensive Care Unit setting in which 1007 patients received Precedex. The mean total dose was 7.4 mcg/kg (range: 0.8 to 84.1), mean dose per hour was 0.5 mcg/kg/hr (range: 0.1 to 6.0) and the mean duration of infusion of 15.9 hours (range: 0.2 to 157.2). The population was between 17 to 88 years of age, 43% ≥65 years of age, 77% male and 93% Caucasian. Treatment-emergent adverse reactions occurring at an incidence of >2% are provided in Table 2. The most frequent adverse reactions were hypotension, bradycardia and dry mouth [see *Warnings and Precautions* (5.2)].

Table 2: Adverse Reactions with an Incidence >2%—Intensive Care Unit Sedation Population <24 hours\*

Adverse Event	All Precedex (N = 1007) (%)	Randomized Precedex (N = 798) (%)	Placebo (N = 400) (%)	Propofol (N = 188) (%)
Hypotension	25%	24%	12%	13%
Hypertension	12%	13%	19%	4%
Nausea	9%	9%	9%	11%
Bradycardia	5%	5%	3%	0
Atrial Fibrillation	4%	5%	3%	7%
Pyrexia	4%	4%	4%	4%
Dry Mouth	4%	3%	1%	1%
Vomiting	3%	3%	5%	3%
Hypovolemia	3%	3%	2%	5%
Atelectasis	3%	3%	3%	6%
Pleural Effusion	2%	2%	1%	6%
Agitation	2%	2%	3%	1%
Tachycardia	2%	2%	4%	1%
Anemia	2%	2%	2%	2%
Hyperthermia	2%	2%	3%	0
Chills	2%	2%	3%	2%
Hyperglycemia	2%	2%	2%	3%
Hypoxia	2%	2%	2%	3%
Post-procedural Hemorrhage	2%	2%	3%	4%
Pulmonary Edema	1%	1%	1%	3%
Hypocalcemia	1%	1%	0	2%
Acidosis	1%	1%	1%	2%
Urine Output Decreased	1%	1%	0	2%
Sinus Tachycardia	1%	1%	1%	2%
Ventricular Tachycardia	<1%	1%	1%	5%
Wheezing	<1%	1%	0	2%
Edema Peripheral	<1%	0	1%	2%

\* 26 subjects in the all Precedex group and 10 subjects in the randomized Precedex group had exposure for greater than 24 hours.

Adverse reaction information was also derived from the placebo-controlled, continuous infusion trials of Precedex for sedation in the surgical intensive care unit setting in which 387 patients received Precedex for less than 24 hours. The most frequently observed treatment-emergent adverse events included hypotension, hypertension, nausea, bradycardia, fever, vomiting, hypoxia, tachycardia and anemia [see Table 3].

Table 3: Treatment-Emergent Adverse Events Occurring in >1% Of All Dexmedetomidine-Treated Patients in the Randomized Placebo-Controlled Continuous Infusion <24 Hours ICU Sedation Studies

Adverse Event	Randomized Dexmedetomidine (N = 387) (%)	Placebo (N = 379) (%)
Hypotension	28%	13%
Hypertension	16%	18%
Nausea	11%	9%
Bradycardia	7%	3%
Fever	5%	4%
Vomiting	4%	6%
Atrial Fibrillation	4%	3%
Hypoxia	4%	4%
Tachycardia	3%	5%
Hemorrhage	3%	4%
Anemia	3%	2%
Dry Mouth	3%	1%
Rigors	2%	3%
Agitation	2%	3%
Hyperpyrexia	2%	3%
Pain	2%	2%
Hyperglycemia	2%	2%
Acidosis	2%	2%
Pleural Effusion	2%	1%
Oliguria	2%	<1%
Thirst	2%	<1%

In a controlled clinical trial, Precedex was compared to midazolam for ICU sedation exceeding 24 hours duration. Key treatment emergent adverse events occurring in dexmedetomidine or midazolam treated patients in the randomized active comparator continuous infusion long-term intensive care unit sedation study are provided in Table 4. The number (%) of subjects who had a dose-related increase in treatment-emergent adverse events by maintenance adjusted dose rate range in the Precedex group is provided in Table 5.

Table 4: Key Treatment-Emergent Adverse Events Occurring in Dexmedetomidine- or Midazolam-Treated Patients in the Randomized Active Comparator Continuous Infusion Long-Term Intensive Care Unit Sedation Study

Adverse Event	Dexmedetomidine (N = 244) (%)	Midazolam (N = 122) (%)
Hypotension <sup>1</sup>	56%	56%
Hypotension Requiring Intervention	28%	27%
Bradycardia <sup>2</sup>	42%	19%
Bradycardia Requiring Intervention	5%	1%
Systolic Hypertension <sup>3</sup>	28%	42%
Tachycardia <sup>4</sup>	25%	44%
Tachycardia Requiring Intervention	10%	10%
Diastolic Hypertension <sup>3</sup>	12%	15%
Hypertension <sup>3</sup>	11%	15%
Hypertension Requiring Intervention <sup>1</sup>	19%	30%
Hypokalemia	9%	13%
Pyrexia	7%	2%
Agitation	7%	6%
Hyperglycemia	7%	2%
Constipation	6%	6%
Hypoglycemia	5%	6%
Respiratory Failure	5%	3%
Renal Failure Acute	2%	1%
Acute Respiratory Distress Syndrome	2%	1%
Generalized Edema	2%	6%
Hypomagnesemia	1%	7%

<sup>1</sup> Includes any type of hypertension.

<sup>2</sup> Hypotension was defined in absolute terms as Systolic blood pressure of <80 mmHg or Diastolic blood pressure of <50 mmHg or in relative terms as ≤30% lower than pre-study drug infusion value.

<sup>3</sup> Bradycardia was defined in absolute terms as <40 bpm or in relative terms as ≤30% lower than pre-study drug infusion value.

<sup>4</sup> Hypertension was defined in absolute terms as Systolic blood pressure >180 mmHg or Diastolic blood pressure of >100 mmHg or in relative terms as ≥30% higher than pre-study drug infusion value.

<sup>5</sup> Tachycardia was defined in absolute terms as >120 bpm or in relative terms as ≥30% greater than pre-study drug infusion value.

The following adverse events occurred between 2 and 5% for Precedex and Midazolam, respectively: renal failure acute (2.5%, 0.8%), acute respiratory distress syndrome (2.5%, 0.8%), and respiratory failure (4.5%, 3.3%).

Table 5: Number (%) of Subjects Who Had a Dose-Related Increase in Treatment Emergent Adverse Events by Maintenance Adjusted Dose Rate Range in the Precedex Group



Table 7: Adverse Reactions Experienced During Post-approval Use of Precedex	
Body System	Preferred Term
Body as a Whole	Fever, hyperpyrexia, hypovolemia, light anesthesia, pain, rigors
Cardiovascular Disorders, General	Blood pressure fluctuation, heart disorder, hypertension, hypotension, myocardial infarction
Central and Peripheral Nervous System Disorders	Dizziness, headache, neuralgia, neuritis, speech disorder, convulsion
Gastrointestinal System Disorders	Abdominal pain, diarrhea, vomiting, nausea
Heart Rate and Rhythm Disorders	Arrhythmia, ventricular arrhythmia, bradycardia, hypoxia, atrioventricular block, cardiac arrest, extrasystoles, atrial fibrillation, heart block, t wave inversion, tachycardia, supraventricular tachycardia, ventricular tachycardia
Liver and Biliary System Disorders	Increased gamma-glutamyl transpeptidase, hepatic function abnormal, hyperbilirubinemia, alanine transaminase, aspartate aminotransferase
Metabolic and Nutritional Disorders	Acidosis, respiratory acidosis, hyperkalemia, increased alkaline phosphatase, thirst, hypoglycemia
Psychiatric Disorders	Agitation, confusion, delirium, hallucination, illusion
Red Blood Cell Disorders	Anemia
Renal Disorders	Blood urea nitrogen increased, oliguria
Respiratory System Disorders	Apnea, bronchospasm, dyspnea, hypercapnia, hypoventilation, hypoxia, pulmonary congestion
Skin and Appendages Disorders	Increased sweating
Vascular Disorders	Hemorrhage
Vision Disorders	Photopsia, abnormal vision

7 DRUG INTERACTIONS

7.1 Anesthetics, Sedatives, Hypnotics, Opioids

Co-administration of Precedex with anesthetics, sedatives, hypnotics, and opioids is likely to lead to an enhancement of effects. Specific studies have confirmed these effects with sevoflurane, isoflurane, propofol, alfentanil, and midazolam. No pharmacokinetic interactions between Precedex and isoflurane, propofol, alfentanil and midazolam have been demonstrated. However, due to possible pharmacodynamic interactions, when co-administered with Precedex, a reduction in dosage of Precedex or the concomitant anesthetic, sedative, hypnotic or opioid may be required.

7.2 Neuromuscular Blockers

In one study of 10 healthy volunteers, administration of Precedex for 45 minutes at a plasma concentration of 1 ng/mL resulted in no clinically meaningful increases in the magnitude of neuromuscular blockade associated with rocuronium administration.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C

There are no adequate and well-controlled studies of Precedex use in pregnant women. In an *in vitro* human placenta study, placental transfer of dexmedetomidine occurred. In a study in the pregnant rat, placental transfer of dexmedetomidine was observed when radiolabeled dexmedetomidine was administered subcutaneously. Thus, fetal exposure should be expected in humans, and Precedex should be used during pregnancy only if the potential benefits justify the potential risk to the fetus.

Teratogenic effects were not observed in rats following subcutaneous administration of dexmedetomidine during the period of fetal organogenesis (from gestation day 5 to 16) with doses up to 200 mcg/kg (representing a dose approximately equal to the maximum recommended human intravenous dose based on body surface area) or in rabbits following intravenous administration of dexmedetomidine during the period of fetal organogenesis (from gestation day 6 to 18) with doses up to 96 mcg/kg (representing approximately half the human exposure at the maximum recommended dose based on plasma area under the time-curve comparison). However, fetal toxicity, as evidenced by increased post-implantation losses and reduced live pups, was observed in rats at a subcutaneous dose of 200 mcg/kg. The no-effect dose in rats was 20 mcg/kg (representing a dose less than the maximum recommended human intravenous dose based on a body surface area comparison). In another reproductive toxicity study when dexmedetomidine was administered subcutaneously to pregnant rats at 8 and 32 mcg/kg (representing a dose less than the maximum recommended human intravenous dose based on a body surface area comparison) from gestation day 16 through weaning, lower offspring weights were observed. Additionally, when offspring of the 32 mcg/kg group were allowed to mate, elevated fetal and embryocidal toxicity and delayed motor development was observed in second generation offspring.

8.2 Labor and Delivery

The safety of Precedex during labor and delivery has not been studied.

8.3 Nursing Mothers

It is not known whether Precedex is excreted in human milk. Radio-labeled dexmedetomidine administered subcutaneously to lactating female rats was excreted in milk. Because many drugs are excreted in human milk, caution should be exercised when Precedex is administered to a nursing woman.

8.4 Pediatric Use

The efficacy, safety, and pharmacokinetics of Precedex in pediatric patients less than 18 years of age have not been established. Therefore, Precedex should not be used in this population.

8.5 Geriatric Use

Intensive Care Unit Sedation

A total of 729 patients in the clinical studies were 65 years of age and over. A total of 200 patients were 75 years of age and over. In patients greater than 65 years of age, a higher incidence of bradycardia and hypotension was observed following administration of Precedex [see *Warnings and Precautions* (5.2)]. Therefore a dose reduction may be considered in patients over 65 years of age [see *Dosage and Administration* (2.2) and *Clinical Pharmacology* (12.3)].

Procedural Sedation

A total of 131 patients in the clinical studies were 65 years of age and over. A total of 47 patients were 75 years of age and over. Hypotension occurred in a higher incidence in Precedex-treated patients 65 years or older (72%) and 75 years or older (74%) as compared to patients <65 years (47%). A reduced loading dose of 0.5 mcg/kg given over 10 minutes is recommended and a reduction in the maintenance infusion should be considered for patients greater than 65 years of age.

8.6 Hepatic Impairment

Since Precedex clearance decreases with increasing severity of hepatic impairment, dose reduction should be considered in patients with impaired hepatic function [see *Dosage and Administration* (2.2) and *Clinical Pharmacology* (12.3)].

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

Precedex (dexmedetomidine hydrochloride) is not a controlled substance.

9.3 Dependence

The dependence potential of Precedex has not been studied in humans. However, since studies in rodents and primates have demonstrated that Precedex exhibits pharmacologic actions similar to those of clonidine, it is possible that Precedex may produce a clonidine-like withdrawal syndrome upon abrupt discontinuation [see *Warnings and Precautions* (5.5)].


10 OVERDOSAGE

The tolerability of Precedex was studied in one study in which healthy subjects were administered doses at and above the recommended dose of 0.2 to 0.7 mcg/kg/hr. The maximum blood concentration achieved in this study was approximately 13 times the upper boundary of the therapeutic range. The most notable effects observed in two subjects who achieved the highest doses were first degree atrioventricular block and second degree heart block. No hemodynamic compromise was noted with the atrioventricular block and the heart block resolved spontaneously within one minute.

Five patients received an overdose of Precedex in the intensive care unit sedation studies. Two of these patients had no symptoms reported; one patient received a 2 mcg/kg loading dose over 10 minutes (twice the recommended loading dose) and one patient received a maintenance infusion of 0.8 mcg/kg/hr. Two other patients who received a 2 mcg/kg loading dose over 10 minutes, experienced bradycardia and/or hypotension. One patient who received a loading bolus dose of undiluted Precedex (19.4 mcg/kg), had cardiac arrest from which he was successfully resuscitated.

11 DESCRIPTION

Precedex (dexmedetomidine hydrochloride) injection is a sterile, nonpyrogenic solution suitable for intravenous infusion following dilution. Dexmedetomidine hydrochloride is the 5-enantiomer of medetomidine and is chemically described as (+)-4-(S)-[1-(2,3-dimethylphenyl)ethyl]-1H-imidazole monohydrochloride. Precedex has a molecular weight of 236.7 and the empirical formula is C<sub>13</sub>H<sub>16</sub>N<sub>2</sub> • HCl and the structural formula is:



Dexmedetomidine hydrochloride is a white or almost white powder that is freely soluble in water and has a pKa of 7.1. Its partition coefficient in-octanol: water at pH 7.4 is 2.89. Precedex is supplied as a clear, colorless, isotonic solution with a pH of 4.5 to 7.0. Each mL of Precedex Injection, Concentrate contains 118 mcg of dexmedetomidine hydrochloride equivalent to 100 mcg of dexmedetomidine and 9 mg of sodium chloride in water. Each mL of Precedex Injection contains 4.72 mcg of dexmedetomidine hydrochloride equivalent to 4 mcg dexmedetomidine and 9 mg of sodium chloride in water. The solution is preservative-free and contains no additives or chemical stabilizers.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Precedex is a relatively selective alpha<sub>2</sub>-adrenergic agonist with sedative properties. Alpha<sub>2</sub> selectivity is observed in animals following slow intravenous infusion of low and medium doses (10–300 mcg/kg). Both alpha<sub>1</sub> and alpha<sub>2</sub> activity is observed following slow intravenous infusion of high doses (≥1000 mcg/kg) or with rapid intravenous administration.

12.2 Pharmacodynamics

In a study in healthy volunteers (N = 10), respiratory rate and oxygen saturation remained within normal limits and there was no evidence of respiratory depression when Precedex was administered by intravenous infusion at doses within the recommended dose range (0.2–0.7 mcg/kg/hr).

12.3 Pharmacokinetics

Following intravenous administration, dexmedetomidine exhibits the following pharmacokinetic parameters: a rapid distribution phase with a distribution half-life (t<sub>1/2</sub>) of approximately 6 minutes; a terminal elimination half-life (t<sub>1/2</sub>) of approximately 2 hours; and steady-state volume of distribution (V<sub>ss</sub>) of approximately 118 liters. Clearance is estimated to be approximately 39 L/h. The mean body weight associated with this clearance estimate was 72 kg.

Dexmedetomidine exhibits linear pharmacokinetics in the dosage range of 0.2 to 0.7 mcg/kg/hr when administered by intravenous infusion for up to 24 hours. Table 8 shows the main pharmacokinetic parameters when Precedex was infused (after appropriate loading doses) at maintenance infusion rates of 0.17 mcg/kg/hr (target plasma concentration of 0.3 ng/mL) for 12 and 24 hours, 0.33 mcg/kg/hr (target plasma concentration of 0.6 ng/mL) for 24 hours, and 0.70 mcg/kg/hr (target plasma concentration of 1.25 ng/mL) for 24 hours.

Table 8: Mean ± SD Pharmacokinetic Parameters				
Parameter	Loading Infusion (min)/Total Infusion Duration (hrs)			
	10 min/12 hrs	10 min/24 hrs	10 min/24 hrs	35 min/24 hrs
	Precedex Target Plasma Concentration (ng/mL) and Dose (mcg/kg/hr)			
	0.3/0.17	0.3/0.17	0.6/0.33	1.25/0.70
t <sub>1/2</sub> <sup>a</sup> , hour	1.78 ± 0.30	2.22 ± 0.59	2.23 ± 0.21	2.50 ± 0.61
Cl, liter/hour	46.3 ± 8.3	43.1 ± 6.5	35.3 ± 6.8	36.5 ± 7.5
V <sub>ss</sub> , liter	88.7 ± 22.9	102.4 ± 20.3	93.6 ± 17.0	99.6 ± 17.8
Avg C <sub>ss</sub> <sup>b</sup> , ng/mL	0.27 ± 0.05	0.27 ± 0.05	0.67 ± 0.10	1.37 ± 0.20

<sup>a</sup> Presented as harmonic mean and pseudo standard deviation.

<sup>b</sup> Mean C<sub>ss</sub> = Average steady-state concentration of Precedex. The mean C<sub>ss</sub> was calculated based on post-dose sampling from 2.5 to 9 hours samples for 12-hour infusion and post-dose sampling from 2.5 to 18 hours for 24-hour infusions.

The loading doses for each of the above indicated groups were 0.5, 0.5, 1, and 2.2 mcg/kg, respectively.

Dexmedetomidine pharmacokinetic parameters after Precedex maintenance doses of 0.2 to 1.4 mcg/kg/hr for >24 hours were similar to the PK parameters after Precedex maintenance dosing for <24 hours in other studies. The values for clearance (CL), volume of distribution (V), and t<sub>1/2</sub> were 39.4 L/hr, 152 L, and 2.67 hours, respectively.

Distribution

The steady-state volume of distribution (V<sub>ss</sub>) of dexmedetomidine was approximately 118 liters. Dexmedetomidine protein binding was assessed in the plasma of normal healthy male and female subjects. The average protein binding was 94% and was constant across the different plasma concentrations tested. Protein binding was similar in males and females. The fraction of Precedex that was bound to plasma proteins was significantly decreased in subjects with hepatic impairment compared to healthy subjects.

The potential for protein binding displacement of dexmedetomidine by fentanyl, ketorolac, theophylline, digoxin and lidocaine was explored *in vitro*, and negligible changes in the plasma protein binding of Precedex were observed. The potential for protein binding displacement of phentanyl, warfarin, ibuprofen, propranolol, theophylline and digoxin by Precedex was explored *in vitro* and none of these compounds appeared to be significantly displaced by Precedex.

Metabolism

Dexmedetomidine undergoes almost complete biotransformation with very little unchanged dexmedetomidine excreted in urine and feces. Biotransformation involves both direct glucuronidation as well as cytochrome P450 mediated metabolism. The major metabolic pathways of dexmedetomidine are: direct N-glucuronidation to inactive metabolites; aliphatic hydroxylation (mediated primarily by CYP2A6) of dexmedetomidine to generate 3-hydroxy-dexmedetomidine, the glucuronide of 3-hydroxy-dexmedetomidine, and 3-carboxy-dexmedetomidine; and N-methylation of dexmedetomidine to generate 3-hydroxy N-methyl-dexmedetomidine, 3-carboxy N-methyl-dexmedetomidine, and dexmedetomidine-N-methyl O-glucuronide.

Elimination

The terminal elimination half-life (t<sub>1/2</sub>) of dexmedetomidine is approximately 2 hours and clearance is estimated to be approximately 39 L/h. A mass balance study demonstrated that after nine days an average of 95% of the radioactivity, following intravenous administration of radiolabeled dexmedetomidine, was recovered in the urine and 4% in the feces. No unchanged dexmedetomidine was detected in the urine. Approximately 85% of the radioactivity recovered in the urine was excreted within 24 hours after the infusion. Fractionation of the radioactivity excreted in urine demonstrated that products of N-glucuronidation accounted for approximately 34% of the cumulative urinary excretion. In addition, aliphatic hydroxylation of parent drug to form 3-hydroxy-dexmedetomidine, the glucuronide of 3-hydroxy-dexmedetomidine, and 3-carboxylic acid-dexmedetomidine together represented approximately 14% of the dose in urine. N-methylation of dexmedetomidine to form 3-hydroxy N-methyl dexmedetomidine, 3-carboxy N-methyl dexmedetomidine, and N-methyl O-glucuronide dexmedetomidine accounted for approximately 18% of the dose in urine. The N-Methyl metabolite itself was a minor circulating component and was undetected in urine. Approximately 28% of the urinary metabolites have not been identified.

Gender

There was no observed difference in Precedex pharmacokinetics due to gender.

Geriatrics

The pharmacokinetic profile of Precedex was not altered by age. There were no differences in the pharmacokinetics of Precedex in young (18–40 years), middle age (41–65 years), and elderly (>65 years) subjects.

Pediatrics

The pharmacokinetic profile of Precedex has not been studied in pediatric patients.

Hepatic Impairment

In subjects with varying degrees of hepatic impairment (Child-Pugh Class A, B, or C), clearance values for Precedex were lower than in healthy subjects. The mean clearance values for patients with mild, moderate, and severe hepatic impairment were 74%, 64% and 53% of those observed in the normal healthy subjects, respectively. Mean clearances for free drug were 59%, 51% and 32% of those observed in the normal healthy subjects, respectively.

Although Precedex is dosed to effect, it may be necessary to consider dose reduction in subjects with hepatic impairment [see *Dosage and Administration* (2.2), *Warnings and Precautions* (5.7)].

Renal Impairment

Precedex pharmacokinetics (C<sub>max</sub>, T<sub>max</sub>, AUC, t<sub>1/2</sub>, CL, and V<sub>ss</sub>) were not significantly different in patients with severe renal impairment (creatinine clearance: <30 mL/min) compared to healthy subjects.

Drug Interactions

*In vitro* studies: *In vitro* studies in human liver microsomes demonstrated no evidence of cytochrome P450 mediated drug interactions that are likely to be of clinical relevance.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Animal carcinogenicity studies have not been performed with dexmedetomidine. Dexmedetomidine was not mutagenic *in vitro*, in either the bacterial reverse mutation assay (*E. coli* and *Salmonella typhimurium*) or the mammalian cell forward mutation assay (mouse lymphoma). Dexmedetomidine was clastogenic in the *in vitro* human lymphocyte chromosome aberration test with, but not without, rat S9 metabolic activation. In contrast, dexmedetomidine was not clastogenic in the *in vitro* human lymphocyte chromosome aberration test with or without human S9 metabolic activation. Although dexmedetomidine was clastogenic in an *in vivo* mouse micronucleus test in NMRI mice, there was no evidence of clastogenicity in CD-1 mice.

Fertility in male or female rats was not affected after daily subcutaneous injections of dexmedetomidine at doses up to 54 mcg/kg (less than the maximum recommended human intravenous dose on a mcg/m<sup>2</sup> basis) administered from 10 weeks prior to mating in males, and 3 weeks prior to mating and during mating in females.

13.2 Animal Pharmacology and/or Toxicology

There were no differences in the adrenocorticotrophic hormone (ACTH)-stimulated cortisol response in dogs following a single dose of dexmedetomidine compared to saline control. However, after continuous subcutaneous infusions of dexmedetomidine at 3 mcg/kg/hr and 10 mcg/kg/hr for one week in dogs (exposures estimated to be within the clinical range), the ACTH-stimulated cortisol response was diminished by approximately 27% and 40%, respectively, compared to saline-treated control animals indicating a dose-dependent adrenal suppression.

14 CLINICAL STUDIES

The safety and efficacy of Precedex has been evaluated in four randomized, double-blind, placebo-controlled multicenter clinical trials in 1185 patients.

14.1 Intensive Care Unit Sedation

Two randomized, double-blind, parallel-group, placebo-controlled multicenter clinical trials included 754 patients being treated in a surgical intensive care unit. All patients were initially intubated and received mechanical ventilation. These trials evaluated the sedative properties of Precedex by comparing the amount of rescue medication (midazolam in one trial and propofol in the second) required to achieve a specified level of sedation (using the standardized Ramsay Sedation Scale) between Precedex and placebo from onset of treatment to extubation or to a total treatment duration of 24 hours. The Ramsay Level of Sedation Scale is displayed in Table 9.

Table 9: Ramsay Level of Sedation Scale	
Clinical Score	Level of Sedation Achieved
6	Asleep, no response
5	Asleep, sluggish response to light glabellar tap or loud auditory stimulus
4	Asleep, but with brisk response to light glabellar tap or loud auditory stimulus
3	Patient responds to commands
2	Patient cooperative, oriented, and tranquil
1	Patient anxious, agitated, or restless

In the first study, 175 patients were randomized to receive placebo and 178 to receive Precedex by intravenous infusion at a dose of 0.4 mcg/kg/hr (with allowed adjustment between 0.2 and 0.7 mcg/kg/hr) following an initial loading infusion of 1 mcg/kg intravenous over 10 minutes. The study drug infusion rate was adjusted to maintain a Ramsay sedation score of ≥3. Patients were allowed to receive “rescue” midazolam as needed to augment the study drug infusion. In addition, morphine sulfate

was administered for pain as needed. The primary outcome measure for this study was the total amount of rescue medication (midazolam) needed to maintain sedation as specified while intubated. Patients randomized to placebo received significantly more midazolam than patients randomized to Precedex (see Table 10).

A second prospective primary analysis assessed the sedative effects of Precedex by comparing the percentage of patients who achieved a Ramsay sedation score of ≥3 during intubation without the use of additional rescue medication. A significantly greater percentage of patients in the Precedex group maintained a Ramsay sedation score of ≥3 without receiving any midazolam rescue compared to the placebo group (see Table 10).

Table 10: Midazolam Use as Rescue Medication During Intubation (ITT)			
Study One	Placebo (N = 175)	Precedex (N = 178)	p-value
	19 mg	5 mg	0.0011*
Standard deviation	53 mg	19 mg	
Categorized Midazolam Use			
0 mg	43 (25%)	108 (61%)	<0.001**
0–4 mg	34 (19%)	36 (20%)	
>4 mg	98 (56%)	34 (19%)	

ITT (intent-to-treat) population includes all randomized patients.

\* ANOVA model with treatment center.

\*\* Chi-square.

A prospective secondary analysis assessed the dose of morphine sulfate administered to patients in the Precedex and placebo groups. On average, Precedex-treated patients received less morphine sulfate for pain than placebo-treated patients (0.47 versus 0.83 mg/h). In addition, 44% (79 of 178 patients) of Precedex patients received no morphine sulfate for pain versus 19% (33 of 175 patients) in the placebo group.

In a second study, 198 patients were randomized to receive placebo and 203 to receive Precedex by intravenous infusion at a dose of 0.4 mcg/kg/hr (with allowed adjustment between 0.2 and 0.7 mcg/kg/hr) following an initial loading infusion of 1 mcg/kg intravenous over 10 minutes. The study drug infusion was adjusted to maintain a Ramsay sedation score of ≥3. Patients were allowed to receive “rescue” propofol as needed to augment the study drug infusion. In addition, morphine sulfate was administered as needed for pain. The primary outcome measure for this study was the total amount of rescue medication (propofol) needed to maintain sedation as specified while intubated.

Patients randomized to placebo received significantly more propofol than patients randomized to Precedex (see Table 11).

A significantly greater percentage of patients in the Precedex group compared to the placebo group maintained a Ramsay sedation score of ≥3 without receiving any propofol rescue (see Table 11).

Table 11: Propofol Use as Rescue Medication During Intubation (ITT)			
Study Two	Placebo (N = 198)	Precedex (N = 203)	p-value
	513 mg	72 mg	<0.0001*
Standard deviation	782 mg	249 mg	
Categorized Propofol Use			
0 mg	47 (24%)	122 (60%)	<0.001**
0–50 mg	30 (15%)	43 (21%)	
>50 mg	121 (61%)	38 (19%)	

\* ANOVA model with treatment center.

\*\* Chi-square

A prospective secondary analysis assessed the dose of morphine sulfate administered to patients in the Precedex and placebo groups. On average, Precedex-treated patients received less morphine sulfate for pain than placebo-treated patients (0.43 versus 0.89 mg/h). In addition, 41% (83 of 203 patients) of Precedex patients received no morphine sulfate for pain versus 15% (30 of 198 patients) in the placebo group.

In a controlled clinical trial, Precedex was compared to midazolam for ICU sedation exceeding 24 hours duration. Precedex was not shown to be superior to midazolam for the primary efficacy endpoint, the percent of time patients were adequately sedated (81% versus 81%). In addition, administration of Precedex for longer than 24 hours was associated with tolerance, tachyphylaxis, and a dose-related increase in adverse events [see *Adverse Reactions* (6.1)].

14.2 Procedural Sedation

The safety and efficacy of Precedex for sedation of non-intubated patients prior to and/or during surgical and other procedures was evaluated in two randomized, double-blind, placebo-controlled multicenter clinical trials. Study 1 evaluated the sedative properties of Precedex in patients having a variety of elective surgeries/procedures performed under monitored anesthesia care. Study 2 evaluated Precedex in patients undergoing awake fiberoptic intubation prior to a surgical or diagnostic procedure.

In Study 1, the sedative properties of Precedex were evaluated by comparing the percent of patients not requiring rescue midazolam to achieve a specified level of sedation using the standardized Observer’s Assessment of Alertness/Sedation Scale (see Table 12).

Table 12: Observer’s Assessment of Alertness/Sedation				
Assessment Categories				
Responsiveness	Speech	Facial Expression	Eyes	Composite Score
Responds readily to name spoken in normal tone	Normal	Normal	Clear, no ptosis	5 (alert)
Lethargic response to name spoken in normal tone	Mild slowing or thickening	Mild relaxation	Glazed or mild ptosis (less than half the eye)	4
Responds only after name is called loudly and/or repeatedly	Slurring or prominent slowing	Marked relaxation (slack jaw)	Glazed and marked ptosis (half the eye or more)	3
Responds only after mild prodding or shaking	Few recognizable words	–	–	2
Does not respond to mild prodding or shaking	–	–	–	1 (deep sleep)

Patients were randomized to receive a loading infusion of either Precedex 1 mcg/kg, Precedex 0.5 mcg/kg, or placebo (normal saline) given over 10 minutes and followed by a maintenance infusion started at 0.6 mcg/kg/hr. The maintenance infusion of study drug could be titrated from 0.2 mcg/kg/hr to 1 mcg/kg/hr to achieve the targeted sedation score (Observer’s Assessment of Alertness/Sedation Scale ≤4). Patients were allowed to receive rescue midazolam as needed to achieve and/or maintain an Observer’s Assessment of Alertness/Sedation Scale ≤4. After achieving the desired level of sedation, a local or regional anesthetic block was performed. Demographic characteristics were similar

between the Precedex and comparator groups. Efficacy results showed that Precedex was more effective than the comparator group when used to sedate non-intubated patients requiring monitored anesthesia care during surgical and other procedures (see Table 13).

In Study 2, the sedative properties of Precedex were evaluated by comparing the percent of patients requiring rescue midazolam to achieve or maintain a specified level of sedation using the Ramsay Sedation Scale score ≥2 (see Table 9). Patients were randomized to receive a loading infusion of Precedex 1 mcg/kg or placebo (normal saline) given over 10 minutes and followed by a fixed maintenance infusion of 0.7 mcg/kg/hr. After achieving the desired level of sedation, topicalization of the airway occurred. Patients were allowed to receive rescue midazolam as needed to achieve and/or maintain a Ramsay Sedation Scale ≥2. Demographic characteristics were similar between the Precedex and comparator groups. For efficacy results see Table 13.

Table 13: Key Efficacy Results of Procedural Sedation Studies						
				Confidence <sup>b</sup>	Mean (SD)	
Study	Loading Infusion Treatment Arm	Number of Patients Enrolled <sup>a</sup>	% Not Requiring Midazolam Rescue	Interval on the Difference vs. Placebo	Total Dose (mg) of Rescue Midazolam Required	Confidence <sup>b</sup> Intervals of the Mean Rescue Dose
Study 1	Precedex 0.5 mcg/kg	134	40	37 (27, 48)	1.4 (1.7)	-2.7 (-3.4, -2.0)
	Precedex 1 mcg/kg	129	54	51 (40, 62)	0.9 (1.5)	-3.1 (-3.8, -2.5)
	placebo	63	3	—	4.1 (3.0)	—
Study 2	Precedex 1 mcg/kg	55	53	39 (20, 57)	1.1 (1.5)	-1.8 (-2.7, -0.9)
	placebo	50	14	—	2.9 (3.0)	—