



Rapamune[®] (sirolimus) Tablets

0.5 mg

Unit Dose - 100 Tablets
(10 Blister Cards of 10 Tablets Each)

ALWAYS DISPENSE WITH MEDICATION GUIDE



NDC 0008-1040-10

Rapamune[®] (sirolimus) Tablets

0.5 mg

For oral use only.

Unit Dose - 100 Tablets
(10 Blister Cards of 10 Tablets Each) **Rx only**



Rapamune[®] (sirolimus) Tablets

0.5 mg

Unit Dose - 100 Tablets
(10 Blister Cards of 10 Tablets Each)

12982100

DOSAGE AND USE: See accompanying prescribing information.

Store at room temperature, 20°C to 25°C (68°F to 77°F).
[See USP Controlled Room Temperature.] Protect from light.

This unit-dose package is not child-resistant. If dispensed for outpatient use, a child-resistant container should be utilized. Keep out of reach of children.

Distributed by
Wyeth Pharmaceuticals Inc
A subsidiary of Pfizer Inc
Philadelphia, PA 19101

MADE IN SWITZERLAND



Rapamune[®] (sirolimus) Tablets

0.5 mg

Unit Dose - 100 Tablets
(10 Blister Cards of 10 Tablets Each)



DOSAGE AND USE: See accompanying prescribing information.

Store at room temperature, 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.] Dispense in tight (USP), light-resistant (USP), and child-resistant containers.

Distributed by
Wyeth Pharmaceuticals Inc
A subsidiary of Pfizer Inc
Philadelphia, PA 19101
MADE IN SWITZERLAND

ALWAYS DISPENSE WITH MEDICATION GUIDE



NDC 0008-1040-05

Rapamune®
(sirolimus) Tablets

0.5 mg

For oral use only.

100 Tablets

Rx only



12112101



Rapamune[®] (sirolimus) Tablets

1 mg

Unit Dose - 100 Tablets
(10 Blister Cards of 10 Tablets Each)

ALWAYS DISPENSE WITH MEDICATION GUIDE



NDC 0008-1041-10

Rapamune[®] (sirolimus) Tablets

1 mg

For oral use only.

Unit Dose - 100 Tablets
(10 Blister Cards of 10 Tablets Each) **Rx only**



Rapamune[®] (sirolimus) Tablets

1 mg

Unit Dose - 100 Tablets
(10 Blister Cards of 10 Tablets Each)

12982200

DOSAGE AND USE: See accompanying prescribing information.

Store at room temperature, 20°C to 25°C (68°F to 77°F).
[See USP Controlled Room Temperature.] Protect from light.

This unit-dose package is not child-resistant. If dispensed for outpatient use, a child-resistant container should be utilized. Keep out of reach of children.

Distributed by
Wyeth Pharmaceuticals Inc
A subsidiary of Pfizer Inc
Philadelphia, PA 19101

MADE IN SWITZERLAND



Rapamune[®] (sirolimus) Tablets

1 mg

Unit Dose - 100 Tablets
(10 Blister Cards of 10 Tablets Each)

00228621

FPO 128: 12982200



DOSAGE AND USE: See accompanying prescribing information.

Store at room temperature, 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.] Dispense in tight (USP), light-resistant (USP), and child-resistant containers.

Distributed by
Wyeth Pharmaceuticals Inc
A subsidiary of Pfizer Inc
Philadelphia, PA 19101
MADE IN SWITZERLAND

ALWAYS DISPENSE WITH MEDICATION GUIDE



NDC 0008-1041-05

Rapamune®
(sirolimus) Tablets

1 mg

For oral use only.

100 Tablets

Rx only



12112401

DOSAGE AND USE: See accompanying prescribing information.

Store at room temperature, 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.] Dispense in tight (USP), light-resistant (USP), and child-resistant containers.

Distributed by
Wyeth Pharmaceuticals Inc
A subsidiary of Pfizer Inc
Philadelphia, PA 19101
MADE IN SWITZERLAND

ALWAYS DISPENSE WITH MEDICATION GUIDE



NDC 0008-1042-05

Rapamune®
(sirolimus) Tablets

2 mg

For oral use only.

100 Tablets

Rx only



FP0128 (80% x 5.5mm)
N3 0008-1042-05 1

12112601

FPO - 128 10 mil
13123001



IMPRINT
AREA

LOT
EXP
MFG



Rapamune®
(sirolimus)
Oral Solution

1 mg/mL

60 mL bottle

ALWAYS DISPENSE WITH MEDICATION GUIDE



NDC 0008-1030-04

Rapamune®
(sirolimus)
Oral Solution

1 mg/mL

60 mL bottle

Rx only



Rapamune®
(sirolimus)
Oral Solution

1 mg/mL

60 mL bottle

DOSAGE AND USE:

See accompanying prescribing information.

Keep out of reach of children.

Store refrigerated at 2° to 8°C
(36° to 46°F).

PROTECT FROM LIGHT.

Use contents of bottle within
1 month after opening.

MADE IN JAPAN



N 3 0008-1030-04 1

Distributed by
Wyeth Pharmaceuticals Inc
A subsidiary of Pfizer Inc
Philadelphia, PA 19101

13123001



FPO - 128 10 mil
13123001

ALWAYS DISPENSE WITH MEDICATION GUIDE

NDC 0008-1030-04



Rapamune®
(sirolimus)
Oral Solution

1 mg/mL

For oral use only.

Each mL of RAPAMUNE contains 1 mg sirolimus,
1.5%-2.5% ethanol

Also contains a mixture of propylene glycol and
phosphatidylcholine derived from soy lecithin
(and other components) and polysorbate 80.
60 mL

Rx only

DOSAGE AND USE: See accompanying prescribing information.
Keep out of reach of children.


Distributed by
Wyeth Pharmaceuticals Inc
A subsidiary of Pfizer Inc
Philadelphia, PA 19101

MADE IN JAPAN



(01) 003000 8 103004 1

FPO



ZZZ
LOT
EXP

YYT22901

Store refrigerated at 2° to 8°C (36° to 46°F). Protect from light.
Use contents within 1 month after opening.



FPO: 128
12965902



Rapamune®
(sirolimus)
Oral Solution
1 mg/mL

12965902

Instructions for Use for RAPAMUNE® (sirolimus) Oral Solution




1 Open the solution bottle. Remove the safety cap by squeezing the tabs on the cap and twisting counterclockwise.



2 The first time you use a bottle of Rapamune Oral Solution, insert the oral syringe adapter (plastic tube with stopper) tightly into the bottle until it is even with the top of the bottle. Do not remove the oral syringe adapter from the bottle once inserted.



3 Use a new disposable amber oral syringe for each dose of Rapamune Oral Solution. Fully push down (depress) on the plunger of the disposable amber oral syringe. Then, tightly insert the oral syringe into the opening in the adapter.




4 Withdraw the prescribed amount of Rapamune Oral Solution. Gently pull back the plunger of the syringe until the bottom of the black line of the plunger is even with the appropriate mark on the syringe. Always keep the bottle in an upright position. If bubbles form in the syringe, empty the syringe into the bottle and repeat steps 3 and 4.



5 If your doctor tells you to carry your medicine with you: If you need to carry your Rapamune Oral Solution in a filled syringe, place a cap securely on the syringe. The cap should snap into place.



6 Place the capped syringe in the enclosed carrying case.



7 Taking a dose of Rapamune Oral Solution: Empty the syringe into a glass or plastic cup containing at least 2 ounces (1/4 cup; 60 mL) of water or orange juice, stir vigorously for 1 minute and drink right away. Refill the container with at least 4 ounces (1/2 cup; 120 mL) of water or orange juice, stir vigorously again and drink the rinse solution. Do not mix Rapamune Oral Solution with apple juice, grapefruit juice, or other liquids. Only glass or plastic cups should be used to mix Rapamune Oral Solution. The syringe and cap should be used only one time and then thrown away.

8 Always store the bottles of medication in the refrigerator. You may store RAPAMUNE Oral Solution that is in a syringe at room temperature up to 77°F (25°C) or in the refrigerator at 36°F to 46°F (2°C to 8°C) for up to 24 hours. See "How should I store RAPAMUNE?" at the end of Instructions for Use provided in the package insert.

60 mL Patient Kit

Rapamune®
(sirolimus)
Oral Solution

1 mg/mL

DOSAGE AND USE:
See accompanying prescribing information.

Refer to back of box for Instructions for Use for RAPAMUNE (sirolimus) Oral Solution

Keep out of reach of children.

ALWAYS DISPENSE WITH MEDICATION GUIDE



NDC 0008-1030-06

Rapamune®
(sirolimus)
Oral Solution

1 mg/mL

For oral use only.

Each mL of RAPAMUNE contains 1 mg sirolimus.
1.5%-2.5% ethanol

Also contains a mixture of propylene glycol and phosphatidylcholine derived from soy lecithin (and other components) and polysorbate 80.

60 mL Patient Kit

Rx only


Each Rapamune Oral Solution carton contains:

- a) A 2 oz. (60 mL fill) amber glass bottle of sirolimus (concentration of 1 mg/mL)
- b) 1 oral syringe adapter for fitting into the neck of the bottle
- c) 30 amber oral syringes and 30 caps for daily dosing
- d) 1 carrying case

Also contains: Package Insert and Medication Guide

You will also need:

- glass or plastic cup
- 6 oz. of water or orange juice only



60 mL Patient Kit

Rapamune®
(sirolimus)
Oral Solution

1 mg/mL

Store refrigerated at
2° to 8°C (36° to 46°F).

PROTECT FROM LIGHT

Use contents of bottle
within 1 month after
opening.

MADE IN JAPAN



N 3 0008-1030-06 5

LOT
EXP.

Distributed by
Wyeth Pharmaceuticals Inc
A subsidiary of Pfizer Inc
Philadelphia, PA 19101

