

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use FAZACLO safely and effectively. See full prescribing information for FAZACLO.

FAZACLO® (clozapine) orally disintegrating tablets

Initial U.S. Approval: 1989

**WARNING: AGRANULOCYTOSIS; ORTHOSTATIC HYPOTENSION, BRADYCARDIA, AND SYNCOPES; SEIZURE; MYOCARDITIS AND CARDIOMYOPATHY; INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS**

*See full prescribing information for complete boxed warning.*

- **Agranulocytosis:** Can lead to serious infection and death. Monitor white blood cell count and absolute neutrophil count prior to and during treatment. Monitor for symptoms of agranulocytosis and infection (2.1, 5.1).
- Because of risk of agranulocytosis, FAZACLO is available only through a restricted program called the FAZACLO Patient Registry. Prescribers, patients, and pharmacies must enroll in the program (5.2).
- **Orthostatic Hypotension, Bradycardia, and Syncope:** Risk is dose-related. Starting dose is 12.5 mg. Titrate gradually and use divided dosages (2.2, 2.6, 5.3).
- **Seizure:** Risk is dose-related. Titrate gradually and use divided doses. Use with caution in patients with history of seizure or risk factors for seizure (2.2, 5.4).
- **Myocarditis and Cardiomyopathy:** Can be fatal. Discontinue and obtain cardiac evaluation if findings suggest these cardiac reactions (5.5).
- **Increased Mortality in Elderly Patients with Dementia-Related Psychosis:** FAZACLO is not approved for this condition (5.6).

**RECENT MAJOR CHANGES**

Required Laboratory Testing Prior to Initiation and During Therapy (2.1)	7/2013
Dosage Adjustments with Concomitant Use of CYP1A2, CYP2D6, CYP3A4 Inhibitors or CYP1A2, CYP3A4 Inducers (2.7)	7/2013
Renal or Hepatic Impairment or CYP2D6 Poor Metabolizers (2.8)	7/2013
Contraindications (4.0)	7/2013
FAZACLO Patient Registry Because of the Risk of Agranulocytosis (5.2)	7/2013
Recurrence of Psychosis and Cholinergic Rebound after Abrupt Discontinuation of FAZACLO (5.18)	7/2013

**INDICATIONS AND USAGE**

FAZACLO is an atypical antipsychotic indicated for:

- Treatment-resistant schizophrenia. Efficacy was established in an active-controlled study (1.1, 14.1).
- Reducing suicidal behavior in patients with schizophrenia or schizoaffective disorder. Efficacy was established in an active-controlled study (1.2, 14.2).

**DOSAGE AND ADMINISTRATION**

- Starting Dose: 12.5 mg once daily or twice daily (2.3).
- Use cautious titration and divided dosage schedule (2.3, 5.3).
- Titration: increase the total daily dosage in increments of 25 mg to 50 mg per day, if well-tolerated (2.3).
- Target dose: 300 mg to 450 mg per day, in divided doses, by the end of 2 weeks (2.3).
- Subsequent increases: increase in increments of 100 mg or less, once or twice weekly (2.3).
- Maximum daily dose: 900 mg (2.3). Tablets rapidly disintegrate after placement in the mouth and may be chewed if desired. No water is needed. Can be administered with or without food (2.3).

**DOSAGE FORMS AND STRENGTHS**

Orally disintegrating tablets: 12.5 mg, 25 mg, 100 mg, 150 mg, and 200 mg (3).

**CONTRAINDICATIONS**

- History of clozapine-induced agranulocytosis or severe granulocytopenia (4.1).
- Known hypersensitivity to clozapine or any other component of FAZACLO (4.2).

**WARNINGS AND PRECAUTIONS**

- **Eosinophilia:** Assess for organ involvement (e.g., myocarditis, pancreatitis, hepatitis, colitis, nephritis). Discontinue if these occur (5.7).
- **QT Interval Prolongation:** Can be fatal. Consider additional risk factors for prolonged QT interval (disorders and drugs) (5.8).
- **Metabolic Changes:** Atypical antipsychotic drugs have been associated with metabolic changes that may increase cardiovascular/ cerebrovascular risk. These metabolic changes include:

- **Hyperglycemia and Diabetes Mellitus:** Monitor for symptoms of hyperglycemia including polydipsia, polyuria, polyphagia, and weakness. Monitor glucose regularly in patients with diabetes or at risk for diabetes (5.9).
- **Dyslipidemia:** Undesirable alterations in lipids have occurred in patients treated with atypical antipsychotics (5.9).
- **Weight Gain:** Significant weight gain has occurred. Monitor weight gain (5.9).

- **Neuroleptic Malignant Syndrome (NMS):** Immediately discontinue and monitor closely. Assess for co-morbid conditions (5.10).
- **Fever:** Evaluate for infection, agranulocytosis, NMS (5.11).
- **Pulmonary Embolism (PE):** Consider PE if respiratory distress, chest, pain, or deep vein thrombosis occur (5.12).
- **Anticholinergic Toxicity:** Use cautiously in presence of specific conditions (e.g., narrow angle glaucoma, use of anticholinergic drugs) (5.13).
- **Interference with Cognitive and Motor Performance:** Advise caution when operating machinery, including automobiles (5.14).

**ADVERSE REACTIONS**

Most common adverse reactions ( $\geq 5\%$ ) were: CNS reactions (sedation, dizziness/vertigo, headache, and tremor); cardiovascular reactions (tachycardia, hypotension, and syncope); autonomic nervous system reactions (hypersalivation, sweating, dry mouth, and visual disturbances); gastrointestinal reactions (constipation and nausea); and fever (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Jazz Pharmaceuticals, Inc., at 1-877-329-2256 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)

**DRUG INTERACTIONS**

- Concomitant use of **Strong CYP1A2 Inhibitors:** Reduce FAZACLO dose to one third when coadministered with strong CYP1A2 inhibitors (e.g., fluvoxamine, ciprofloxacin, enoxacin) (2.7, 7.1).
- Concomitant use of **Strong CYP3A4 Inducers:** is not recommended (2.7, 7.1).
- **Discontinuation of CYP1A2 or CYP3A4 Inducers:** Consider reducing FAZACLO dose when CYP1A2 (e.g., tobacco smoke) or CYP3A4 inducers (e.g., carbamazepine) are discontinued (2.7, 7.1).

**USE IN SPECIFIC POPULATIONS**

- **Nursing Mothers:** Discontinue drug or discontinue nursing, taking into consideration importance of drug to mother (8.3).

See 17 for PATIENT COUNSELING INFORMATION.

Revised: July 2013

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**WARNING: AGRANULOCYTOSIS; ORTHOSTATIC HYPOTENSION, BRADYCARDIA, AND SYNCOPES; SEIZURE; MYOCARDITIS AND CARDIOMYOPATHY; INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS**

**1 INDICATIONS AND USAGE**

- 1.1 Treatment Resistant Schizophrenia
- 1.2 Reduction in the Risk of Recurrent Suicidal Behavior in Schizophrenia or Schizoaffective Disorders

**2 DOSAGE AND ADMINISTRATION**

- 2.1 Required Laboratory Testing Prior to Initiation and During Therapy
- 2.2 Important Administration Instructions
- 2.3 Dosing Information
- 2.4 Maintenance Treatment
- 2.5 Discontinuation of Treatment
- 2.6 Re-Initiation of Treatment
- 2.7 Dosage Adjustment with Concomitant Use of CYP1A2, CYP2D6, CYP3A4 Inhibitors or CYP1A2, CYP3A4 Inducers
- 2.8 Renal or Hepatic Impairment, or CYP2D6 Poor Metabolizers

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

- 4.1 History of Clozapine-induced Agranulocytosis or Severe Granulocytopenia
- 4.2 Hypersensitivity

**5 WARNINGS AND PRECAUTIONS**

- 5.1 Agranulocytosis
- 5.2 FAZACLO Patient Registry Because of the Risk of Agranulocytosis
- 5.3 Orthostatic Hypotension, Bradycardia, and Syncope
- 5.4 Seizures
- 5.5 Myocarditis and Cardiomyopathy
- 5.6 Increased Mortality in Elderly Patients with Dementia-Related Psychosis
- 5.7 Eosinophilia
- 5.8 QT Interval Prolongation
- 5.9 Metabolic Changes
- 5.10 Neuroleptic Malignant Syndrome
- 5.11 Fever
- 5.12 Pulmonary Embolism
- 5.13 Anticholinergic Toxicity
- 5.14 Interference with Cognitive and Motor Performance
- 5.15 Tardive Dyskinesia
- 5.16 Patients with Phenylketonuria
- 5.17 Cerebrovascular Adverse Reactions

5.18 Recurrence of Psychosis and Cholinergic Rebound after Abrupt Discontinuation of FAZACLO

**6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

**7 DRUG INTERACTIONS**

- 7.1 Potential for Other Drugs to Affect FAZACLO
  - 7.2 Potential for FAZACLO to Affect Other Drugs
- 8 USE IN SPECIFIC POPULATIONS**
- 8.1 Pregnancy
  - 8.3 Nursing Mothers
  - 8.4 Pediatric Use
  - 8.5 Geriatric Use
  - 8.6 Patients with Renal or Hepatic Impairment
  - 8.7 CYP2D6 Poor Metabolizers

**10 OVERDOSAGE**

- 10.1 Overdosage Experience
- 10.2 Management of Overdosage

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**14 CLINICAL STUDIES**

- 14.1 Treatment Resistant Schizophrenia
- 14.2 Recurrent Suicidal Behavior in Schizophrenia or Schizoaffective Disorder

**16 HOW SUPPLIED/STORAGE AND HANDLING**

- 16.1 How Supplied
- 16.2 Storage and Handling

**17 PATIENT COUNSELING INFORMATION**

\*Sections or subsections omitted from the Full Prescribing Information are not listed.

## FULL PRESCRIBING INFORMATION

### **WARNING: AGRANULOCYTOSIS; ORTHOSTATIC HYPOTENSION, BRADYCARDIA, AND SYNCOPE; SEIZURE; MYOCARDITIS AND CARDIOMYOPATHY; INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS**

#### **Agranulocytosis**

Clozapine treatment has caused agranulocytosis, defined as an absolute neutrophil count (ANC) less than 500/mm<sup>3</sup>. Agranulocytosis can lead to serious infection and death. Prior to initiating treatment with FAZACLO, obtain a baseline white blood cell count (WBC) and ANC. The ANC must be greater than or equal to 2000/mm<sup>3</sup> and the WBC must be greater than or equal to 3500/mm<sup>3</sup> for a patient to begin treatment with FAZACLO. During treatment, patients must have regular monitoring of ANC and WBC. Discontinue FAZACLO and do not rechallenge if the ANC is less than 1000/mm<sup>3</sup> or the WBC is less than 2000/mm<sup>3</sup>. Advise patients to immediately report symptoms consistent with agranulocytosis or infection (e.g., fever, weakness, lethargy, or sore throat) [see *Dosage and Administration* (2.1) and *Warnings and Precautions* (5.1)].

Because of the risk of agranulocytosis, FAZACLO is available only through a restricted program called the FAZACLO Patient Registry. Under the FAZACLO Patient Registry, prescribers, patients, and pharmacies must enroll in the program [see *Warnings and Precautions* (5.2)].

#### **Orthostatic Hypotension, Bradycardia, Syncope**

Orthostatic hypotension, bradycardia, syncope, and cardiac arrest have occurred with clozapine treatment. The risk is highest during the initial titration period, particularly with rapid dose escalation. These reactions can occur with the first dose, with doses as low as 12.5 mg per day. Initiate treatment at 12.5 mg once or twice daily; titrate slowly; and use divided dosages. Use FAZACLO cautiously in patients with cardiovascular/cerebrovascular disease or conditions predisposing to hypotension (e.g., dehydration, use of antihypertensive medications) [see *Dosage and Administration* (2.3, 2.6) and *Warnings and Precautions* (5.3)].

#### **Seizures**

Seizures have occurred with clozapine treatment. The risk is dose-related. Initiate treatment at 12.5 mg, titrate gradually, and use divided dosing. Use caution when administering FAZACLO to patients with a history of seizures or other predisposing risk factors for seizure (CNS pathology, medications that lower the seizure threshold, alcohol abuse). Caution patients about engaging in any activity where sudden loss of consciousness could cause serious risk to themselves or others [see *Dosage and Administration* (2.3), *Warnings and Precautions* (5.4)].

#### **Myocarditis and Cardiomyopathy**

Fatal myocarditis and cardiomyopathy have occurred with clozapine treatment. Discontinue FAZACLO and obtain a cardiac evaluation upon suspicion of these reactions. Generally, patients with FAZACLO-related myocarditis or cardiomyopathy should not be

**rechallenged with FAZACLO. Consider the possibility of myocarditis or cardiomyopathy if chest pain, tachycardia, palpitations, dyspnea, fever, flu-like symptoms, hypotension, or ECG changes occur [see Warnings and Precautions (5.5)].**

**Increased Mortality in Elderly Patients with Dementia-Related Psychosis**

**Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. FAZACLO is not approved for use in patients with dementia-related psychosis [see Warnings and Precautions (5.6)].**

## **1 INDICATIONS AND USAGE**

### **1.1 Treatment Resistant Schizophrenia**

FAZACLO is indicated for the treatment of severely ill patients with schizophrenia who fail to respond adequately to standard antipsychotic treatment. Because of the significant risk of agranulocytosis and seizure associated with its use, FAZACLO should be used only in patients who have failed to respond adequately to standard antipsychotic treatment [see Warnings and Precautions (5.1, 5.4)].

The effectiveness of clozapine in treatment-resistant schizophrenia was demonstrated in a 6-week, randomized, double-blind, active-controlled study comparing clozapine and chlorpromazine in patients who had failed other antipsychotics [see Clinical Studies (14.1)].

### **1.2 Reduction in the Risk of Recurrent Suicidal Behavior in Schizophrenia or Schizoaffective Disorder**

FAZACLO is indicated for reducing the risk of recurrent suicidal behavior in patients with schizophrenia or schizoaffective disorder who are judged to be at chronic risk for re-experiencing suicidal behavior, based on history and recent clinical state. Suicidal behavior refers to actions by a patient that put him/herself at risk for death.

The effectiveness of clozapine in reducing the risk of recurrent suicidal behavior was demonstrated over a two-year treatment period in the InterSePT™ trial [see Clinical Studies (14.2)].

## **2 DOSAGE AND ADMINISTRATION**

### **2.1 Required Laboratory Testing Prior to Initiation and During Therapy**

Prior to initiating treatment with FAZACLO, obtain a complete blood count (CBC) with differential. The absolute neutrophil count (ANC) must be greater than or equal to 2000/mm<sup>3</sup> and the WBC must be greater than or equal to 3500/mm<sup>3</sup> in order to initiate treatment. To continue treatment, the ANC and WBC must be monitored regularly [see Warnings and Precautions (5.1)].

## **2.2 Important Administration Instructions**

FAZACLO orally disintegrating tablets should be immediately placed in the mouth after removing the tablet from the blister pack or bottle. The tablet disintegrates rapidly after placement in the mouth. The tablets can be allowed to disintegrate, or they may be chewed. They may be swallowed with saliva. No water is necessary for administration.

The orally disintegrating tablets in a blister pack should be left in the unopened blister until the time of use. Just prior to use, peel the foil from the blister and gently remove the orally disintegrating tablet. Do not push the tablets through the foil, because this could damage the tablet.

## **2.3 Dosing Information**

The starting dose is 12.5 mg once daily or twice daily. The total daily dose can be increased in increments of 25 mg to 50 mg per day, if well-tolerated, to achieve a target dose of 300 mg to 450 mg per day (administered in divided doses) by the end of 2 weeks. Subsequently, the dose can be increased once weekly or twice weekly, in increments of up to 100 mg. The maximum dose is 900 mg per day. To minimize the risk of orthostatic hypotension, bradycardia, and syncope, it is necessary to use this low starting dose, gradual titration schedule, and divided dosages [*see Warnings and Precautions (5.3)*].

FAZACLO can be taken with or without food [*see Pharmacokinetics (12.3)*].

## **2.4 Maintenance Treatment**

Generally, it is recommended that patients responding to FAZACLO continue maintenance treatment on their effective dose beyond the acute episode.

## **2.5 Discontinuation of Treatment**

In the event of planned termination of FAZACLO therapy, reduce the dose gradually over a period of 1 to 2 weeks. If abrupt discontinuation is necessary (because of agranulocytosis or another medical condition, for example), monitor carefully for the recurrence of psychotic symptoms and symptoms related to cholinergic rebound such as profuse sweating, headache, nausea, vomiting, and diarrhea.

## **2.6 Re-Initiation of Treatment**

When restarting FAZACLO in patients who have discontinued FAZACLO (i.e., 2 days or more since the last dose), re-initiate with 12.5 mg once daily or twice daily. This is necessary to minimize the risk of hypotension, bradycardia, and syncope [*see Warnings and Precautions (5.3)*]. If that dose is well tolerated, the dose may be increased to the previously therapeutic dose more quickly than recommended for initial treatment.

## 2.7 Dosage Adjustments with Concomitant Use of CYP1A2, CYP2D6, CYP3A4 Inhibitors or CYP1A2, CYP3A4 Inducers

Dose adjustments may be necessary in patients with concomitant use of: strong CYP1A2 inhibitors (e.g., fluvoxamine, ciprofloxacin, or enoxacin); moderate or weak CYP1A2 inhibitors (e.g., oral contraceptives, or caffeine); CYP2D6 or CYP3A4 inhibitors (e.g., cimetidine, escitalopram, erythromycin, paroxetine, bupropion, fluoxetine, quinidine, duloxetine, terbinafine, or sertraline); CYP3A4 inducers (e.g., phenytoin, carbamazepine, St. John's wort, and rifampin); or CYP1A2 inducers (e.g., tobacco smoking) (Table 1) [see *Drug Interactions* (7)].

**Table 1: Dosage Adjustments in Patients Taking Concomitant Medications**

Co-Medications	Scenarios		
	Initiating FAZACLO while taking a co-medication	Adding a co-medication while taking FAZACLO	Discontinuing a co-medication while continuing FAZACLO
Strong CYP1A2 Inhibitors	Use one third of the FAZACLO dose.		Increase FAZACLO dose based on clinical response.
Moderate or Weak CYP1A2 Inhibitors	Monitor for adverse reactions. Consider reducing the FAZACLO dose if necessary.		Monitor for lack of effectiveness. Consider increasing FAZACLO dose if necessary.
CYP2D6 or CYP3A4 Inhibitors			
Strong CYP3A4 Inducers	Concomitant use is not recommended. However, if the inducer is necessary, it may be necessary to increase the FAZACLO dose. Monitor for decreased effectiveness.		Reduce FAZACLO dose based on clinical response.
Moderate or Weak CYP1A2 or CYP3A4 Inducers	Monitor for decreased effectiveness. Consider increasing the FAZACLO dose if necessary.		Monitor for adverse reactions. Consider reducing the FAZACLO dose if necessary.

## 2.8 Renal or Hepatic Impairment or CYP2D6 Poor Metabolizers

It may be necessary to reduce the FAZACLO dose in patients with significant renal or hepatic impairment, or in CYP2D6 poor metabolizers [see *Use in Specific Populations* (8.6, 8.7)].

## 3 DOSAGE FORMS AND STRENGTHS

FAZACLO is available as 12.5 mg, 25 mg, 100 mg, 150 mg, and 200 mg round, yellow, orally disintegrating tablets.

## 4 CONTRAINDICATIONS

### 4.1 History of Clozapine-induced Agranulocytosis or Severe Granulocytopenia

FAZACLO is contraindicated in patients with a history of clozapine-induced agranulocytosis or severe granulocytopenia [see *Warnings and Precautions* (5.1)].

### 4.2 Hypersensitivity

FAZACLO is contraindicated in patients with a history of hypersensitivity to clozapine (e.g., photosensitivity, vasculitis, erythema multiforme, or Stevens-Johnson Syndrome) or any other component of FAZACLO [see *Adverse Reactions* (6.2)].

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Agranulocytosis

#### Background

Agranulocytosis, defined as an ANC of less than  $500/\text{mm}^3$ , has been estimated to occur in association with clozapine use at a cumulative incidence at 1 year of approximately 1.3%, based on the occurrence of 15 US cases out of 1743 patients exposed to clozapine during its clinical testing prior to domestic marketing. All of these cases occurred at a time when the need for close monitoring of WBC counts was already recognized. A hematologic risk analysis was conducted based upon the available information in the *Clozapine National Nonrechallenge Masterfile* for US patients. Based upon a cut-off date of April 30, 1995, the incidence rates of agranulocytosis based upon a weekly monitoring schedule rose steeply during the first two months of therapy, peaking in the third month. Among clozapine patients who continued the drug beyond the third month, the weekly incidence of agranulocytosis fell a substantial degree. After 6 months, the weekly incidence of agranulocytosis declines still further; however, it never reaches zero. It should be noted that any type of reduction in the frequency of monitoring WBC counts may result in an increased incidence of agranulocytosis.

#### Risk Factors

Experience from clinical development, as well as from examples in the medical literature, suggests that patients who have developed agranulocytosis during clozapine therapy are at increased risk of subsequent episodes of agranulocytosis. Analysis of WBC count data from the *Clozapine National Nonrechallenge Masterfile* also suggests that patients who have an initial episode of moderate leukopenia ( $3000/\text{mm}^3 \geq \text{WBC count} \geq 2000/\text{mm}^3$ ) are at an increased risk of subsequent episodes of agranulocytosis. Except for bone-marrow suppression during initial clozapine therapy, there are no other established risk factors based on worldwide experience for the development of agranulocytosis in association with clozapine use. However, a disproportionate number of the US cases of agranulocytosis occurred in patients of Jewish background compared to the overall proportion of such patients exposed during domestic development of clozapine. Most of the US cases of agranulocytosis occurred within 4–10 weeks of exposure, but neither dose nor duration is a reliable predictor of this problem. Agranulocytosis

associated with other antipsychotic drugs has been reported to occur with a greater frequency in women, the elderly, and in patients who are cachectic or have serious underlying medical illness; such patients may also be at particular risk with FAZACLO, although this has not been definitively demonstrated.

**WBC Count and ANC Clinical Monitoring Schedule**

FAZACLO is available only through a distribution system that ensures monitoring of WBC count and ANC according to the schedule described below prior to delivery of the next supply of medication.

As described in Table 2, patients who are being treated with FAZACLO must have a baseline WBC count and ANC before initiation of treatment, and a WBC count and ANC every week for the first 6 months. Thereafter, if acceptable WBC counts and ANCs (WBC count  $\geq 3500/\text{mm}^3$  and ANC  $\geq 2000/\text{mm}^3$ ) have been maintained during the first 6 months of continuous therapy, WBC counts and ANCs can be monitored every 2 weeks for the next 6 months. Thereafter, if acceptable WBC counts and ANCs (WBC count  $\geq 3500/\text{mm}^3$  and ANC  $\geq 2000/\text{mm}^3$ ) have been maintained during the second 6 months of continuous therapy, WBC count and ANC can be monitored every 4 weeks.

When treatment with FAZACLO is discontinued (regardless of the reason), WBC count and ANC must be monitored weekly for at least 4 weeks from the day of discontinuation or until WBC count  $\geq 3500/\text{mm}^3$  and ANC  $\geq 2000/\text{mm}^3$ .

Table 2 provides a summary of the frequency of monitoring that should occur based on various stages of therapy (e.g., initiation of therapy) or results from WBC count and ANC monitoring tests (e.g., moderate leukopenia). The text that follows should be consulted for additional details regarding the treatment of patients under the various conditions (e.g., severe leukopenia).

Advise patients to immediately report the appearance of signs/symptoms consistent with agranulocytosis or infection (e.g., fever, weakness, lethargy, or sore throat) at any time during FAZACLO therapy. Such patients should have a WBC count and an ANC performed promptly.

**Table 2. Frequency of Monitoring Based on Stage of Therapy or Results from WBC Count and ANC Monitoring Tests**

Situation	Hematological Values for Monitoring	Frequency of WBC Count and ANC Monitoring
Initiation of therapy	WBC count $\geq 3500/\text{mm}^3$ and ANC $\geq 2000/\text{mm}^3$  Note: Do not initiate in patients with a history of clozapine-induced agranulocytosis or severe granulocytopenia.	Weekly for 6 months

<b>Situation</b>	<b>Hematological Values for Monitoring</b>	<b>Frequency of WBC Count and ANC Monitoring</b>
6 to 12 months of therapy	WBC $\geq$ 3500/mm <sup>3</sup> and ANC $\geq$ 2000/mm <sup>3</sup>	Every 2 weeks for 6 months
12 months of therapy	WBC $\geq$ 3500/mm <sup>3</sup> and ANC $\geq$ 2000/mm <sup>3</sup>	Every 4 weeks ad infinitum
Immature forms present	N/A	Repeat WBC and ANC
Discontinuation of therapy	N/A	Weekly for at least 4 weeks from day of discontinuation or until WBC $\geq$ 3500/mm <sup>3</sup> and ANC $\geq$ 2000/mm <sup>3</sup>
Substantial drop in WBC or ANC	Single drop or cumulative drop within 3 weeks of: WBC $\geq$ 3000/mm <sup>3</sup> or ANC $\geq$ 1500/mm <sup>3</sup>	<ol style="list-style-type: none"> <li>1. Repeat WBC and ANC</li> <li>2. If repeat values are: WBC 3000/mm<sup>3</sup> to 3500 and ANC &gt; 2000/mm<sup>3</sup>, then monitor twice weekly</li> </ol>
Mild leukopenia and/or Mild granulocytopenia	If WBC 3000/mm <sup>3</sup> to < 3500/mm <sup>3</sup> and/or ANC 1500/mm <sup>3</sup> to < 2000/mm <sup>3</sup>	Twice weekly until WBC > 3500/mm <sup>3</sup> and ANC > 2000/mm <sup>3</sup> then return to previous monitoring frequency
Moderate leukopenia and/or Moderate granulocytopenia	WBC 2000/mm <sup>3</sup> to < 3000/mm <sup>3</sup> and/or ANC 1000/mm <sup>3</sup> to < 1500/mm <sup>3</sup>	<ol style="list-style-type: none"> <li>1. Interrupt therapy</li> <li>2. Daily until WBC &gt; 3000/mm<sup>3</sup> and ANC &gt; 1500/mm<sup>3</sup></li> <li>3. Twice weekly until WBC &gt; 3500/mm<sup>3</sup> and ANC &gt; 2000/mm<sup>3</sup></li> <li>4. May rechallenge when WBC &gt; 3500/mm<sup>3</sup> and ANC &gt; 2000/mm<sup>3</sup></li> <li>5. If rechallenged, monitor weekly for 1 year before returning to the usual monitoring schedule of every 2 weeks for 6 months and then every 4 weeks ad infinitum</li> </ol>
Severe leukopenia and/or Severe granulocytopenia	WBC count < 2000/mm <sup>3</sup> and/or ANC < 1000/mm <sup>3</sup>	<ol style="list-style-type: none"> <li>1. Discontinue treatment and do not rechallenge patient</li> <li>2. Monitor until normal and for at least four weeks from day of discontinuation as follows: <ul style="list-style-type: none"> <li>- Daily until WBC &gt; 3000/mm<sup>3</sup> and ANC &gt; 1500/mm<sup>3</sup></li> <li>- Twice weekly until WBC &gt; 3500/mm<sup>3</sup> and ANC &gt; 2000/mm<sup>3</sup></li> <li>- Weekly after WBC &gt; 3500/mm<sup>3</sup></li> </ul> </li> </ol>
Agranulocytosis	ANC < 500/mm <sup>3</sup>	<ol style="list-style-type: none"> <li>1. Discontinue treatment and do not rechallenge patient</li> <li>2. Monitor until normal and for at least four weeks</li> </ol>

Situation	Hematological Values for Monitoring	Frequency of WBC Count and ANC Monitoring
		<p>from day of discontinuation as follows:</p> <ul style="list-style-type: none"><li>– Daily until <math>\text{WBC} &gt; 3000/\text{mm}^3</math> and <math>\text{ANC} &gt; 1500/\text{mm}^3</math></li><li>– Twice weekly until <math>\text{WBC} &gt; 3500/\text{mm}^3</math> and <math>\text{ANC} &gt; 2000/\text{mm}^3</math></li><li>– Weekly after <math>\text{WBC} &gt; 3500/\text{mm}^3</math></li></ul>

WBC = White blood cell

ANC = Absolute neutrophil count

#### Decrements in WBC Count and/or ANC

Consult Table 2 above to determine how to monitor patients who experience decrements in WBC count and/or ANC at any point during treatment. Additionally, patients should be carefully monitored for flu-like symptoms or other symptoms suggestive of infection.

#### Nonrechallengeable Patients

If the total WBC count falls below  $2000/\text{mm}^3$  or the ANC falls below  $1000/\text{mm}^3$ , bone-marrow aspiration should be considered to ascertain granulopoietic status and patients should not be rechallenged with FAZACLO. Protective isolation with close observation may be indicated if granulopoiesis is determined to be deficient. Should evidence of infection develop, the patient should have appropriate cultures performed and an appropriate antibiotic regimen instituted.

Patients discontinued from clozapine therapy due to significant granulopoietic suppression have been found to develop agranulocytosis upon rechallenge, often with a shorter latency on re-exposure. To reduce the chances of rechallenge occurring in patients who have experienced significant bone-marrow suppression during FAZACLO therapy, a single, national master file (i.e., Nonrechallengeable Database) is confidentially maintained.

#### Treatment of Rechallengeable Patients

Patients may be rechallenged with FAZACLO if their WBC count does not fall below  $2000/\text{mm}^3$  and the ANC does not fall below  $1000/\text{mm}^3$ . However, analysis of the data from the *Clozapine National Registry* suggests that patients who have an initial episode of moderate leukopenia ( $3000/\text{mm}^3 > \text{WBC count} \geq 2000/\text{mm}^3$ ) have up to a 12-fold increased risk of having a subsequent episode of agranulocytosis when rechallenged as compared to the full cohort of patients treated with clozapine. Although FAZACLO therapy may be resumed if no symptoms of infection develop and when the WBC count rises above  $3500/\text{mm}^3$  and the ANC rises above  $2000/\text{mm}^3$ , prescribers are strongly advised to consider whether the benefit of continuing FAZACLO treatment outweighs the increased risk of agranulocytosis.


Analyses of the *Clozapine National Registry* have shown an increased risk of having a subsequent episode of granulopoietic suppression up to a year after recovery from the initial episode. Therefore, as noted in Table 2, patients must undergo weekly WBC count and ANC

monitoring for one year following recovery from an episode of moderate leukopenia and/or moderate granulocytopenia regardless of when the episode develops. If acceptable WBC counts and ANC (WBC count  $\geq 3500/\text{mm}^3$  and ANC  $\geq 2000/\text{mm}^3$ ) have been maintained during the year of weekly monitoring, WBC counts can be monitored every 2 weeks for the next 6 months. If acceptable WBC counts and ANC (WBC count  $\geq 3500/\text{mm}^3$  and ANC  $\geq 2000/\text{mm}^3$ ) continue to be maintained during the 6 months of every-2-week monitoring, WBC counts can be monitored every 4 weeks thereafter, ad infinitum.

**Interruptions in Therapy**

Figure 1 provides instructions regarding re-initiating therapy and subsequently the frequency of WBC count and ANC monitoring after a period of interruption.

**Figure 1. Resuming Monitoring Frequency after Interruption of Therapy**



\*Transitions to reduce frequency of monitoring only permitted if all WBC counts  $\geq 3500/\text{mm}^3$  and ANCs  $\geq 2000/\text{mm}^3$ .

## 5.2 FAZACLO Patient Registry Because of the Risk of Agranulocytosis

Because of the risk of agranulocytosis, FAZACLO is available only through a restricted program called the FAZACLO Patient Registry. Under the FAZACLO Patient Registry, prescribers, patients, pharmacies, and distributors must enroll in the program.

Required components of the FAZACLO Patient Registry are:

- Healthcare professionals who prescribe FAZACLO must enroll in the program and comply with the Registry requirements.
- Pharmacies that dispense FAZACLO must enroll in the program and comply with the Registry requirements.
- Routine monitoring and submission of laboratory results (WBC and ANC) is required during treatment with FAZACLO [*see Warnings and Precautions (5.1)*].
- Patients that receive FAZACLO must be enrolled in a registry.

Further information is available at <https://www.fazacloregistry.com> or 1-877-329-2256.

## 5.3 Orthostatic Hypotension, Bradycardia, and Syncope

Hypotension, bradycardia, syncope, and cardiac arrest have occurred with clozapine treatment. The risk is highest during the initial titration period, particularly with rapid dose-escalation. These reactions can occur with the first dose, at doses as low as 12.5 mg. These reactions can be fatal. The syndrome is consistent with neurally mediated reflex bradycardia (NMRB).

Treatment must begin at a maximum dose of 12.5 mg once daily or twice daily. The total daily dose can be increased in increments of 25 mg to 50 mg per day, if well-tolerated, to a target dose of 300 mg to 450 mg per day (administered in divided doses) by the end of 2 weeks. Subsequently, the dose can be increased weekly or twice weekly, in increments of up to 100 mg. The maximum dose is 900 mg per day. Use cautious titration and a divided dosage schedule to minimize the risk of serious cardiovascular reactions [*see Dosage and Administration (2.3)*]. Consider reducing the dose if hypotension occurs. When restarting patients who have had even a brief interval off FAZACLO (i.e., 2 days or more since the last dose), re-initiate treatment at 12.5 mg dose once daily or twice daily [*see Dosage and Administration (2.6)*].

Use FAZACLO cautiously in patients with cardiovascular disease (history of myocardial infarction or ischemia, heart failure, or conduction abnormalities), cerebrovascular disease, and conditions which would predispose patients to hypotension (e.g., concomitant use of antihypertensives, dehydration and hypovolemia).

## 5.4 Seizures

Seizure has been estimated to occur in association with clozapine use at a cumulative incidence at one year of approximately 5%, based on the occurrence of one or more seizures in 61 of 1743 patients exposed to clozapine during its clinical testing prior to domestic marketing (i.e., a

crude rate of 3.5%). The risk of seizure is dose-related. Initiate treatment with a low dose (12.5 mg), titrate slowly, and use divided dosing [*see Dosage and Administration (2.3)*].

Use caution when administering FAZACLO to patients with a history of seizures or other predisposing risk factors for seizure (e.g., head trauma or other CNS pathology, use of medications that lower the seizure threshold or alcohol abuse). Because of the substantial risk of seizure associated with FAZACLO use, caution patients about engaging in any activity where sudden loss of consciousness could cause serious risk to themselves or others (e.g., driving an automobile, operating complex machinery, swimming, climbing).

## **5.5 Myocarditis and Cardiomyopathy**

Myocarditis and cardiomyopathy have occurred with the use of clozapine. These reactions can be fatal. Discontinue FAZACLO and obtain a cardiac evaluation upon suspicion of myocarditis or cardiomyopathy. Generally, patients with a history of clozapine-associated myocarditis or cardiomyopathy should not be rechallenged with FAZACLO. However, if the benefit of FAZACLO treatment is judged to outweigh the potential risks of recurrent myocarditis or cardiomyopathy, the clinician may consider rechallenge with FAZACLO in consultation with a cardiologist, after a complete cardiac evaluation, and under close monitoring.

Consider the possibility of myocarditis or cardiomyopathy in patients receiving FAZACLO who present with chest pain, dyspnea, persistent tachycardia at rest, palpitations, fever, flu-like symptoms, hypotension, other signs or symptoms of heart failure, or electrocardiographic findings (low voltages, ST-T abnormalities, arrhythmias, right axis deviation, and poor R wave progression). Myocarditis most frequently presents within the first two months of FAZACLO treatment. Symptoms of cardiomyopathy generally occur later than clozapine-associated myocarditis and usually after 8 weeks of treatment. However, myocarditis and cardiomyopathy can occur at any period during treatment with FAZACLO. It is common for nonspecific flu-like symptoms such as malaise, myalgia, pleuritic chest pain, and low-grade fevers to precede more overt signs of heart failure. Typical laboratory findings include elevated troponin I or T, elevated creatinine kinase-MB, peripheral eosinophilia, and elevated C-reactive protein (CRP). Chest roentgenogram may demonstrate cardiac silhouette enlargement, and cardiac imaging (echocardiogram, radionucleotide studies, or cardiac catheterization) may reveal evidence of left ventricular dysfunction.

## **5.6 Increased Mortality in Elderly Patients with Dementia-Related Psychosis**

Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Analyses of 17 placebo-controlled trials (modal duration of 10 weeks), largely in patients taking atypical antipsychotic drugs, revealed a risk of death in drug-treated patients of between 1.6 to 1.7 times the risk of death in placebo-treated patients. Over the course of a typical 10-week controlled trial, the rate of death in drug-treated patients was about 4.5%, compared to a rate of about 2.6% in the placebo group. Although the causes of death were varied, most of the deaths appeared to be either cardiovascular (e.g., heart failure, sudden death) or infectious (e.g., pneumonia) in nature. Observational studies suggest that, similar to atypical antipsychotic drugs, treatment with conventional antipsychotic drugs may increase mortality in this population. The extent to which the findings of increased mortality in observational studies

may be attributed to the antipsychotic drug as opposed to some characteristic(s) of the patients is not clear. FAZACLO is not approved for the treatment of patients with dementia-related psychosis [see *Boxed Warning*].

### **5.7 Eosinophilia**

Eosinophilia, defined as a blood eosinophil count of greater than 700/mm<sup>3</sup>, has occurred with clozapine treatment. In clinical trials, approximately 1% of patients developed eosinophilia. Clozapine-related eosinophilia usually occurs during the first month of treatment. In some patients, it has been associated with myocarditis, pancreatitis, hepatitis, colitis, and nephritis. Such organ involvement could be consistent with a drug reaction with eosinophilia and systemic symptoms syndrome (DRESS), also known as drug induced hypersensitivity syndrome (DIHS). If eosinophilia develops during FAZACLO treatment, evaluate promptly for signs and symptoms of systemic reactions, such as rash or other allergic symptoms, myocarditis, or other organ-specific disease associated with eosinophilia. If clozapine-related systemic disease is suspected, discontinue FAZACLO immediately.

If a cause of eosinophilia unrelated to clozapine is identified (e.g., asthma, allergies, collagen vascular disease, parasitic infections, and specific neoplasms), treat the underlying cause and continue FAZACLO.

Clozapine-related eosinophilia has also occurred in the absence of organ involvement and can resolve without intervention. There are reports of successful rechallenge after discontinuation of clozapine, without recurrence of eosinophilia. In the absence of organ involvement, continue FAZACLO under careful monitoring. If the total eosinophil count continues to increase over several weeks in the absence of systemic disease, the decision to interrupt FAZACLO therapy and rechallenge after the eosinophil count decreases should be based on the overall clinical assessment, in consultation with an internist or hematologist.

### **5.8 QT Interval Prolongation**

QT prolongation, Torsades de Pointes and other life-threatening ventricular arrhythmias, cardiac arrest, and sudden death have occurred with clozapine treatment. When prescribing FAZACLO, consider the presence of additional risk factors for QT prolongation and serious cardiovascular reactions. Conditions that increase these risks include the following: history of QT prolongation, long QT syndrome, family history of long QT syndrome or sudden cardiac death, significant cardiac arrhythmia, recent myocardial infarction, uncompensated heart failure, treatment with other medications that cause QT prolongation, treatment with medications that inhibit the metabolism of FAZACLO, and electrolyte abnormalities.

Prior to initiating treatment with FAZACLO, perform a careful physical examination, medical history, and concomitant medication history. Consider obtaining a baseline ECG and serum chemistry panel. Correct electrolyte abnormalities. Discontinue FAZACLO if the QTc interval exceeds 500 msec. If patients experience symptoms consistent with Torsades de Pointes or other arrhythmias (e.g., syncope, presyncope, dizziness, or palpitations), obtain a cardiac evaluation and discontinue FAZACLO.

Use caution when administering concomitant medications that prolong the QT interval or inhibit the metabolism of FAZACLO. Drugs that cause QT prolongation include: specific antipsychotics (e.g., ziprasidone, iloperidone, chlorpromazine, thioridazine, mesoridazine, droperidol, pimozide), specific antibiotics (e.g., erythromycin, gatifloxacin, moxifloxacin, sparfloxacin), Class 1A antiarrhythmic medications (e.g., quinidine, procainamide) or Class III antiarrhythmics (e.g., amiodarone, sotalol), and others (e.g., pentamidine, levomethadyl acetate, methadone, halofantrine, mefloquine, dolasetron mesylate, probucol or tacrolimus). FAZACLO is primarily metabolized by CYP isoenzymes 1A2, 2D6, and 3A4. Concomitant treatment with inhibitors of these enzymes can increase the concentration of FAZACLO [see *Drug Interactions (7.1)* and *Clinical Pharmacology (12.3)*].

Hypokalemia and hypomagnesemia increase the risk of QT prolongation. Hypokalemia can result from diuretic therapy, diarrhea, and other causes. Use caution when treating patients at risk for significant electrolyte disturbance, particularly hypokalemia. Obtain baseline measurements of serum potassium and magnesium levels, and periodically monitor electrolytes. Correct electrolyte abnormalities before initiating treatment with FAZACLO.

## 5.9 Metabolic Changes

Atypical antipsychotic drugs, including clozapine have been associated with metabolic changes that can increase cardiovascular and cerebrovascular risk. These metabolic changes include hyperglycemia, dyslipidemia, and body weight gain. While atypical antipsychotic drugs may produce some metabolic changes, each drug in the class has its own specific risk profile.

### Hyperglycemia and Diabetes Mellitus

Hyperglycemia, in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, has been reported in patients treated with atypical antipsychotics including clozapine. Assessment of the relationship between atypical antipsychotic use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Given these confounders, the relationship between atypical antipsychotic use and hyperglycemia-related adverse reactions is not completely understood. However, epidemiological studies suggest an increased risk of treatment-emergent, hyperglycemia-related adverse reactions in patients treated with the atypical antipsychotics. Precise risk estimates for hyperglycemia-related adverse reactions in patients treated with atypical antipsychotics are not available.

Patients with an established diagnosis of diabetes mellitus who are started on FAZACLO should be monitored regularly for worsening of glucose control. Patients with risk factors for diabetes mellitus (e.g., obesity, family history of diabetes) who are starting treatment with atypical antipsychotics should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics should be monitored for symptoms of hyperglycemia including polydipsia, polyuria, polyphagia, and weakness. Patients who develop symptoms of hyperglycemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some cases, hyperglycemia has

resolved when the atypical antipsychotic was discontinued; however, some patients required continuation of anti-diabetic treatment despite discontinuation of the suspect drug.

In a pooled data analysis of 8 studies in adult subjects with schizophrenia, the mean changes in fasting glucose concentration in the clozapine and chlorpromazine groups were +11 mg/dL and +4 mg/dL respectively. A higher proportion of the clozapine group demonstrated categorical increases from baseline in fasting glucose concentrations, compared to the chlorpromazine group (Table 3). The clozapine doses were 100-900 mg per day (mean modal dose: 512 mg per day). The maximum chlorpromazine dose was 1800 mg per day (mean modal dose: 1029 mg per day). The median duration of exposure was 42 days for clozapine and chlorpromazine.

**Table 3. Categorical Changes in Fasting Glucose Level in Studies in Adult Subjects with Schizophrenia**

Laboratory Parameter	Category Change (at least once) from Baseline	Treatment Arm	N	n (%)
Fasting Glucose	Normal (< 100 mg/dL) to High ( $\geq$ 126 mg/dL)	Clozapine	198	53 (27)
		Chlorpromazine	135	14 (10)
	Borderline (100 to 125 mg/dL) to High ( $\geq$ 126 mg/dL)	Clozapine	57	24 (42)
		Chlorpromazine	43	12 (28)

#### Dyslipidemia

Undesirable alterations in lipids have occurred in patients treated with atypical antipsychotics, including clozapine. Clinical monitoring, including baseline and periodic follow-up lipid evaluations in patients using FAZACLO, is recommended.

In a pooled data analysis of 10 studies in adult subjects with schizophrenia, clozapine treatment was associated with increases in serum total cholesterol. No data were collected on LDL and HDL cholesterol. The mean increase in total cholesterol was 13 mg/dL in the clozapine group and 15 mg/dL in the chlorpromazine group. In a pooled data analysis of 2 studies in adult subjects with schizophrenia, clozapine treatment was associated with increases in fasting serum triglyceride. The mean increase in fasting triglyceride was 71 mg/dL (54%) in the clozapine group and 39 mg/dL (35%) in the chlorpromazine group (Table 4). In addition, clozapine treatment was associated with categorical increases in serum total cholesterol and triglyceride, as illustrated in Table 5. The proportion of patients with categorical increases in total cholesterol or fasting triglyceride increased with the duration of exposure. The median duration of clozapine and chlorpromazine exposure was 45 days and 38 days, respectively. The clozapine dose range was 100 mg to 900 mg daily; the maximum chlorpromazine dose was 1800 mg daily.

**Table 4. Mean Changes in Total Cholesterol and Triglyceride Concentration in Studies in Adult Subjects with Schizophrenia**

Treatment Arm	Baseline Total Cholesterol Concentration (mg/dL)	Change from Baseline mg/dL (%)
Clozapine (N=334)	184	+13 (7)
Chlorpromazine (185)	182	+15 (8)
	Baseline Triglyceride Concentration (mg/dL)	Change from Baseline mg/dL (%)
Clozapine (N=6)	130	+71 (54)
Chlorpromazine (N=7)	110	+39 (35)

**Table 5. Categorical Changes in Lipid Concentrations in Studies in Adult Subjects with Schizophrenia**

Laboratory Parameter	Category Change (at least once) from Baseline	Treatment Arm	N	n (%)
Total Cholesterol (random or fasting)	Increase by $\geq 40$ mg/dL	Clozapine	334	111 (33 )
		Chlorpromazine	185	46 (25)
	Normal (< 200 mg/dL) to High ( $\geq 240$ mg/dL)	Clozapine	222	18 (8)
		Chlorpromazine	132	3 (2)
	Borderline (200 - 239 mg/dL) to High ( $\geq 240$ mg/dL)	Clozapine	79	30 (38)
		Chlorpromazine	34	14 (41)
Triglycerides (fasting)	Increase by $\geq 50$ mg/dL	Clozapine	6	3 (50)
		Chlorpromazine	7	3(43)
	Normal (< 150 mg/dL) to High ( $\geq 200$ mg/dL)	Clozapine	4	0 (0)
		Chlorpromazine	6	2 (33)
	Borderline ( $\geq 150$ mg/dL and < 200 mg/dL) to High ( $\geq 200$ mg/dL)	Clozapine	1	1 (100)
		Chlorpromazine	1	0 (0)

#### Weight Gain

Weight gain has occurred with the use of antipsychotics, including clozapine. Monitor weight during treatment with FAZACLO. Table 6 summarizes the data on weight gain by the duration of exposure pooled from 11 studies with clozapine and active comparators. The median duration

of exposure was 609, 728, and 42 days, in the clozapine, olanzapine, and chlorpromazine group, respectively.

**Table 6. Mean Change in Body Weight (kg) by duration of exposure from studies in adult subjects with schizophrenia**

Metabolic Parameter	Exposure Duration	Clozapine (N = 669)		Olanzapine (N = 442)		Chlorpromazine (N = 155)	
		n	Mean	n	Mean	n	Mean
Weight change from baseline	2 weeks (Day 11 – 17)	6	+0.9	3	+0.7	2	-0.5
	4 weeks (Day 21 – 35)	23	+0.7	8	+0.8	17	+0.6
	8 weeks (Day 49 – 63)	12	+1.9	13	+1.8	16	+0.9
	12 weeks (Day 70 – 98)	17	+2.8	5	+3.1	0	0
	24 weeks (154 – 182)	42	-0.6	12	+5.7	0	0
	48 weeks (Day 322 – 350)	3	+3.7	3	+13.7	0	0

Table 7 summarizes pooled data from 11 studies in adult subjects with schizophrenia demonstrating weight gain  $\geq 7\%$  of body weight relative to baseline. The median duration of exposure was 609, 728, and 42 days, in the clozapine, olanzapine, and chlorpromazine group, respectively.

**Table 7. Proportion of adult subjects in schizophrenia studies with weight gain  $\geq 7\%$  relative to baseline body weight**

Weight Change	Clozapine	Olanzapine	Chlorpromazine
N	669	442	155
$\geq 7\% \text{ (inclusive)}$	236 (35%)	203 (46%)	13 (8%)

## 5.10 Neuroleptic Malignant Syndrome

Antipsychotic drugs including FAZACLO can cause a potentially fatal symptom complex referred to as Neuroleptic Malignant Syndrome (NMS). Clinical manifestations of NMS include hyperpyrexia, muscle rigidity, altered mental status, and autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmias). Associated findings can include elevated creatine phosphokinase (CPK), myoglobinuria, rhabdomyolysis, and acute renal failure.

The diagnostic evaluation of patients with this syndrome is complicated. It is important to consider the presence of other serious medical conditions (e.g., agranulocytosis, infection, heat stroke, primary CNS pathology, central anticholinergic toxicity, extrapyramidal symptoms, and drug fever).

The management of NMS should include (1) immediate discontinuation of antipsychotic drugs and other drugs not essential to concurrent therapy, (2) intensive symptomatic treatment and medical monitoring, and (3) treatment of comorbid medical conditions. There is no general agreement about specific pharmacological treatments for NMS.

If a patient requires antipsychotic drug treatment after recovery from NMS, the potential reintroduction of drug therapy should be carefully considered. NMS can recur. Monitor closely if restarting treatment with antipsychotics.

NMS has occurred with clozapine monotherapy and with concomitant CNS-active medications, including lithium.

## **5.11 Fever**

During clozapine therapy, patients have experienced transient, clozapine-related fever. The peak incidence is within the first 3 weeks of treatment. While this fever is generally benign and self-limited, it may necessitate discontinuing treatment. The fever can be associated with an increase or decrease in WBC count. Carefully evaluate patients with fever to rule out agranulocytosis or infection. Consider the possibility of NMS [*see Warnings and Precautions (5.10)*].

## **5.12 Pulmonary Embolism**

Pulmonary embolism and deep vein thrombosis have occurred in patients treated with clozapine. Consider the possibility of pulmonary embolism in patients who present with deep-vein thrombosis, acute dyspnea, chest pain, or with other respiratory signs and symptoms. Whether pulmonary embolus and deep vein thrombosis can be attributed to clozapine or some characteristic(s) of patients is not clear.

## **5.13 Anticholinergic Toxicity**

FAZACLO has potent anticholinergic effects. Treatment with FAZACLO can result in CNS and peripheral anticholinergic toxicity. Use with caution in the presence of narrow-angle glaucoma, concomitant anticholinergic medications, prostatic hypertrophy, or other conditions in which anticholinergic effects can lead to significant adverse reactions.

Treatment with FAZACLO can result in gastrointestinal adverse reactions, including constipation, intestinal obstruction, fecal impaction, and paralytic ileus. Such reactions can be fatal. Constipation should be initially treated by ensuring adequate hydration and use of ancillary therapy such as bulk laxatives. Consultation with a gastroenterologist is advisable in more serious cases.

## **5.14 Interference with Cognitive and Motor Performance**

FAZACLO can cause sedation and impairment of cognitive and motor performance. Caution patients about operating hazardous machinery, including automobiles, until they are reasonably certain that FAZACLO does not affect them adversely. These reactions may be dose-related. Consider reducing the dose if they occur.

## **5.15 Tardive Dyskinesia**

Tardive dyskinesia (TD) has occurred in patients treated with antipsychotic drugs, including clozapine. The syndrome consists of potentially irreversible, involuntary, dyskinetic movements. The risk of TD and the likelihood that it will become irreversible are believed to increase with greater durations of treatment and higher total cumulative doses. However, the syndrome can develop after relatively brief treatment periods at low doses. Prescribe FAZACLO in a manner that is most likely to minimize the risk of developing TD. Use the lowest effective dose and the shortest duration necessary to control symptoms. Periodically assess the need for continued treatment. Consider discontinuing treatment if TD occurs. However, some patients may require treatment with FAZACLO despite the presence of the syndrome.

There is no known treatment for TD. However, the syndrome may remit partially or completely if treatment is discontinued. Antipsychotic treatment, itself, may suppress (or partially suppress) the signs and symptoms, and it has the potential to mask the underlying process. The effect of symptom suppression on the long-term course of TD is unknown.

## **5.16 Patients with Phenylketonuria**

Phenylketonuric patients should be informed that FAZACLO contains phenylalanine (a component of aspartame). Each 12.5 mg, orally disintegrating tablet contains 0.87 mg phenylalanine. Each 25 mg, orally disintegrating tablet contains 1.74 mg phenylalanine. Each 100 mg, orally disintegrating tablet contains 6.96 mg phenylalanine. Each 150 mg, orally disintegrating tablet contains 10.44 mg phenylalanine. Each 200 mg, orally disintegrating tablet contains 13.92 mg phenylalanine.

## **5.17 Cerebrovascular Adverse Reactions**

In controlled trials, elderly patients with dementia-related psychosis treated with some atypical antipsychotics had an increased risk (compared to placebo) of cerebrovascular adverse reactions (e.g., stroke, transient ischemic attack), including fatalities. The mechanism for this increased risk is not known. An increased risk cannot be excluded for FAZACLO or other antipsychotics or other patient populations. FAZACLO should be used with caution in patients with risk factors for cerebrovascular adverse reactions.

## **5.18 Recurrence of Psychosis and Cholinergic Rebound after Abrupt Discontinuation of FAZACLO**

If abrupt discontinuation of FAZACLO is necessary (because of agranulocytosis or another medical condition, for example), monitor carefully for the recurrence of psychotic symptoms and adverse reactions related to cholinergic rebound, such as profuse sweating, headache, nausea, vomiting and diarrhea.

## **6 ADVERSE REACTIONS**

The following adverse reactions are discussed in more detail in other sections of the labeling:

- Agranulocytosis [see Warnings and Precautions (5.1)].

- Orthostatic Hypotension, Bradycardia, and Syncope [*see Warnings and Precautions (5.3)*].
- Seizures [*see Warnings and Precautions (5.4)*].
- Myocarditis and Cardiomyopathy [*see Warnings and Precautions (5.5)*].
- Increased Mortality in Elderly Patients with Dementia-Related Psychosis [*see Warnings and Precautions (5.6)*].
- Eosinophilia [*see Warnings and Precautions (5.7)*].
- QT Interval Prolongation [*see Warnings and Precautions (5.8)*].
- Metabolic Changes (Hyperglycemia and Diabetes Mellitus, Dyslipidemia, and Weight Gain) [*see Warnings and Precautions (5.9)*].
- Neuroleptic Malignant Syndrome [*see Warnings and Precautions (5.10)*].
- Fever [*see Warnings and Precautions (5.11)*].
- Pulmonary Embolism [*see Warnings and Precautions (5.12)*].
- Anticholinergic Toxicity [*see Warnings and Precautions (5.13)*].
- Interference with Cognitive and Motor Performance [*see Warnings and Precautions (5.14)*].
- Tardive Dyskinesia [*see Warnings and Precautions (5.15)*].
- Patients with Phenylketonuria [*see Warnings and Precautions (5.16)*].
- Cerebrovascular Adverse Reactions [*see Warnings and Precautions (5.17)*].
- Recurrence of Psychosis and Cholinergic Rebound after Abrupt Discontinuation [*see Warnings and Precautions (5.18)*].

## 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The most commonly reported adverse reactions ( $\geq 5\%$ ) across clozapine clinical trials were: CNS reactions, including sedation, dizziness/vertigo, headache, and tremor; cardiovascular reactions, including tachycardia, hypotension, and syncope; autonomic nervous system reactions, including hypersalivation, sweating, dry mouth, and visual disturbances; gastrointestinal

reactions, including constipation and nausea; and fever. Table 8 summarizes the most commonly reported adverse reactions ( $\geq 5\%$ ) in clozapine-treated patients (compared to chlorpromazine-treated patients) in the pivotal, 6-week, controlled trial in treatment-resistant schizophrenia.

**Table 8. Common Adverse Reactions ( $\geq 5\%$ ) in the 6-Week, Randomized, Chlorpromazine-controlled Trial in Treatment-Resistant Schizophrenia**

Adverse Reaction	Clozapine (N = 126) (%)	Chlorpromazine (N = 142) (%)
Sedation	21	13
Tachycardia	17	11
Constipation	16	12
Dizziness	14	16
Hypotension	13	38
Fever (hyperthermia)	13	4
Hypersalivation	13	1
Hypertension	12	5
Headache	10	10
Nausea/vomiting	10	12
Dry mouth	5	20

Table 9 summarizes the adverse reactions reported in clozapine-treated patients at a frequency of 2% or greater across all clozapine studies (excluding the 2-year InterSePT™ Study). These rates are not adjusted for duration of exposure.

**Table 9. Adverse Reactions ( $\geq 2\%$ ) Reported in Clozapine-treated Patients (N=842) across all Clozapine Studies (excluding the 2-year InterSePT™ Study)**

Body System Adverse Reaction	Clozapine N = 842	Percentage of Patients
<b>Central Nervous System</b>		
Drowsiness/Sedation	39	
Dizziness/Vertigo	19	
Headache	7	
Tremor	6	
Syncope	6	
Disturbed Sleep/Nightmares	4	
Restlessness	4	
Hypokinesia/Akinesia	4	

<b>Body System</b> <b>Adverse Reaction</b>	<b>Clozapine</b> <b>N = 842</b>	<b>Percentage of</b> <b>Patients</b>
Agitation	4	
Seizures (convulsions)	3†	
Rigidity	3	
Akathisia	3	
Confusion	3	
Fatigue	2	
Insomnia	2	
<b>Cardiovascular</b>		
Tachycardia	25†	
Hypotension	9	
Hypertension	4	
<b>Gastrointestinal</b>		
Constipation	14	
Nausea	5	
Abdominal Discomfort/Heartburn	4	
Nausea/Vomiting	3	
Vomiting	3	
Diarrhea	2	
<b>Urogenital</b>		
Urinary Abnormalities	2	
<b>Autonomic Nervous System</b>		
Salivation	31	
Sweating	6	
Dry Mouth	6	
Visual Disturbances	5	
<b>Skin</b>		
Rash	2	
<b>Hemic/Lymphatic</b>		
Leukopenia/Decreased WBC/Neutropenia	3	
<b>Miscellaneous</b>		
Fever	5	
Weight Gain	4	

† Rate based on population of approximately 1700 exposed during premarket clinical evaluation of clozapine.

Table 10 summarizes the most commonly reported adverse reactions ( $\geq 10\%$  of the clozapine or olanzapine group) in the InterSePT™ Study. This was an adequate and well-controlled, two-year study evaluating the efficacy of clozapine relative to olanzapine in reducing the risk of suicidal behavior in patients with schizophrenia or schizoaffective disorder. The rates are not adjusted for duration of exposure.

**Table 10. Incidence of Adverse Reactions in Patients Treated with Clozapine or Olanzapine in the InterSePT™ Study ( $\geq 10\%$  in the clozapine or olanzapine group)**

Adverse Reactions	Clozapine N = 479	Olanzapine N = 477
	% Reporting	% Reporting
Salivary hypersecretion	48%	6%
Somnolence	46%	25%
Weight increased	31%	56%
Dizziness (excluding vertigo)	27%	12%
Constipation	25%	10%
Insomnia	20%	33%
Nausea	17%	10%
Vomiting	17%	9%
Dyspepsia	14%	8%

### Dystonia

*Class effect:* Symptoms of dystonia, prolonged abnormal contractions of muscle groups, may occur in susceptible individuals during the first few days of treatment. Dystonic symptoms include: spasm of the neck muscles, sometimes progressing to tightness of the throat, swallowing difficulty, difficulty breathing, and/or protrusion of the tongue. While these symptoms can occur at low doses, they occur more frequently and with greater severity with high potency and at higher doses of first generation antipsychotic drugs. An elevated risk of acute dystonia is observed in males and younger age groups.

### **6.2 Postmarketing Experience**

The following adverse reactions have been identified during post-approval use of clozapine. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

#### Central Nervous System

Delirium, EEG abnormal, myoclonus, paresthesia, possible cataplexy, status epilepticus, obsessive compulsive symptoms, and post-discontinuation cholinergic rebound adverse reactions.

#### Cardiovascular System

Atrial or ventricular fibrillation, ventricular tachycardia, QT interval prolongation, Torsades de Pointes, myocardial infarction, cardiac arrest, and periorbital edema.

#### Gastrointestinal System

Acute pancreatitis, dysphagia, salivary gland swelling.

**Hepatobiliary System**

Cholestasis, hepatitis, jaundice, hepatotoxicity, hepatic steatosis, hepatic necrosis, hepatic fibrosis, hepatic cirrhosis, liver injury (hepatic, cholestatic, and mixed), and liver failure.

**Urogenital System**

Acute interstitial nephritis, nocturnal enuresis, priapism, and renal failure.

**Skin**

Hypersensitivity reactions: photosensitivity, vasculitis, erythema multiforme, and Stevens-Johnson Syndrome.

**Musculoskeletal System**

Myasthenic syndrome and rhabdomyolysis.

**Respiratory System**

Aspiration, pleural effusion, pneumonia, lowers respiratory tract infection.

**Hemic and Lymphatic System**

Deep-vein thrombosis, elevated hemoglobin/hematocrit, erythrocyte sedimentation rate (ESR) increased, sepsis, thrombocytosis, and thrombocytopenia.

**Vision Disorders**

Narrow-angle glaucoma.

**Miscellaneous**

Creatine phosphokinase elevation, hyperuricemia, hyponatremia, and weight loss.

**7 DRUG INTERACTIONS**

**7.1 Potential for Other Drugs to Affect FAZACLO**

Clozapine is a substrate for many cytochrome P450 isozymes, in particular CYP1A2, CYP3A4, and CYP2D6. Use caution when administering FAZACLO concomitantly with drugs that are inducers or inhibitors of these enzymes.

**CYP1A2 Inhibitors**

Concomitant use of FAZACLO and CYP1A2 inhibitors can increase plasma levels of clozapine, potentially resulting in adverse reactions. Reduce the FAZACLO dose to one third of the original dose when FAZACLO is coadministered with strong CYP1A2 inhibitors (e.g., fluvoxamine, ciprofloxacin, or enoxacin). The FAZACLO dose should be increased to the original dose when

coadministration of strong CYP1A2 inhibitors is discontinued [*see Dosage and Administration (2.7), Clinical Pharmacology (12.3)*].

Moderate or weak CYP1A2 inhibitors include oral contraceptives and caffeine. Monitor patients closely when FAZACLO is coadministered with these inhibitors. Consider reducing the FAZACLO dosage if necessary [*see Dosage and Administration (2.7)*].

#### CYP2D6 and CYP3A4 Inhibitors

Concomitant treatment with FAZACLO and CYP2D6 or CYP3A4 inhibitors (e.g., cimetidine, escitalopram, erythromycin, paroxetine, bupropion, fluoxetine, quinidine, duloxetine, terbinafine, or sertraline) can increase FAZACLO levels and lead to adverse reactions [*see Clinical Pharmacology (12.3)*]. Use caution and monitor patients closely when using such inhibitors. Consider reducing the FAZACLO dose [*see Dosage and Administration (2.7)*].

#### CYP1A2 and CYP3A4 Inducers

Concomitant treatment with drugs that induce CYP1A2 or CYP3A4 can decrease the plasma concentration of clozapine, resulting in decreased effectiveness of FAZACLO. Tobacco smoke is a moderate inducer of CYP1A2. Strong CYP3A4 inducers include carbamazepine, phenytoin, St. John's wort, and rifampin. It may be necessary to increase the FAZACLO dose if used concomitantly with inducers of these enzymes. However, concomitant use of FAZACLO and strong CYP3A4 inducers is not recommended [*see Dosage and Administration (2.7)*].

Consider reducing the FAZACLO dosage when discontinuing coadministered enzyme inducers; because discontinuation of inducers can result in increased clozapine plasma levels and an increased risk of adverse reactions [*see Dosage and Administration (2.7)*].

#### Drugs that Cause QT Interval Prolongation

Use caution when administering concomitant medications that prolong the QT interval or inhibit the metabolism of FAZACLO. Drugs that cause QT prolongation include: specific antipsychotics (e.g., ziprasidone, iloperidone, chlorpromazine, thioridazine, mesoridazine, droperidol, and pimozide), specific antibiotics (e.g., erythromycin, gatifloxacin, moxifloxacin, sparfloxacin), Class 1A antiarrhythmics (e.g., quinidine, procainamide) or Class III antiarrhythmics (e.g., amiodarone, sotalol), and others (e.g., pentamidine, levomethadyl acetate, methadone, halofantrine, mefloquine, dolasetron mesylate, probucol or tacrolimus) [*see Warnings and Precautions (5.8)*].

## **7.2 Potential for FAZACLO to Affect Other Drugs**

Concomitant use of FAZACLO with other drugs metabolized by CYP2D6 can increase levels of these CYP2D6 substrates. Use caution when coadministering FAZACLO with other drugs that are metabolized by CYP2D6. It may be necessary to use lower doses of such drugs than usually prescribed. Such drugs include specific antidepressants, phenothiazines, carbamazepine, and Type 1C antiarrhythmics (e.g., propafenone, flecainide, and encainide).

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

Pregnancy Category B

#### Risk Summary

There are no adequate or well-controlled studies of clozapine in pregnant women.

Reproduction studies have been performed in rats and rabbits at doses up to 0.4 and 0.9 times, respectively, the maximum recommended human dose (MRHD) of 900 mg/day on a mg/m<sup>2</sup> body surface area basis. The studies revealed no evidence of impaired fertility or harm to the fetus due to clozapine. Because animal reproduction studies are not always predictive of human response, FAZACLO should be used during pregnancy only if clearly needed.

#### Clinical Considerations

Consider the risk of exacerbation of psychosis when discontinuing or changing treatment with antipsychotic medications during pregnancy and postpartum. Consider early screening for gestational diabetes for patients treated with antipsychotic medications [see *Warnings and Precautions* (5.9)]. Neonates exposed to antipsychotic drugs during the third trimester of pregnancy are at risk for extrapyramidal and/or withdrawal symptoms following delivery. Monitor neonates for symptoms of agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress, and feeding difficulties. The severity of complications can vary from self-limited symptoms to some neonates requiring intensive care unit support and prolonged hospitalization.

#### Animal Data

In embryofetal developmental studies, clozapine had no effects on maternal parameters, litter sizes, or fetal parameters when administered orally to pregnant rats and rabbits during the period of organogenesis at doses up to 0.4 and 0.9 times, respectively, the MRHD of 900 mg/day on a mg/m<sup>2</sup> body surface area basis.

In peri/postnatal developmental studies, pregnant female rats were administered clozapine over the last third of pregnancy and until day 21 postpartum. Observations were made on fetuses at birth and during the postnatal period; the offspring were allowed to reach sexual maturity and mated. Clozapine caused a decrease in maternal body weight but had no effects on litter size or body weights of either F1 or F2 generations at doses up to 0.4 times the MRHD of 900 mg/day on a mg/m<sup>2</sup> body surface area basis.

### 8.3 Nursing Mothers

FAZACLO is present in human milk. Because of the potential for serious adverse reactions in nursing infants from FAZACLO, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

## **8.4 Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

## **8.5 Geriatric Use**

There have not been sufficient numbers of geriatric patients in clinical studies utilizing FAZACLO to determine whether those over 65 years of age differ from younger subjects in their response to FAZACLO.

Orthostatic hypotension and tachycardia can occur with clozapine treatment [*see Boxed Warning and Warnings and Precautions (5.3)*]. Elderly patients, particularly those with compromised cardiovascular functioning, may be more susceptible to these effects.

Elderly patients may be particularly susceptible to the anticholinergic effects of clozapine, such as urinary retention and constipation [*see Warnings and Precautions (5.13)*].

Carefully select FAZACLO doses in elderly patients, taking into consideration their greater frequency of decreased hepatic, renal, or cardiac function, as well as other concomitant disease and other drug therapy. Clinical experience suggests that the prevalence of tardive dyskinesia appears to be highest among the elderly; especially elderly women [*see Warnings and Precautions (5.15)*].

## **8.6 Patients with Renal or Hepatic Impairment**

Dose reduction may be necessary in patients with significant impairment of renal or hepatic function. Clozapine concentrations may be increased in these patients, because clozapine is almost completely metabolized and then excreted [*see Dosage and Administration (2.8), Clinical Pharmacology (12.3)*].

## **8.7 CYP2D6 Poor Metabolizers**

Dose reduction may be necessary in patients who are CYP2D6 poor metabolizers. Clozapine concentrations may be increased in these patients, because clozapine is almost completely metabolized and then excreted [*see Dosage and Administration (2.8), Clinical Pharmacology (12.3)*].

# **10 OVERDOSAGE**

## **10.1 Overdosage Experience**

The most commonly reported signs and symptoms associated with FAZACLO overdose are: sedation, delirium, coma, tachycardia, hypotension, respiratory depression or failure; and hypersalivation. There are reports of aspiration pneumonia, cardiac arrhythmias, and seizure. Fatal overdoses have been reported with clozapine, generally at doses above 2500 mg. There have also been reports of patients recovering from overdoses well in excess of 4 g.



## 10.2 Management of Overdosage

For the most up-to-date information on the management of FAZACLO overdosage, contact a certified Regional Poison Control Center (1-800-222-1222). Telephone numbers of certified Regional Poison Control Centers are listed in the *Physicians' Desk Reference*®, a registered trademark of Thomson PDR. Establish and maintain an airway; ensure adequate oxygenation and ventilation. Monitor cardiac status and vital signs. Use general symptomatic and supportive measures. There are no specific antidotes for FAZACLO.

In managing overdosage, consider the possibility of multiple-drug involvement.

## 11 DESCRIPTION

FAZACLO® (clozapine), an atypical antipsychotic drug, is a tricyclic dibenzodiazepine derivative, 8-chloro-11-(4-methyl-1-piperazinyl)-5H-dibenzo [b,e] [1,4] diazepine. The structural formula is:



Mol. Wt. 326.83

FAZACLO is available as yellow, orally disintegrating tablets of 12.5 mg, 25 mg, 100 mg, 150 mg, and 200 mg for oral administration without water. FAZACLO tablets may be chewed.

Each orally disintegrating tablet contains clozapine equivalent to 12.5 mg, 25 mg, 100 mg, 150 mg, and 200 mg.

The active component of FAZACLO is clozapine. The remaining components are aminoalkyl methacrylate copolymer E, mannitol, aspartame, microcrystalline cellulose\*, silicified microcrystalline cellulose\*\*, crospovidone, natural and artificial mint flavor, sodium bicarbonate, citric acid, ferric oxide (yellow), and magnesium stearate

\*12.5, 25, and 100 mg tablets

\*\*150 and 200 mg tablets

THIS PRODUCT CONTAINS ASPARTAME AND IS NOT INTENDED FOR USE BY INFANTS. PHENYLKETONURICS: CONTAINS PHENYLALANINE [*see Warnings and Precautions (5.16)*. Phenylalanine is a component of aspartame. Each 12.5 mg, orally disintegrating tablet contains 1.6 mg aspartame, thus, 0.87 mg phenylalanine. Each 25 mg, orally disintegrating tablet contains 3.1 mg aspartame, thus, 1.74 mg phenylalanine. Each 100 mg,

orally disintegrating tablet contains 12.4 mg aspartame, thus, 6.96 mg phenylalanine. Each 150 mg, orally disintegrating tablet contains 18.6 mg aspartame, thus, 10.44 mg phenylalanine. Each 200 mg, orally disintegrating tablet contains 24.8 mg aspartame, thus, 13.92 mg phenylalanine. The allowable daily intake of aspartame is 50 mg per kilogram of body weight per day.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

The mechanism of action of FAZACLO is unknown. However, it has been proposed that the therapeutic efficacy of FAZACLO in schizophrenia is mediated through antagonism of the dopamine type 2 ( $D_2$ ) and the serotonin type 2A (5-HT<sub>2A</sub>) receptors. FAZACLO also acts as an antagonist at adrenergic, cholinergic, histaminergic and other dopaminergic and serotonergic receptors.

### 12.2 Pharmacodynamics

Clozapine demonstrated binding affinity to the following receptors: histamine H<sub>1</sub> ( $K_i$  1.1 nM), adrenergic  $\alpha_{1A}$  ( $K_i$  1.6 nM), serotonin 5-HT<sub>6</sub> ( $K_i$  4 nM), serotonin 5-HT<sub>2A</sub> ( $K_i$  5.4 nM), muscarinic M<sub>1</sub> ( $K_i$  6.2 nM), serotonin 5-HT<sub>7</sub> ( $K_i$  6.3 nM), serotonin 5-HT<sub>2C</sub> ( $K_i$  9.4 nM), dopamine D<sub>4</sub> ( $K_i$  24 nM), adrenergic  $\alpha_{2A}$  ( $K_i$  90 nM), serotonin 5-HT<sub>3</sub> ( $K_i$  95 nM), serotonin 5-HT<sub>1A</sub> ( $K_i$  120 nM), dopamine D<sub>2</sub> ( $K_i$  160 nM), dopamine D<sub>1</sub> ( $K_i$  270 nM), dopamine D<sub>5</sub> ( $K_i$  454 nM), and dopamine D<sub>3</sub> ( $K_i$  555 nM).

FAZACLO causes little or no prolactin elevation.

Clinical electroencephalogram (EEG) studies demonstrated that clozapine increases delta and theta activity and slows dominant alpha frequencies. Enhanced synchronization occurs. Sharp wave activity and spike and wave complexes may also develop. Patients have reported an intensification of dream activity during clozapine therapy. REM sleep was found to be increased to 85% of the total sleep time. In these patients, the onset of REM sleep occurred almost immediately after falling asleep.

### 12.3 Pharmacokinetics

#### Absorption

In man, clozapine tablets (25 and 100 mg) are equally bioavailable relative to a clozapine solution. FAZACLO Oral Suspension is bioequivalent to clozapine marketed tablets.

Following oral administration of 100 mg to 800 mg FAZACLO, once daily, the average steady-state peak plasma concentration was 275 ng/mL (range: 105 to 723 ng/mL), occurring at the average of 2.2 hours (range: 1 to 3.5 hours) after dosing. The average minimum concentration at steady state was 75 ng/mL (range: 11 to 198 ng/mL).

When FAZACLO was administered after a high fat meal there was no effect on the AUC<sub>ss</sub> or C<sub>minss</sub>, however C<sub>max</sub> was reduced about 20%, and there was a slight delay in T<sub>max</sub> of 0.5 hour

from a median  $T_{max}$  of 2.0 hours under fasted conditions to 2.5 hours under fed conditions. The decrease in  $C_{max}$  is not considered clinically relevant. Therefore FAZACLO may be taken without regard to meals.

### Distribution

Clozapine is approximately 97% bound to serum proteins. The interaction between clozapine and other highly protein-bound drugs has not been fully evaluated but may be important.

### Metabolism and Excretion

FAZACLO is almost completely metabolized prior to excretion, and only trace amounts of unchanged drug are detected in the urine and feces. FAZACLO is a substrate for many cytochrome P450 isozymes, in particular CYP1A2, CYP2D6, and CYP3A4. Approximately 50% of the administered dose is excreted in the urine and 30% in the feces. The demethylated, hydroxylated, and *N*-oxide derivatives are components in both urine and feces. Pharmacological testing has shown the desmethyl metabolite (norclozapine) to have only limited activity, while the hydroxylated and *N*-oxide derivatives were inactive. The mean elimination half-life of clozapine after a single 75 mg dose was 8 hours (range: 4-12 hours), compared to a mean elimination half-life of 12 hours (range: 4-66 hours), after achieving steady state with 100 mg twice daily dosing.

A comparison of single-dose and multiple-dose administration of clozapine demonstrated that the elimination half-life increased significantly after multiple dosing relative to that after single-dose administration, suggesting the possibility of concentration-dependent pharmacokinetics. However, at steady state, approximately dose-proportional changes with respect to AUC (area under the curve), peak, and minimum clozapine plasma concentrations were observed after administration of 37.5, 75, and 150 mg twice daily.

### Drug-Drug Interaction Studies

#### Fluvoxamine

A pharmacokinetic study was conducted in 16 schizophrenic patients who received clozapine under steady-state conditions. After coadministration of fluvoxamine for 14 days, mean trough concentrations of clozapine and its metabolites, *N*-desmethylclozapine and clozapine *N*-oxide, were elevated about three-fold compared to baseline steady state concentrations.

#### Paroxetine, Fluoxetine, and Sertraline

In a study of schizophrenic patients (n=14) who received clozapine under steady-state conditions, coadministration of paroxetine produced only minor changes in the levels of clozapine and its metabolites. However, other published reports describe modest elevations (less than two-fold) of clozapine and metabolite concentrations when clozapine was taken with paroxetine, fluoxetine, and sertraline.

### Specific Population Studies

#### Renal or Hepatic Impairment

No specific pharmacokinetic studies were conducted to investigate the effects of renal or hepatic impairment on the pharmacokinetics of clozapine. Higher clozapine plasma concentrations are likely in patients with significant renal or hepatic impairment when given usual doses.

#### CYP2D6 Poor Metabolizers

A subset (3%–10%) of the population has reduced activity of CYP2D6 (CYP2D6 poor metabolizers). These individuals may develop higher than expected plasma concentrations of clozapine when given usual doses.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### Carcinogenesis

No carcinogenic potential was demonstrated in long-term studies in mice and rats at doses up to 0.3 times and 0.4 times, respectively, the maximum recommended human dose (MRHD) of 900 mg/day on a mg/m<sup>2</sup> body surface area basis.

#### Mutagenesis

Clozapine was not genotoxic when tested in the following gene mutation and chromosomal aberration tests: the bacterial Ames test, the *in vitro* mammalian V79 in Chinese hamster cells, the *in vitro* unscheduled DNA synthesis in rat hepatocytes or the *in vivo* micronucleus assay in mice.

#### Impairment of Fertility

Clozapine had no effect on any parameters of fertility, pregnancy, fetal weight or postnatal development when administered orally to male rats 70 days before mating and to female rats for 14 days before mating at doses up to 0.4 times the MRHD of 900 mg/day on a mg/m<sup>2</sup> body surface area basis.

## **14 CLINICAL STUDIES**

### **14.1 Treatment-Resistant Schizophrenia**

The efficacy of clozapine in treatment-resistant schizophrenia was established in a multicenter, randomized, double-blind, active-controlled (chlorpromazine) study of clozapine in patients with a DSM-III diagnosis of schizophrenia who had inadequate responses to at least 3 different antipsychotics (from at least 2 different chemical classes) during the preceding 5 years. The antipsychotic trials must have been judged adequate; the antipsychotic dosages must have been

equivalent to or greater than 1000 mg per day of chlorpromazine for a period of at least 6 weeks, each without significant reduction of symptoms. There must have been no period of good functioning within the preceding 5 years. Patients must have had a baseline score of at least 45 on the investigator-rated Brief Psychiatric Rating Scale (BPRS). On the 18-item BPRS, 1 indicates the absence of symptoms, and 7 indicates severe symptoms; the maximum potential total BPRS score is 126. At baseline, the mean BPRS score was 61. In addition, patients must have had a score of at least 4 on at least two of the following four individual BPRS items: conceptual disorganization, suspiciousness, hallucinatory behavior, and unusual thought content. Patients must have had a Clinical Global Impressions – Severity Scale score of at least 4 (moderately ill).

In the prospective, lead-in phase of the trial, all patients (N=305) initially received single-blind treatment with haloperidol (the mean dose was 61 mg per day) for 6 weeks. More than 80% of patients completed the 6-week trial. Patients with an inadequate response to haloperidol (n=268) were randomized to double-blind treatment with clozapine (N=126) or chlorpromazine (N=142). The maximum daily clozapine dose was 900 mg; the mean daily dose was > 600 mg. The maximum daily chlorpromazine dose was 1800 mg; the mean daily dose was > 1200 mg.

The primary endpoint was treatment response, predefined as a decrease in BPRS score of at least 20% and either (1) a CGI-S score of ≤ 3 (mildly ill), or (2) a BPRS score of ≤ 35, at the end of 6 weeks of treatment. Approximately 88% of patients from the clozapine and chlorpromazine groups completed the 6-week trial. At the end of six weeks, 30% of the clozapine group responded to treatment, and 4% of the chlorpromazine group responded to treatment. The difference was statistically significant ( $p < 0.001$ ). The mean change in total BPRS score was -16 and -5 in the clozapine and chlorpromazine group, respectively; the mean change in the 4 key BPRS item scores was -5 and -2 in the clozapine and chlorpromazine group, respectively; and the mean change in CGI-S score was -1.2 and -0.4, in the clozapine and chlorpromazine group, respectively. These changes in the clozapine group were statistically significantly greater than in the chlorpromazine group ( $p < 0.001$  in each analysis).

## **14.2 Recurrent Suicidal Behavior in Schizophrenia or Schizoaffective Disorder**

The effectiveness of clozapine in reducing the risk of recurrent suicidal behavior was assessed in the International Suicide Prevention Trial (InterSePT™, a trademark of Novartis Pharmaceuticals Corporation). This was a prospective, randomized, open-label, active-controlled, multicenter, international, parallel-group comparison of clozapine (CLOZARIL®) versus olanzapine (Zyprexa®, a registered trademark of Eli Lilly and Company) in 956 patients with schizophrenia or schizoaffective disorder (DSM-IV) who were judged to be at risk for recurrent suicidal behavior. Only about one-fourth of these patients (27%) were considered resistant to standard antipsychotic drug treatment. To enter the trial, patients must have met one of the following criteria:

- They had attempted suicide within the three years prior to their baseline evaluation.
- They had been hospitalized to prevent a suicide attempt within the three years prior to their baseline evaluation.

- They demonstrated moderate-to-severe suicidal ideation with a depressive component within one week prior to their baseline evaluation.
- They demonstrated moderate-to-severe suicidal ideation accompanied by command hallucinations to do self-harm within one week prior to their baseline evaluation.

Dosing regimens for each treatment group were determined by individual investigators and were individualized by patient. Dosing was flexible, with a dose range of 200–900 mg/day for clozapine and 5–20 mg/day for olanzapine. For the 956 patients who received clozapine or olanzapine in this study, there was extensive use of concomitant psychotropics: 84% with antipsychotics, 65% with anxiolytics, 53% with antidepressants, and 28% with mood stabilizers. There was significantly greater use of concomitant psychotropic medications among the patients in the olanzapine group.


The primary efficacy measure was time to (1) a significant suicide attempt, including a completed suicide; (2) hospitalization due to imminent suicide risk, including increased level of surveillance for suicidality for patients already hospitalized; or (3) worsening of suicidality severity as demonstrated by “much worsening” or “very much worsening” from baseline in the Clinical Global Impression of Severity of Suicidality as assessed by the Blinded Psychiatrist (CGI-SS-BP) scale. A determination of whether or not a reported event met criterion 1 or 2 above was made by the Suicide Monitoring Board (SMB), a group of experts blinded to patient data.

A total of 980 patients were randomized to the study and 956 received study medication. Sixty-two percent of the patients were diagnosed with schizophrenia, and the remainder (38%) were diagnosed with schizoaffective disorder. Only about one-fourth of the total patient population (27%) was identified as “treatment-resistant” at baseline. There were more males than females in the study (61% of all patients were male). The mean age of patients entering the study was 37 years of age (range 18–69). Most patients were Caucasian (71%), 15% were Black, 1% were Asian, and 13% were classified as being of “other” races.

Patients treated with clozapine had a statistically significant longer delay in the time to recurrent suicidal behavior in comparison with olanzapine. This result should be interpreted only as evidence of the effectiveness of clozapine in delaying time to recurrent suicidal behavior and not a demonstration of the superior efficacy of clozapine over olanzapine.

The probability of experiencing (1) a significant suicide attempt, including a completed suicide, or (2) hospitalization because of imminent suicide risk, including increased level of surveillance for suicidality for patients already hospitalized, was lower for clozapine patients than for olanzapine patients at Week 104: clozapine 24% versus olanzapine 32%; 95% CI of the difference: 2%, 14% (Figure 2).

**Figure 2. Cumulative Probability of a Significant Suicide Attempt or Hospitalization to Prevent Suicide in Patients with Schizophrenia or Schizoaffective Disorder at High Risk of Suicidality**



## 16 HOW SUPPLIED/STORAGE AND HANDLING

### 16.1 How Supplied

#### 12.5 mg

1/4-inch diameter round, yellow tablet debossed with “A05” on one side.

Bottles of 100: 100 tablets in an HDPE bottle with child-resistant cap ... NDC No. 18860-101-10

#### 25 mg

5/16-inch diameter round, yellow tablet debossed with “A06” on one side.

Bottles of 100: 100 tablets in an HDPE bottle with child-resistant cap ... NDC No. 18860-102-10

Cartons of 48 for Institutional Use Only: 8 cards, 6 non child-resistant blisters per card ..... NDC No. 18860-102-01

#### 100 mg

1/2-inch diameter round, yellow tablet debossed with “A08” on one side.

Bottles of 100: 100 tablets in an HDPE bottle with child-resistant cap ... NDC No. 18860-104-10

Cartons of 48 for Institutional Use Only: 8 cards, 6 non child-resistant blisters per card ..... NDC No. 18860-104-01

### **150 mg**

9/16-inch diameter round, yellow tablet debossed with “A09” on one side.

Bottles of 100: 100 tablets in an HDPE bottle with child-resistant cap ... NDC No. 18860-105-10

Cartons of 48 for Institutional Use Only: 8 cards, 6 non child-resistant blisters per card ..... NDC No. 18860-105-01

### **200 mg**

5/8-inch diameter round, yellow tablet debossed with “A10” on one side.

Bottles of 100: 100 tablets in an HDPE bottle with child-resistant cap ... NDC No. 18860-106-10

Cartons of 48 for Institutional Use Only: 8 cards, 6 non child-resistant blisters per card ..... NDC No. 18860-106-01

## **16.2 Storage and Handling**

Store FAZACLO at 20°C to 25°C (68° to 77°F); excursions permitted to 15°C to 30°C (59° to 86°F). (See USP Controlled Room Temperature.) Protect from moisture.

**KEEP OUT OF REACH OF CHILDREN.**

FAZACLO tablets must remain in the original package until used by the patient.

Drug dispensing should not ordinarily exceed a weekly supply. If a patient is eligible for WBC count and ANC testing every 2 weeks, then a 2-week supply of FAZACLO can be dispensed. If a patient is eligible for WBC count and ANC testing every 4 weeks, then a 4-week supply of FAZACLO can be dispensed. Dispensing should be contingent upon the WBC count and ANC testing results.

## **17 PATIENT COUNSELING INFORMATION**

Discuss the following issues with patients and caregivers:

- **Agranulocytosis:** Prior to initiating treatment, educate patients and caregivers about the significant risk of developing agranulocytosis. Advise them to immediately report the appearance of signs or symptoms consistent with agranulocytosis or infection (e.g., fever; mucus membrane ulcers; skin, pharyngeal, vaginal, urinary, or pulmonary infection; or extreme weakness or lethargy) at any time during FAZACLO therapy [*see Warnings and Precautions (5.1)*].
  - Inform patients and caregivers that FAZACLO will be made available only through a special program designed to ensure the required blood monitoring in order to reduce the

risk of developing agranulocytosis. Inform patients and caregivers the WBC count and ANC will be monitored as follows:

- Weekly blood tests are required for the first 6 months.
- If acceptable WBC counts and ANCs (WBC count  $\geq 3500/\text{mm}^3$  and ANC  $\geq 2000/\text{mm}^3$ ) have been maintained during the first 6 months of continuous therapy, then WBC counts and ANCs can be monitored every 2 weeks for the next 6 months.
- Thereafter, if acceptable WBC counts and ANCs have been maintained during the second 6 months of continuous therapy, WBC counts and ANCs can be monitored every 4 weeks.
- **Orthostatic Hypotension, Bradycardia, and Syncope:** Inform patients and caregivers about the risk of orthostatic hypotension and syncope, especially during the period of initial dose titration. Instruct them to strictly follow the clinician's instructions for dosage and administration [see *Dosage and Administration (2.3)* and *Warnings and Precautions (5.3)*].
- **Seizures:** Inform patients and caregivers about the significant risk of seizure during FAZACLO treatment. Caution them about driving and any other potentially hazardous activity while taking FAZACLO [see *Warnings and Precautions (5.4)*].
- **Metabolic Changes (hyperglycemia and diabetes mellitus, dyslipidemia, weight gain):** Educate patients and caregivers about the risk of metabolic changes and the need for specific monitoring. The risks include hyperglycemia and diabetes mellitus, dyslipidemia, weight gain, and cardiovascular reactions. Educate patients and caregivers about the symptoms of hyperglycemia (high blood sugar) and diabetes mellitus (e.g., polydipsia, polyuria, polyphagia, and weakness). Monitor all patients for these symptoms. Patients who are diagnosed with diabetes or have risk factors for diabetes (obesity, family history of diabetes) should have their fasting blood glucose monitored before beginning treatment and periodically during treatment. Patients who develop symptoms of hyperglycemia should have assessments of fasting glucose. Clinical monitoring of weight is recommended [see *Warnings and Precautions (5.9)*].
- **Patients with Phenylketonuria:** Inform patients and caregivers that FAZACLO contains phenylalanine (a component of aspartame) [see *Warnings and Precautions (5.16)*].
- **Interference with Cognitive and Motor Performance:** Caution patients about operating hazardous machinery, including automobiles, until they are reasonably certain that FAZACLO does not affect them adversely [see *Warnings and Precautions (5.14)*].
- **Missed Doses and Re-initiating Treatment:** Inform patients and caregivers that if the patient misses taking FAZACLO for more than 2 days, they should not restart their medication at the same dosage but should contact their physician for dosing instructions [see *Dosage and Administration (2.6)* and *Warnings and Precautions (5.3)*].
- **QT Interval Prolongation:** Advise patients to consult their clinician immediately if they feel faint, lose consciousness or have signs or symptoms suggestive of arrhythmia. Instruct patients to not take FAZACLO with other drugs that cause QT interval prolongation. Instruct patients to inform their clinicians that they are taking FAZACLO before any new drug [see *Warnings and Precautions (5.8)* and *Drug Interactions (7.1)*].

- Concomitant Medication: Advise patients and caregivers to notify the physician if they are taking or plan to take any prescription or over-the-counter drugs or alcohol. [see *Drug Interactions (7)*].
- Pregnancy: Patients and caregivers should notify the physician if the patient becomes pregnant or intends to become pregnant during therapy. [see *Use in Specific Populations (8.1)*].
- Nursing: Advise patients and caregivers that the patient should not breast feed an infant if they are taking FAZACLO. [see *Use in Specific Populations (8.3)*].
- Administration: Patients should be advised that FAZACLO tablets should remain in the original package until immediately before use [see *Dosage and Administrations (2.2)*].

Distributed by:

**Jazz Pharmaceuticals, Inc.**  
Palo Alto, CA 94304 USA  
U.S. Patent Nos. 6,024,981; 6,221,392; and 6,106,861  
See U.S. Patent No. 6,155,423 for product sold in blister pack.

FazaClo® is a registered trademark of Jazz Pharmaceuticals plc or its subsidiaries.  
FZCLpc-13-01 Revised: July 2013  
Printed in USA  
Medical Affairs 1-877-FAZACLO (1-877-329-2256)