

Drug Facts**Active ingredient** *Purpose*
(in each capsule)Aspirin (NSAID)* 325mg . . . Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug**Uses**

- for the temporary relief of minor aches and pains associated with:
 - a cold • headache • backache • toothache
 - premenstrual & menstrual cramps
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

↓ Continue Reading 

Drug Facts (continued)

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling
- shock • asthma (wheezing)

This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you

Drug Facts (continued)

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days

Reference ID: **3244830**

Drug Facts (continued)

- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, consult a doctor get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
 - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Drug Facts (continued)**Other information**

- save carton for full directions and warnings
- store at 15 - 30°C (59 - 86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

Inactive ingredients anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Questions or comments


Call 1-800-123-4567 8:30 AM - 4:00 PM ET,
Monday - Friday



PLx Pharma Inc.
8285 El Rio, Suite 130
Houston, TX 77054
Tel: 713-842-1249
Fax: 713-842-3052

Distributed by: Company XXX
Visit us at www.companyxxx.com

Fold Out Panel



NDC: 1234-5678-90

ASPIRIN
Aspirin Capsules, 325mg
Pain Reliever / Fever Reducer (NSAID)



30 Liquid Filled Capsules

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL WITH REPEATED TEXT "SEALED FOR YOUR PROTECTION" UNDER CAP IS MISSING OR BROKEN.

Drug Facts	Purpose	Active ingredient (in each capsule)	Uses	Warnings	Continue Reading
		Aspirin (NSAID)* 325mg . . . Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug	<ul style="list-style-type: none"> • for the temporary relief of minor aches and pains associated with: <ul style="list-style-type: none"> • a cold • headache • backache • toothache • premenstrual & menstrual cramps • minor pain of arthritis • temporarily reduces fever 	<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p>	<p>↓  Continue Reading</p>

Drug facts continued under label

copy code
Lot: XXXXXXXX
Exp: XX/XX/XXXX

No Print
Glue Area

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

No Print
Glue Area

Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
 - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at 15 - 30°C (59 - 86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

Inactive ingredients

anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Questions or comments

Call 1-800-123-4567 8:30 AM - 4:00 PM ET, Monday - Friday



non printing
dieline

ASPIRIN
Pain Reliever / Fever Reducer (NSAID)
Aspirin Capsules, 325mg
30 Liquid Filled Capsules



TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL WITH REPEATED TEXT "SEALED FOR YOUR PROTECTION" UNDER CAP IS MISSING OR BROKEN.

30 Liquid Filled Capsules

ASPIRIN
Aspirin Capsules, 325mg
Pain Reliever / Fever Reducer (NSAID)

NDC# 1234-5678-90

RETAIN CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule)

Aspirin (NSAID)* 325mg Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purpose

Uses

- for the temporary relief of minor aches and pains associated with:
 - a cold
 - headache
 - backache
 - toothache
 - premenstrual & menstrual cramps
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older



6 28037 13550 4

LOT & EXP

Reference ID: 3244830

Drug Facts**Active ingredient (in each capsule) Purpose**

Aspirin (NSAID)* 325mg Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of minor aches and pains associated with:
 - a cold • headache • backache • toothache
 - premenstrual & menstrual cramps
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling
- shock • asthma (wheezing)

This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Reference ID: 3244830

↓ Continue Reading

**Drug Facts (continued)****Stop use and ask doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use.

It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, consult a doctor get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
 - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at 15 - 30°C (59 - 86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

Inactive ingredients anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Questions or comments

Call 1-800-123-4567 8:30 AM - 4:00 PM ET, Monday - Friday



PLx Pharma Inc.
8285 El Rio, Suite 130
Houston, TX 77054
Tel: 713-842-1249
Fax: 713-842-3052
Distributed by: Company XXX
Visit us at www.companyxxx.com

NDC#: 1234-5678-90

ASPIRIN

Aspirin Capsules, 325mg

Pain Reliever / Fever Reducer (NSAID)

120 Liquid Filled Capsules

Drug Facts	Purpose
Active ingredient (in each capsule) Aspirin (NSAID)* 325mg Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug	Purpose Pain reliever/fever reducer
Uses	
<ul style="list-style-type: none"> • for the temporary relief of minor aches and pains associated with: <ul style="list-style-type: none"> • a cold • headache • backache • toothache • premenstrual & menstrual cramps • minor pain of arthritis • temporarily reduces fever 	
Warnings	
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.	
Allergy alert: Aspirin may cause a severe allergic reaction which may include: <ul style="list-style-type: none"> • hives • facial swelling • shock • asthma (wheezing) This product contains soy.	
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: <ul style="list-style-type: none"> • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed 	
Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.	
Ask a doctor before use if	
<ul style="list-style-type: none"> • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma 	
Ask a doctor or pharmacist before use if you are	
<ul style="list-style-type: none"> • taking a prescription drug for diabetes, gout or arthritis 	
<p>→ Continue Reading</p>	

PEEL HERE FOR MORE DRUG FACTS



Drug facts continued under label

copy code

Lot: XXXXXXXX
Exp: XX/XXXXXX

FOR POSITION ONLY

No Print
Glue Area

120 Liquid Filled Capsules
Pain Reliever / Fever Reducer (NSAID)
ASPIRIN



TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL WITH REPEATED
TEXT "SEALED FOR YOUR PROTECTION" UNDER CAP IS MISSING OR BROKEN.

RETAIN CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each capsule)	Purpose
Aspirin (NSAID)* 325mg	Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of minor aches and pains associated with:
 - a cold • headache • backache • toothache
 - premenstrual & menstrual cramps • minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling
- shock • asthma (wheezing)

This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older

NDC#: 1234-5678-90

120 Liquid Filled Capsules

ASPIRIN
Aspirin Capsules, 325mg
Pain Reliever / Fever Reducer (NSAID)

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
 - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at 15 - 30°C (59 - 86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

Inactive ingredients

anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Questions or comments

Call 1-800-123-4567 8:30 AM - 4:00 PM ET, Monday - Friday

FOR POSITION ONLY

No Print
Glue Area



non printing
dieline

PLx Pharma Inc.
8285 El Rio, Suite 130
Houston, TX 77054
Tel: 713-842-1249
Fax: 713-842-3052

Distributed by: Company XXX
Visit us at www.companyxxx.com

Lot & Exp



ASPIRIN

Aspirin Capsules, 325mg
Pain Reliever / Fever Reducer (NSAID)

Pharmaceutics International, Hunt Valley, MD 21031

ASPIRIN

Aspirin Capsules, 325mg
(NSAID)

EXPIRES: 00/00/2013 LOT#: 012345

Pharmaceutics International
Hunt Valley, MD 21031



ASPIRIN

Aspirin Capsules, 325mg
(NSAID)

EXPIRES: 00/00/2013 LOT#: 012345

Pharmaceutics International
Hunt Valley, MD 21031



ASPIRIN

Aspirin Capsules, 325mg
(NSAID)

EXPIRES: 00/00/2013 LOT#: 012345

Pharmaceutics International
Hunt Valley, MD 21031



TO OPEN:

Separate at the perforations and peel back paper from arrow.

Reference ID: B214830

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

• take more or for a longer time than directed



ASPIRIN

Aspirin Capsules, 325mg
(NSAID)

EXPIRES: 00/00/2013 LOT#: 012345

Pharmaceutics International
Hunt Valley, MD 21031

ASPIRIN

Aspirin Capsules, 325mg
(NSAID)

EXPIRES: 00/00/2013 LOT#: 012345

Pharmaceutics International
Hunt Valley, MD 21031

ASPIRIN

Aspirin Capsules, 325mg
(NSAID)

EXPIRES: 00/00/2013 LOT#: 012345

Pharmaceutics International
Hunt Valley, MD 21031

ASPIRIN

Aspirin Capsules, 325mg
(NSAID)

EXPIRES: 00/00/2013 LOT#: 012345

Pharmaceutics International
Hunt Valley, MD 21031

FOR POSITION ONLY

TAMPER EVIDENT: DO NOT USE IF BLISTER IS OPEN OR TORN OR IF THE BAND AROUND THE CAPSULE IS MISSING OR BROKEN

Drug Facts

Active ingredient (in each capsule)

Purpose

Aspirin (NSAID)* 325mg Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of minor aches and pains associated with:
 - a cold • headache • backache • toothache
 - premenstrual & menstrual cramps • minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling
- shock • asthma (wheezing)

This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed.

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn

Drug Facts (continued)

- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood • have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
- do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings • store at 15 - 30°C (59 - 86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blister is open or torn or if the band around the capsule is missing or broken

FOR POSITION ONLY

FOR POSITION ONLY

Drug Facts (continued)

Inactive ingredients anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Drug Facts (continued)

Questions or comments

Call 1-800-123-4567 8:30 AM - 4:00 PM ET, Monday - Friday



7 Liquid Filled Capsules

ASPIRIN
Aspirin Capsules, 325mg
Pain Reliever / Fever Reducer (NSAID)

NDC#: 1234-5678-90

READ AND KEEP CARTON
FOR COMPLETE WARNINGS
AND INFORMATION



Visit us at www.aspirinxxx.com
Distributed by: Company XXX



PLX Pharma Inc.
8285 El Rio, Suite 130
Houston, TX 77054
Tel: 713-842-1749
Fax: 713-842-3052

Reference ID: 3244830

non printing
deline

FOR POSITION ONLY

Drug Facts

Active ingredient (in each capsule) **Purpose**
Aspirin (NSAID)* 325mg Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of minor aches and pains associated with:
 - a cold • headache • backache • toothache
 - premenstrual & menstrual cramps • minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
• hives • facial swelling
• shock • asthma (wheezing)
This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn

Drug Facts (continued)

- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood • have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
- do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings • store at 15 - 30°C (59 - 86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blister is open or torn or if the band around the capsule is missing or broken



FOR POSITION ONLY

FOR POSITION ONLY

Drug Facts (continued)

Inactive ingredients anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, medium chain triglycerides, pharmaceutical ink, potassium chloride, soy lecithin, titanium dioxide

Drug Facts (continued)

Questions or comments

Call 1-800-123-4567 8:30 AM - 4:00 PM ET, Monday - Friday

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

ASPIRIN Aspirin Capsules, 325mg (NSAID)

28 Liquid Filled Capsules

ASPIRIN
Aspirin Capsules, 325mg
Pain Reliever / Fever Reducer (NSAID)

NDC#: 1234-5678-90

TAMPER EVIDENT: DO NOT USE IF BLISTER IS OPEN OR TORN OR IF THE BAND AROUND THE CAPSULE IS MISSING OR BROKEN

ASPIRIN
Aspirin Capsules, 325mg
Pain Reliever / Fever Reducer (NSAID)



LOT & EXP

Distributed by: Company XXX
Visit us at www.companyxxx.com

Plx Pharma Inc.
8285 El Rio, Suite 130
Houston, TX 77054
Tel: 713-842-1249
Fax: 713-842-3052



ASPIRIN
Aspirin Capsules, 325mg
Pain Reliever / Fever Reducer (NSAID)

Reference ID: 3244830

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER
01/14/2013