



**COLORS**

Note: PMS Swatch on proof does not represent exact color (b) (4)

AUTHORIZED SIGNATURE \_\_\_\_\_ DATE \_\_\_\_\_

PROOF APPROVED  REPROOF REQUIRED

**HEADING:**  
Helvetica Bold  
Condensed Italic  
10 pt.

**SUBHEADS:**  
Helvetica Bold  
Condensed Italic  
8 pt.

**BODY TEXT:**  
Helvetica Condensed/  
Helvetica Bold Condensed  
8 pt.

(b) (4)

G&W Labs

G&W dwg# 1813

2+3/8 x 1 x 4+3/4

reverse tuck, slit lock - v-notched

.016 sbs

pgm# s5806

approved

date

<b>Drug Facts</b>	
<b>Active ingredient (in each rectal suppository)</b> Acetaminophen 120 mg	<b>Purpose</b> Pain reliever/fever reducer
<b>Uses temporarily</b> • reduces fever • relieves minor aches, pains, and headache	
<b>Warnings</b> For rectal use only <b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if your child inserts • more than 5 suppositories in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen <b>Allergy alert:</b> acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.	
<b>Do not use</b> • if your child is allergic to acetaminophen • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	

<b>Drug Facts (continued)</b>	
Ask a doctor before use if your child has liver disease	
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin	
<b>Stop use and ask a doctor if</b> • fever lasts more than 3 days (72 hours), or recurs • you need to use this product for pain for more than 5 days continuously Severe or recurrent pain, or high or continued fever may indicate a serious illness.	
<b>Keep out of reach of children.</b> If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for children even if you do not notice any signs or symptoms.	
<b>Directions</b> This product does not contain directions or complete warnings for adult use. • do not use more than directed • remove foil wrapper • insert suppository well up into rectum • children 3-6 years • 1 suppository every 4 to 6 hours while symptoms persist • do not exceed 5 suppositories in any 24-hour period • children under 3 years: ask a doctor	

<b>Drug Facts (continued)</b>	<b>Drug Facts (continued)</b>
<b>Other information</b> • for your safety, suppositories are packaged in tamper-evident sealed foil. Do not use if foil is torn or open. • store at 8°-25°C (46°-77°F)	<b>Inactive ingredients</b> butylated hydroxyanisole, butylated hydroxytoluene, glyceryl stearate, hydrogenated vegetable oil, polyethylene glycol 100 stearate, polysorbate 80
<b>Questions?</b> call 1-800-922-1038 Monday-Friday 9am-4pm EST	

G&W Laboratories, Inc., 111 Coolidge Street, South Plainfield, NJ 07080 Visit our website @ www.gwllabs.com

NDC 0713-0118-12

**Pediatric**  
**ACEPHEN™**  
**ACETAMINOPHEN SUPPOSITORIES USP**

ASPIRIN FREE **120 mg**

**12 Rectal Suppositories** Pain Reliever • Fever Reducer

FOR YOUR SAFETY, SUPPOSITORIES ARE PACKAGED IN TAMPER-EVIDENT SEALED FOIL. DO NOT USE IF SEAL IS TORN OR OPEN.

UPC  
N3-0713-0118-12-

10-11812GW23

50 RECTAL SUPPOSITORIES  
 ASPIRIN FREE  
**120 mg**  
 ACETAMINOPHEN  
 SUPPOSITORIES USP  
**ACEPHEN™**  
 Pediatric  
 G&W

Pain Reliever • Fever Reducer

**Drug Facts**

**Active ingredient (in each rectal suppository)**      **Purpose**  
 Acetaminophen 120 mg ..... Pain reliever/fever reducer

**Uses temporarily** • reduces fever • relieves minor aches, pains, and headache

**Warnings**  
**For rectal use only**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child inserts  
 • more than 5 suppositories in 24 hours, which is the maximum daily amount  
 • with other drugs containing acetaminophen

**Allergy alert:** acetaminophen may cause severe skin reactions.  
 Symptoms may include: • skin reddening • blisters • rash  
 If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
 • if your child is allergic to acetaminophen  
 • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if your child has liver disease**  
**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin**

**Stop use and ask a doctor if**  
 • fever lasts more than 3 days (72 hours), or recurs  
 • you need to use this product for pain for more than 5 days continuously  
 Severe or recurrent pain, or high or continued fever may indicate a serious illness.

**Keep out of reach of children.** If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**  
 This product does not contain directions or complete warnings for adult use.  
 • do not use more than directed  
 • remove foil wrapper  
 • insert suppository well up into rectum  
 • children 3-6 years  
 • 1 suppository every 4 to 6 hours while symptoms persist  
 • do not exceed 5 suppositories in any 24-hour period  
 • children under 3 years: ask a doctor

**Other information**  
 • for your safety, suppositories are packaged in tamper-evident sealed foil. Do not use if foil is torn or open.  
 • store at 8°-25°C (46°-77°F)

**Inactive ingredients** butylated hydroxyanisole, butylated hydroxytoluene, glyceryl stearate, hydrogenated vegetable oil, polyethylene glycol 100 stearate, polysorbate 80

**Questions?** call 1-800-922-1038  
 Monday-Friday 9am-4pm EST



G&W Laboratories, Inc.  
 111 Coolidge Street  
 South Plainfield, NJ 07080  
 Visit our website @ www.gwlab.com



NDC 0713-0118-50

Pediatric  
**ACEPHEN™**  
 ACETAMINOPHEN  
 SUPPOSITORIES USP

ASPIRIN FREE      **120 mg**  
 Pain Reliever • Fever Reducer

50 RECTAL SUPPOSITORIES



**TAMPER-EVIDENT:**  
 Individually sealed in foil  
 for your protection.  
 Do not use if foil is torn or open.

UPC Code Placement  
 N3-0713-0118-50-

10-11850GW8



**COLORS**  
 Note: PMS Swatch on proof does not represent exact color.  
 (b) (4)

AUTHORIZED SIGNATURE      DATE  
 \_\_\_\_\_  
 \_\_\_\_\_

PROOF APPROVED     REPROOF REQUIRED

HEADING:  
 Helvetica Bold  
 Condensed Italic  
 10 pt.  
 SUBHEADS:  
 Helvetica Bold  
 Condensed Italic  
 8 pt.  
 BODY TEXT:  
 Helvetica  
 Condensed/  
 Helvetica Bold  
 Condensed  
 6 pt.

AREA  
LOT & EXP

HEADING:  
Helvetica Bold  
Condensed Italic  
10 pt.

SUBHEADS:  
Helvetica Bold  
Condensed Italic  
8 pt.

BODY TEXT:  
Helvetica Condensed/  
Helvetica Bold Condensed  
6 pt.

Drug Facts	
<b>Active ingredient (in each rectal suppository)</b>	<b>Purpose</b>
Acetaminophen 120 mg	Pain reliever/fever reducer
<b>Uses</b> temporarily • reduces fever • relieves minor aches, pains, and headache	
<b>Warnings</b>	
<b>For rectal use only</b>	
<b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if your child inserts more than 5 suppositories in 24 hours, which is the maximum daily amount	
• with other drugs containing acetaminophen	
<b>Allergy alert:</b> acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash	
If a skin reaction occurs, stop use and seek medical help right away.	
<b>Do not use</b>	
• if your child is allergic to acetaminophen	
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	
<b>Ask a doctor before use if your child has liver disease</b>	
<b>Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin</b>	
<b>Stop use and ask a doctor if</b>	
• fever lasts more than 3 days (72 hours), or recurs	
• you need to use this product for pain for more than 5 days continuously	
Severe or recurrent pain, or high or continued fever may indicate a serious illness.	
<b>Keep out of reach of children.</b> If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for children even if you do not notice any signs or symptoms.	
<b>Directions</b>	
This product does not contain directions or complete warnings for adult use.	
• do not use more than directed	
• remove foil wrapper	
• insert suppository well up into rectum	
• children 3-6 years	
• 1 suppository every 4 to 6 hours while symptoms persist	
• do not exceed 5 suppositories in any 24-hour period	
• children under 3 years: ask a doctor	
<b>Other information</b>	
• for your safety, suppositories are packaged in tamper-evident sealed foil. Do not use if foil is torn or open.	
• store at 8°-25°C (46°-77°F)	
<b>Inactive ingredients</b> butylated hydroxyanisole, butylated hydroxytoluene, glyceryl stearate, hydrogenated vegetable oil, polyethylene glycol 100 stearate, polysorbate 80	
<b>Questions?</b> call 1-800-922-1038 Monday-Friday 9am-4pm EST	

Established since 1919, G&W Laboratories, Inc. is one of the largest manufacturers of quality suppositories in the United States. We take special care to ensure that all products are of the finest quality you can buy



G&W Laboratories, Inc.  
111 Coolidge Street  
South Plainfield, NJ 07080  
Visit our website @ www.gwlab.com



NDC 0713-0118-01

# Pediatric ACEPHEN™ ACETAMINOPHEN SUPPOSITORIES USP

ASPIRIN FREE **120 mg**

Pain Reliever • Fever Reducer

100 Rectal Suppositories

FOR YOUR SAFETY, SUPPOSITORIES  
ARE PACKAGED IN TAMPER-EVIDENT  
SEALED FOIL. DO NOT USE IF  
FOIL IS TORN OR OPEN.

80% UPC  
N3-0713-0118-01



### COLORS

Note: PMS Swatch on proof does not represent exact color. (b) (4)

AUTHORIZED SIGNATURE

DATE

PROOF APPROVED

REPROOF REQUIRED

10-11801GW4

-----  
**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**  
-----

/s/  
-----

THERESA M MICHELE  
03/25/2014