

12 of 5
PP#



<p>Drug Facts (continued)</p> <p>Warnings</p> <ul style="list-style-type: none"> • Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 4 doses in 24 hours, which is the maximum daily amount for ages 6 to 11 months, which is the maximum more than 5 doses in 24 hours, which is the maximum daily amount for ages 12 to 36 months. • Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> • rash • blisters • If a skin reaction occurs, stop use and seek medical help right away. <p>Do not use</p> <ul style="list-style-type: none"> • if you are allergic to acetaminophen. • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> • the child has liver disease. • the child is taking the blood thinning drug warfarin. 	<p>Drug Facts (continued)</p> <p>Warnings</p> <ul style="list-style-type: none"> • Stop use and ask a doctor if: <ul style="list-style-type: none"> • fever lasts more than 3 days (72 hours), or recurs. • pain lasts more than 5 days or gets worse. • new symptoms occur. • redness or swelling is present in the painful area. • These may be signs of a serious condition. • Keep out of reach of children. If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children. • Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> • rash • blisters • If a skin reaction occurs, stop use and seek medical help right away. <p>Do not use</p> <ul style="list-style-type: none"> • For rectal use only. • if you are allergic to acetaminophen. <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> • the child has liver disease. • the child is taking the blood thinning drug warfarin.
---	---

<p>Drug Facts (continued)</p> <p>Warnings</p> <ul style="list-style-type: none"> • Stop use and ask a doctor if: <ul style="list-style-type: none"> • fever lasts more than 3 days (72 hours), or recurs. • pain lasts more than 5 days or gets worse. • new symptoms occur. • redness or swelling is present in the painful area. • These may be signs of a serious condition. • Keep out of reach of children. If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children. • Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> • rash • blisters • If a skin reaction occurs, stop use and seek medical help right away. <p>Do not use</p> <ul style="list-style-type: none"> • For rectal use only. • if you are allergic to acetaminophen. <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> • the child has liver disease. • the child is taking the blood thinning drug warfarin. 	<p>Drug Facts (continued)</p> <p>Warnings</p> <ul style="list-style-type: none"> • Stop use and ask a doctor if: <ul style="list-style-type: none"> • fever lasts more than 3 days (72 hours), or recurs. • pain lasts more than 5 days or gets worse. • new symptoms occur. • redness or swelling is present in the painful area. • These may be signs of a serious condition. • Keep out of reach of children. If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children. • Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> • rash • blisters • If a skin reaction occurs, stop use and seek medical help right away. <p>Do not use</p> <ul style="list-style-type: none"> • For rectal use only. • if you are allergic to acetaminophen. <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> • the child has liver disease. • the child is taking the blood thinning drug warfarin.
---	---

<p>Drug Facts</p> <p>Active ingredient (in each rectal suppository)</p> <p>Acetaminophen, USP 80 mg Pain reliever/fever reducer</p> <p>Purposes</p> <ul style="list-style-type: none"> • relieves minor aches, pains, and headache temporarily • reduces fever 	<p>Drug Facts</p> <p>Active ingredient (in each rectal suppository)</p> <p>Acetaminophen, USP 80 mg Pain reliever/fever reducer</p> <p>Purposes</p> <ul style="list-style-type: none"> • relieves minor aches, pains, and headache temporarily • reduces fever
---	---

NDC 0472-1200-06

Infants' *Ages 6-36 months*



FeverAll[®]

Acetaminophen Suppositories

Pain reliever/fever reducer

80 mg

6 Rectal Suppositories

80 mg each

<p>Drug Facts (continued)</p> <p>Other information - store at 2-27°C (35°-80°F)</p> <ul style="list-style-type: none"> • do not use if imprinted suppository wrapper is opened or damaged <p>Inactive ingredients</p> <p>glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80</p> <p>Questions? 1-800-432-6534 between 9 am and 4 pm EST, Monday-Friday.</p> <p>12000613M1 VC101916</p>	<p>Drug Facts (continued)</p> <p>Other information - store at 2-27°C (35°-80°F)</p> <ul style="list-style-type: none"> • do not use if imprinted suppository wrapper is opened or damaged <p>Inactive ingredients</p> <p>glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80</p> <p>Questions? 1-800-432-6534 between 9 am and 4 pm EST, Monday-Friday.</p> <p>12000613M1 VC101916</p>
---	---

FeverAll[®]

Acetaminophen
Suppositories
Pain reliever/fever reducer

12000613M1 VC101916

3 04721 200006 1



Actavis
Div. of Actavis Mid-Atlantic, LLC, Lexington, NC 28059, USA. Made in USA.

Temper-evident: Suppositories are individually wrapped. Do not use if imprinted wrapper is opened or damaged.

(b) (4)

FeverAll.
Acetaminophen Suppositories
Pain reliever/fever reducer
80 mg each

Infants'
Ages 6-36 months
FeverAll.
Acetaminophen Suppositories
 Pain reliever/fever reducer

50 Rectal Suppositories **80 mg each**

NOT FOR RETAIL SALE
NDC 0472-1200-50



Dist. by Actavis Mid Atlantic LLC
 Lincolnton, NC 28092 USA Made in USA

**Tamper-evident: Suppositories are individually wrapped.
 Do not use if imprinted wrapper is opened or damaged.**

12000813M1 VC101917



3 04721 20050 4

Drug Facts (continued)

Active ingredient (in each rectal suppository) Acetaminophen, USP 80 mg

Purposes Pain reliever/fever reducer

Uses temporarily • reduces fever • relieves minor aches, pains, and headache

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes • more than 4 doses in 24 hours, which is the maximum daily amount for ages 6 to 11 months • more than 5 doses in 24 hours, which is the maximum daily amount for ages 12 to 36 months • with other drugs containing acetaminophen
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
 If a skin reaction occurs, stop use and seek medical help right away.
 For rectal use only.

Drug Facts (continued)

Do not use
 • if you are allergic to acetaminophen,
 • with any other drug containing acetaminophen (prescription or nonprescription),
 • if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if
 • the child has liver disease,
 • the child is taking the blood thinning drug warfarin.


Stop use and ask a doctor if
 • fever lasts more than 3 days (72 hours), or recurs,
 • pain lasts more than 3 days or gets worse,
 • new symptoms occur,
 • redness or swelling is present in the painful area.
 These may be signs of a serious condition.

Keep out of reach of children. If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

Other information
 • store at 2°-27°C (35°-80°F)
 • do not use if imprinted suppository wrapper is opened or damaged

Inactive ingredients glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polyorbate 80

Questions? 1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.



12 of 5
PP#



<p>Ask a doctor before use if</p> <ul style="list-style-type: none"> the child has liver disease. the child is taking the blood thinning drug warfarin. <p>Do not use</p> <ul style="list-style-type: none"> if you are allergic to acetaminophen. with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <p>Warnings</p> <p>Liver warnings: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount.</p> <p>Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin redness hives skin peeling <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>For rectal use only.</p> <p>Do not use</p> <ul style="list-style-type: none"> if you are allergic to acetaminophen. with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> the child has liver disease. the child is taking the blood thinning drug warfarin. 	<p>Drug Facts (continued)</p> <p>Warnings</p> <p>Liver warnings: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount.</p> <p>Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin redness hives skin peeling <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>For rectal use only.</p> <p>Do not use</p> <ul style="list-style-type: none"> if you are allergic to acetaminophen. with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> the child has liver disease. the child is taking the blood thinning drug warfarin.
---	--

<p>Drug Facts (continued)</p> <p>Warnings</p> <p>Liver warnings: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount.</p> <p>Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin redness hives skin peeling <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>For rectal use only.</p> <p>Do not use</p> <ul style="list-style-type: none"> if you are allergic to acetaminophen. with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> the child has liver disease. the child is taking the blood thinning drug warfarin. 	<p>Drug Facts (continued)</p> <p>Warnings</p> <p>Liver warnings: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount.</p> <p>Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin redness hives skin peeling <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>For rectal use only.</p> <p>Do not use</p> <ul style="list-style-type: none"> if you are allergic to acetaminophen. with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> the child has liver disease. the child is taking the blood thinning drug warfarin.
--	--

<p>Drug Facts</p> <p>Active ingredient (in each rectal suppository)</p> <p>Acetaminophen, USP 120 mg Pain reliever/fever reducer</p> <p>Uses</p> <p>temporarily</p> <ul style="list-style-type: none"> reduces fever relieves minor aches, pains, and headache 	<p>Drug Facts (continued)</p> <p>Warnings</p> <p>Liver warnings: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount.</p> <p>Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin redness hives skin peeling <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>For rectal use only.</p> <p>Do not use</p> <ul style="list-style-type: none"> if you are allergic to acetaminophen. with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> the child has liver disease. the child is taking the blood thinning drug warfarin.
---	--

NDC 0472-1201-06

Children's Ages 3-6 years **120 mg**



FeverAll[®]

Acetaminophen Suppositories

Pain reliever/fever reducer

6 Rectal Suppositories **120 mg each**



M 047211201006 8

Other information • store at 2-27°C (35°-80°F)

do not use if imprinted suppository wrapper is opened or damaged

Inactive ingredients

glycerol monoacetate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80

Questions? 1-800-432-6534 between 9 am and 4 pm EST, Monday-Friday.

12010613M1 VC101919

Actavis

Made in USA

20092 USA

LLC, Lincoln, NC

© Actavis Inc.

Do not use if imprinted wrapper is opened or damaged.

Suppositories are individually wrapped.

(b) (4)



Drug Facts (continued)

Do not use

- in children under 6 years
- if you are allergic to acetaminophen.
- with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease.
- you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- fever lasts more than 3 days (72 hours), or recurs.
- pain gets worse or lasts more than 10 days.
- new symptoms occur.
- redness or swelling is present in the painful area.

These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- carefully insert suppository well up into the rectum
- remove wrapper

Questions? 1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.





Drug Facts

Active ingredient (in each rectal suppository) Acetaminophen, USP 325 mg

Purposes Pain reliever/fever reducer

Uses temporarily • reduces fever • relieves minor aches, pains, and headache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if • a child 6 to 12 years takes more than 5 doses in 24 hours • an adult or child 12 years and older takes more than 6 doses in 24 hours, which is the maximum daily amount • taken with other drugs containing acetaminophen • an adult takes 3 or more alcoholic drinks everyday while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

For rectal use only.





Drug Facts (continued)

Other Information

- store at 2°-27°C (35°-80°F)
- do not use if imprinted suppository wrapper is opened or damaged

Inactive Ingredients glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80





Drug Facts (continued)

Dosing Chart

Age	Dose
under 6 years	do not use
6 to 12 years	Use 1 suppository every 4 to 6 hours. (maximum of 5 doses in 24 hours)
adults and children 12 years and older	Use 2 suppositories every 4 to 6 hours. (maximum of 6 doses)





actavis

Dist. by Actavis Mid Atlantic LLC
 Lincolnton, NC 28092 USA Made in USA

**Tamper-evident: Suppositories are individually wrapped.
 Do not use if imprinted wrapper is opened or damaged.**



12020813Q1 VC101922
 3 04721120250 8



Jr. Strength

Ages 6-12 years

NOT FOR RETAIL SALE
 NDC 0472-1202-50



FeverAll®
 Acetaminophen Suppositories
 Pain reliever/fever reducer

50 Rectal Suppositories

325 mg each





FeverAll®
 Acetaminophen Suppositories
 Pain reliever/fever reducer

325 mg each



(b) (4)

Drug Facts (continued)

Do not use

- in children under 12 years.
- if you are allergic to acetaminophen.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease.
- you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- fever lasts more than 3 days (72 hours), or recurs.
- pain gets worse or lasts more than 10 days.
- new symptoms occur.
- redness or swelling is present in the painful area.

These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not use more than directed
- remove wrapper
- carefully insert suppository well up into the rectum
- adults and children 12 years and older:
 - 1 suppository every 4 to 6 hours while symptoms last
- children under 12 years: ask a doctor

Other information • store at 2°-27°C (35°-80°F)

- do not use if imprinted suppository wrapper is opened or damaged

Inactive ingredients glycerol monostearate, polyoxethylene stearate, polysorbate 80

Questions? 1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.

Drug Facts

Active ingredient (in each rectal suppository) Acetaminophen, USP 650 mg

Purposes Pain reliever/fever reducer

Uses temporarily • reduces fever • relieves minor aches, pains, and headache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if an adult or child 12 years and older takes more than 6 doses in 24 hours, which is the maximum daily amount • taken with other drugs containing acetaminophen • an adult takes 3 or more alcoholic drinks everyday while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

For racial use only.

Drug Facts (continued)

Directions

- do not use more than directed
- remove wrapper
- carefully insert suppository well up into the rectum
- adults and children 12 years and older:
 - 1 suppository every 4 to 6 hours while symptoms last
- children under 12 years: ask a doctor

Other information • store at 2°-27°C (35°-80°F)

- do not use if imprinted suppository wrapper is opened or damaged

Inactive ingredients glycerol monostearate, polyoxethylene stearate, polysorbate 80

Questions? 1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.

Drug Facts (continued)

Do not use

- in children under 12 years.
- if you are allergic to acetaminophen.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease.
- you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- fever lasts more than 3 days (72 hours), or recurs.
- pain gets worse or lasts more than 10 days.
- new symptoms occur.
- redness or swelling is present in the painful area.

These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.



FeverAll
Acetaminophen Suppositories
Pain reliever/fever reducer
650 mg each

Adults' Ages 12 years and older

FeverAll[®]

Acetaminophen Suppositories

Pain reliever/fever reducer

50 Rectal Suppositories **650 mg each**

NOT FOR RETAIL SALE
NDC 0472-1203-50



Dist. by Actavis Mid Atlantic LLC
Lincolnton, NC 28092 USA Made in USA

Tamper-evident: Suppositories are individually wrapped. Do not use if imprinted wrapper is opened or damaged.

12030813J1 VC101923



3 04721 20350 5

(b) (4)