

**ORIGINAL
BLISTER PACK**

NDC 0135-0474-18

Nicorette[®]

nicotine polacrilex gum, 2mg • stop smoking aid

Gum

**200 PIECES
2mg EACH**



**2
mg**

**FOR THOSE WHO SMOKE
THEIR FIRST CIGARETTE
MORE THAN 30 MINUTES
AFTER WAKING UP.**

If you smoke your first cigarette
WITHIN 30 MINUTES of waking
up, use Nicorette 4mg Gum

**Coated
FOR BOLD FLAVOR**

White Ice Mint[®]

104154XA



200 PIECES, 2mg EACH (2-100 piece cartons)

Nicorette[®]

nicotine polacrilex gum, 2mg • stop smoking aid

Gum

How To Open the Nicorette[®] Gum Blister Pack



To remove the gum, tear off single unit.



Peel off backing, starting at corner with loose edge.



Push gum through foil.

Personalized Quit Plan



Visit www.nicorette.com to enroll in a free, personalized quit plan. **Nicorette Committed Quitters[®]** is a personalized stop smoking program that will help you understand your smoking habits and determine how to best overcome your cravings. Track, target, and tame your temptations with **Nicorette Committed Quitters[®]**. Only available at www.nicorette.com

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicorette[®] gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicorette[®] gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

Drug Facts

Active ingredient (in each chewing piece) **Purpose**
Nicotine polacrilex (equal to 2mg nicotine) Stop smoking aid

Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information

Drug Facts (continued)

- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- each piece contains: calcium 94mg, sodium 11mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients

acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, starch, sucralose, titanium dioxide, xylitol

Questions or comments? call toll-free 1-800-419-4766 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)

104029XA

Lot:
Exp:

Distributed by
GlaxoSmithKline Consumer Healthcare, L.P.
Moon Township, PA 15108 Made in Sweden
©2014 GSK

NICORETTE, NICORETTE COMMITTED QUITTERS, WHITE ICE MINT and the NICORETTE sunburst design are registered trademarks of the GSK group of companies.

851982



3 0766-7750-31 7



Datamatrix Code
104029XA
5mm

NO COPY THIS AREA

THIS AREA TO BE
DIECUT OUT AND
DISCARDED

Helvetica Neue 67 Medium Condensed 6pt

Helvetica Neue 67 Medium Condensed 6pt

© Helvetica Neue 55 Roman 6pt

©2014 GSK 104040XA



Data Matrix
104040XA

**ORIGINAL
BLISTER PACK**

NDC 0135-241-17

Nicorette[®]

nicotine polacrilex gum, 2mg • stop smoking aid

Gum

200 PIECES
2mg EACH



2
mg

**FOR THOSE WHO SMOKE
THEIR FIRST CIGARETTE
MORE THAN 30 MINUTES
AFTER WAKING UP.**

If you smoke your first cigarette
WITHIN 30 MINUTES of waking
up, use Nicorette 4mg Gum

Coated
FOR BOLD FLAVOR

Fruit Chill[™]

104156XA



200 PIECES, 2mg EACH (2-100 piece cartons)

Nicorette[®]

nicotine polacrilex gum, 2mg • stop smoking aid

Gum

How To Open the Nicorette[®] Gum Blister Pack



To remove the gum, tear off single unit.



Peel off backing, starting at corner with loose edge.



Push gum through foil.

Personalized Quit Plan



Visit www.nicorette.com to enroll in a free, personalized quit plan. **Nicorette Committed Quitters[®]** is a personalized stop smoking program that will help you understand your smoking habits and determine how to best overcome your cravings. Track, target, and tame your temptations with **Nicorette Committed Quitters[®]**. Only available at www.nicorette.com

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicorette[®] gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicorette[®] gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

Drug Facts

Active ingredient (in each chewing piece) Purpose
Nicotine polacrilex (equal to 2mg nicotine) Stop smoking aid

Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information

Drug Facts (continued)

- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- each piece contains: calcium 94mg, sodium 11mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium, carnauba wax, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium bicarbonate, sodium carbonate, sucralose, titanium dioxide, xylitol

Questions or comments? call toll-free 1-800-419-4766 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)


104021XA

Lot:
Exp:

Distributed by
GlaxoSmithKline Consumer Healthcare, L.P.
Moon Township, PA 15108 Made in Sweden
©2014 GSK

NICORETTE, NICORETTE FRUIT CHILL, NICORETTE COMMITTED QUITTERS and the NICORETTE sunburst design are registered trademarks of the GSK group of companies.

851985



3 0766-7867-53 4



NO COPY THIS AREA

THIS AREA TO BE
DIECUT OUT AND
DISCARDED

Helvetica Neue 67 Medium Condensed 6pt

Helvetica Neue 67 Medium Condensed 6pt

© Helvetica Neue 55 Roman 6pt

©2014 GSK 104020XA



Data Matrix
104020XA

**ORIGINAL
BLISTER PACK**

NDC 0135-0225-17

Nicorette[®]

nicotine polacrilex gum, 2mg • stop smoking aid

Gum

200 PIECES
2mg EACH



**FOR THOSE WHO SMOKE
THEIR FIRST CIGARETTE
MORE THAN 30 MINUTES
AFTER WAKING UP.**

If you smoke your first cigarette
WITHIN 30 MINUTES of waking
up, use Nicorette 4mg Gum

Coated
FOR BOLD FLAVOR

Fresh Mint[™]

104158XA



200 PIECES, 2mg EACH (2-100 piece cartons)

Nicorette[®]

nicotine polacrilex gum, 2mg • stop smoking aid

Gum

How To Open the Nicorette[®] Gum Blister Pack



To remove the gum, tear off single unit.



Peel off backing, starting at corner with loose edge.



Push gum through foil.

Personalized Quit Plan



Visit www.nicorette.com to enroll in a free, personalized quit plan. **Nicorette Committed Quitters[®]** is a personalized stop smoking program that will help you understand your smoking habits and determine how to best overcome your cravings. Track, target, and tame your temptations with **Nicorette Committed Quitters[®]**. Only available at www.nicorette.com

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew **at least 9 pieces** of Nicorette[®] gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicorette[®] gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

Drug Facts

Active ingredient (in each chewing piece) **Purpose**
Nicotine polacrilex (equal to 2mg nicotine) Stop smoking aid

Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information

Drug Facts (continued)

- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- **each piece contains:** calcium 94mg, sodium 11mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium, carnauba wax, edible ink, gum base, magnesium oxide, menthol, peppermint oil, sodium bicarbonate, sodium carbonate, titanium dioxide, xylitol

Questions or comments? call toll-free 1-800-419-4766 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)

104025XA

Lot:
Exp:

Distributed by
GlaxoSmithKline Consumer Healthcare, L.P.
Moon Township, PA 15108 Made in Sweden
©2014 GSK

NICORETTE, NICORETTE COMMITTED QUITTERS and the NICORETTE sunburst design are registered trademarks of the GSK group of companies.



NO COPY THIS AREA

THIS AREA TO BE
DIECUT OUT AND
DISCARDED

©2014 GSK 104036XA



Data Matrix
104036XA

Helvetica Neue 67 Medium Condensed 6pt

Helvetica Neue 67 Medium Condensed 6pt

©) Helvetica Neue 55 Roman 6pt

**ORIGINAL
BLISTER PACK**

NDC 0135-0475-18

Nicorette[®]

nicotine polacrilex gum, 4mg • stop smoking aid

Gum

**200 PIECES
4mg EACH**



**4
mg**

**FOR THOSE WHO SMOKE
THEIR FIRST CIGARETTE
WITHIN 30 MINUTES
OF WAKING UP.**

If you smoke your first
cigarette **MORE THAN
30 MINUTES** after waking
up, use Nicorette 2mg Gum

**Coated
FOR BOLD FLAVOR**

White Ice Mint[®]

104156XA



200 PIECES, 4mg EACH (2-100 piece cartons)

Nicorette[®]

nicotine polacrilex gum, 4mg • stop smoking aid

Gum

How To Open the Nicorette[®] Gum Blister Pack



To remove the gum, tear off single unit.



Peel off backing, starting at corner with loose edge.



Push gum through foil.

Personalized Quit Plan



Visit www.nicorette.com to enroll in a free, personalized quit plan. **Nicorette Committed Quitters[®]** is a personalized stop smoking program that will help you understand your smoking habits and determine how to best overcome your cravings. Track, target, and tame your temptations with **Nicorette Committed Quitters[®]**. Only available at www.nicorette.com

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicorette[®] gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicorette[®] gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

Drug Facts

Active ingredient (in each chewing piece) **Purpose**
Nicotine polacrilex (equal to 4mg nicotine) Stop smoking aid

Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information

Drug Facts (continued)

- begin using the gum on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine gum
- if you smoke your first cigarette within 30 minutes of waking up, use 4mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- each piece contains: calcium 94mg, sodium 13mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients

acesulfame potassium, carnauba wax, D&C yellow #10 AI, lake, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium carbonate, starch, sucralose, titanium dioxide, xylitol

Questions or comments? call toll-free 1-800-419-4766 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)


104041XA

Lot:
Exp:

Distributed by
GlaxoSmithKline Consumer Healthcare, L.P.
Moon Township, PA 15108 Made in Sweden
©2014 GSK

NICORETTE, NICORETTE COMMITTED QUITTERS, WHITE ICE MINT and the NICORETTE sunburst design are registered trademarks of the GSK group of companies.

851991



3 0766-7760-31 6



NO COPY THIS AREA

THIS AREA TO BE
DIECUT OUT AND
DISCARDED

Helvetica Neue 67 Medium Condensed 6pt

Helvetica Neue 67 Medium Condensed 6pt

© Helvetica Neue 55 Roman 6pt

©2014 GSK 104040XA



Data Matrix
104040XA

**ORIGINAL
BLISTER PACK**

NDC 0135-242-17

Nicorette[®]

nicotine polacrilex gum, 4mg • stop smoking aid

Gum

**200 PIECES
4mg EACH**



**4
mg**

**FOR THOSE WHO SMOKE
THEIR FIRST CIGARETTE
WITHIN 30 MINUTES
OF WAKING UP.**

If you smoke your first
cigarette **MORE THAN
30 MINUTES** after waking
up, use Nicorette 2mg Gum

**Coated
FOR BOLD FLAVOR**

Fruit Chill[™]

104157XA



Datamatrix Code
104157XA
5mm

200 PIECES, 4mg EACH (2-100 piece cartons)

Nicorette[®]

nicotine polacrilex gum, 4mg • stop smoking aid

Gum

How To Open the Nicorette[®] Gum Blister Pack



To remove the gum, tear off single unit.



Peel off backing, starting at corner with loose edge.



Push gum through foil.

Personalized Quit Plan



Visit www.nicorette.com to enroll in a free, personalized quit plan. **Nicorette Committed Quitters[®]** is a personalized stop smoking program that will help you understand your smoking habits and determine how to best overcome your cravings. Track, target, and tame your temptations with **Nicorette Committed Quitters[®]**. Only available at www.nicorette.com

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicorette[®] gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicorette[®] gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

Drug Facts

Active ingredient (in each chewing piece) **Purpose**
Nicotine polacrilex (equal to 4mg nicotine) Stop smoking aid

Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information

Drug Facts (continued)

- begin using the gum on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine gum
- if you smoke your first cigarette within 30 minutes of waking up, use 4mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- each piece contains: calcium 94mg, sodium 13mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium, carnauba wax, D&C yellow #10 Al. lake, edible ink, flavor, gum base, hypromellose, magnesium oxide, menthol, peppermint oil, polysorbate 80, sodium carbonate, sucralose, titanium dioxide, xylitol

Questions or comments? call toll-free 1-800-419-4766 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)


104033XA

Lot:
Exp:

Distributed by
GlaxoSmithKline Consumer Healthcare, L.P.
Moon Township, PA 15108 Made in Sweden
©2014 GSK

NICORETTE, NICORETTE FRUIT CHILL, NICORETTE COMMITTED QUITTERS and the NICORETTE sunburst design are registered trademarks of the GSK group of companies.

851995



3 0766-7867-63 3



NO COPY THIS AREA

THIS AREA TO BE
DIECUT OUT AND
DISCARDED

Helvetica Neue 67 Medium Condensed 6pt

Helvetica Neue 67 Medium Condensed 6pt

© Helvetica Neue 55 Roman 6pt

©2014 GSK 104020XA



Data Matrix
104020XA

**ORIGINAL
BLISTER PACK**

NDC 0135-0226-17

Nicorette[®]

nicotine polacrilex gum, 4mg • stop smoking aid

Gum

**200 PIECES
4mg EACH**



**4
mg**

FOR THOSE WHO SMOKE
THEIR FIRST CIGARETTE
WITHIN 30 MINUTES
OF WAKING UP.

If you smoke your first
cigarette **MORE THAN
30 MINUTES** after waking
up, use Nicorette 2mg Gum

**Coated
FOR BOLD FLAVOR**

Fresh Mint[™]

104159XA



200 PIECES, 4mg EACH (2-100 piece cartons)

Nicorette[®]

nicotine polacrilex gum, 4mg • stop smoking aid

Gum

How To Open the Nicorette[®] Gum Blister Pack



To remove the gum, tear off single unit.



Peel off backing, starting at corner with loose edge.



Push gum through foil.

Personalized Quit Plan



Visit www.nicorette.com to enroll in a free, personalized quit plan. **Nicorette Committed Quitters[®]** is a personalized stop smoking program that will help you understand your smoking habits and determine how to best overcome your cravings. Track, target, and tame your temptations with **Nicorette Committed Quitters[®]**. Only available at www.nicorette.com

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicorette[®] gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicorette[®] gum for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

Drug Facts

Active ingredient **Purpose**
(in each chewing piece)
Nicotine polacrilex (equal to 4mg nicotine) Stop smoking aid

Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information

Drug Facts (continued)

- begin using the gum on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine gum
- if you smoke your first cigarette within 30 minutes of waking up, use 4mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- each piece contains: calcium 94mg, sodium 13mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium, carnauba wax, D&C yellow #10 Al. lake, edible ink, gum base, magnesium oxide, menthol, peppermint oil, sodium carbonate, titanium dioxide, xylitol

Questions or comments? call toll-free 1-800-419-4766 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)

104037XA

Lot:
Exp:

Distributed by
GlaxoSmithKline Consumer Healthcare, L.P.
Moon Township, PA 15108 Made in Sweden
©2014 GSK

NICORETTE, NICORETTE COMMITTED QUITTERS and the NICORETTE sunburst design are registered trademarks of the GSK group of companies.



NO COPY THIS AREA

THIS AREA TO BE
DIECUT OUT AND
DISCARDED

©2014 GSK 104036XA



Data Matrix
104036XA

Helvetica Neue 67 Medium Condensed 6pt

Helvetica Neue 67 Medium Condensed 6pt

©) Helvetica Neue 55 Roman 6pt

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
10/30/2014