

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use MIACALCIN nasal spray safely and effectively. See full prescribing information for MIACALCIN nasal spray.

**MIACALCIN® (calcitonin-salmon) nasal spray, for intranasal use**  
**Initial U.S. Approval: 1975**

-----**RECENT MAJOR CHANGES**-----

Indications and Usage (1.2)	03/2014
Warnings and Precautions (5.4)	03/2014

-----**INDICATIONS AND USAGE**-----

Miacalcin nasal spray is a calcitonin, indicated for the treatment of postmenopausal osteoporosis in women greater than 5 years postmenopause when alternative treatments are not suitable. Fracture reduction efficacy has not been demonstrated (1.1).

Limitations of Use:

- Due to the possible association between malignancy and calcitonin-salmon use, the need for continued therapy should be re-evaluated on a periodic basis (1.2, 5.4)
- Miacalcin nasal spray has not been shown to increase bone mineral density in early postmenopausal women (1.2)

-----**DOSAGE AND ADMINISTRATION**-----

- For intranasal use only: one spray (200 International Units) per day, alternating nostrils daily (2.1)
- Prior to first use, allow the bottle to reach room temperature and prime the pump (2.2)
- Ensure adequate calcium and vitamin D intake (2.3)

-----**DOSAGE FORMS AND STRENGTHS**-----

Nasal Spray: 2200 International Units per mL of calcitonin-salmon in a 3.7 mL fill glass bottle with screw-on pump. Each actuation delivers 200 International Units of calcitonin-salmon (3)

-----**CONTRAINDICATIONS**-----

Hypersensitivity to calcitonin-salmon or any of the excipients (4)

-----**WARNINGS AND PRECAUTIONS**-----

- Serious hypersensitivity reactions including anaphylactic shock have been reported. Consider skin testing prior to treatment in patients with suspected hypersensitivity to calcitonin-salmon (5.1)
- Hypocalcemia has been reported. Ensure adequate intake of calcium and vitamin D (5.2)
- Nasal adverse reactions, including severe ulceration can occur. Periodic nasal examinations are recommended (5.3)
- Malignancy: A meta-analysis of 21 clinical trials suggests an increased risk of overall malignancies in calcitonin-salmon-treated patients (5.4, 6.1)
- Circulating antibodies to calcitonin-salmon may develop, and may cause loss of response to treatment (5.5)

-----**ADVERSE REACTIONS**-----

Most common adverse reactions (3% or greater) are rhinitis, epistaxis and other nasal symptoms, back pain, arthralgia, and headache (6)

**To report SUSPECTED ADVERSE REACTIONS, contact Novartis Pharmaceuticals Corporation at 1-888-669-6682 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

-----**DRUG INTERACTIONS**-----

- Concomitant use of calcitonin-salmon and lithium may lead to a reduction in plasma lithium concentrations due to increased urinary clearance of lithium. The dose of lithium may require adjustment (7)

-----**USE IN SPECIFIC POPULATIONS**-----

- There are no data to support use in children (8.4)
- Nasal reactions are more common in elderly patients (8.5)

**See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.**

**Revised: 03/2014**

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

- 1.1 Treatment of Postmenopausal Osteoporosis
- 1.2 Important Limitations of Use

**2 DOSAGE AND ADMINISTRATION**

- 2.1 Basic Dosing Information
- 2.2 Priming (Activation) of Pump
- 2.3 Recommendations for Calcium and Vitamin D Supplementation

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

- 5.1 Hypersensitivity Reactions
- 5.2 Hypocalcemia
- 5.3 Nasal Adverse Reactions
- 5.4 Malignancy
- 5.5 Antibody Formation
- 5.6 Urine Sediment Abnormalities

**6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience
- 6.3 Immunogenicity

**7 DRUG INTERACTIONS**

**8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

**10 OVERDOSAGE**

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**14 CLINICAL STUDIES**

**16 HOW SUPPLIED/STORAGE AND HANDLING**

**17 PATIENT COUNSELING INFORMATION**

\*Sections or subsections omitted from the full prescribing information are not listed.

---

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

#### 1.1 Treatment of Postmenopausal Osteoporosis

Miacalcin nasal spray is indicated for the treatment of postmenopausal osteoporosis in women greater than 5 years postmenopause. Fracture reduction efficacy has not been demonstrated. Miacalcin nasal spray should be reserved for patients for whom alternative treatments are not suitable (e.g., patients for whom other therapies are contraindicated or for patients who are intolerant or unwilling to use other therapies).

#### 1.2 Important Limitations of Use

- Due to the possible association between malignancy and calcitonin-salmon use, the need for continued therapy should be re-evaluated on a periodic basis [see *Warnings and Precautions* (5.4)].
- Miacalcin nasal spray has not been shown to increase spinal bone mineral density in early postmenopausal women.

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Basic Dosing Information

The recommended dose of Miacalcin nasal spray is 1 spray (200 International Units) per day administered intranasally, alternating nostrils daily.

#### 2.2 Priming (Activation) of Pump

Unopened Miacalcin nasal spray should be stored in the refrigerator. Before using the first dose of Miacalcin nasal spray, the patient should wait until it has reached room temperature. To prime the pump before it is used for the first time, the bottle should be held upright and the two white side arms of the pump depressed toward the bottle, repeat until a full spray is released. The pump is primed once the first full spray is emitted. To administer, the nozzle should first be carefully placed into the nostril while the patient's head is in the upright position, then the pump should be firmly depressed toward the bottle. The pump should not be primed before each daily dose.

#### 2.3 Recommendations for Calcium and Vitamin D Supplementation

Patients who use Miacalcin nasal spray should receive adequate calcium (at least 1000 mg elemental calcium per day) and vitamin D (at least 400 International Units per day).

### 3 DOSAGE FORMS AND STRENGTHS

Miacalcin nasal spray consists of one glass bottle and one screw-on pump. The bottle contains 3.7 mL of calcitonin-salmon clear solution at a concentration of 2200 International Units per mL. A primed pump delivers 0.09 mL (200 International Units) calcitonin-salmon per actuation.

### 4 CONTRAINDICATIONS

Hypersensitivity to calcitonin-salmon or any of the excipients. Reactions have included anaphylactic shock, anaphylaxis, bronchospasm, and swelling of the tongue or throat[see *Warnings and Precautions* (5.1)].

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Hypersensitivity Reactions

Serious hypersensitivity reactions have been reported in patients receiving Miacalcin nasal spray, e.g., bronchospasm, swelling of the tongue or throat, anaphylaxis and anaphylactic shock. Reports of serious

hypersensitivity reactions with injectable calcitonin-salmon have also been reported, including reports of death attributed to anaphylaxis. The usual provisions should be made for emergency treatment if such a reaction occurs. Hypersensitivity reactions should be differentiated from generalized flushing and hypotension [*see Contraindications (4)*].

For patients with suspected hypersensitivity to calcitonin-salmon, skin testing should be considered prior to treatment utilizing a dilute, sterile solution of a calcitonin-salmon injectable product. Healthcare providers may wish to refer patients who require skin testing to an allergist. A detailed skin testing protocol is available from the Medical Services Department of Novartis Pharmaceuticals Corporation.

## 5.2 Hypocalcemia

Hypocalcemia associated with tetany (i.e. muscle cramps, twitching) and seizure activity has been reported with calcitonin therapy. Hypocalcemia must be corrected before initiating therapy with Miacalcin nasal spray. Other disorders affecting mineral metabolism (such as vitamin D deficiency) should also be effectively treated. In patients with these conditions, serum calcium and symptoms of hypocalcemia should be monitored during therapy with Miacalcin nasal spray. Use of Miacalcin nasal spray is recommended in conjunction with an adequate intake of calcium and vitamin D [*see Dosage and Administration (2.3)*].

## 5.3 Nasal Adverse Reactions

Adverse reactions related to the nose including rhinitis and epistaxis have been reported. Development of mucosal alterations may occur. Therefore, periodic nasal examinations with visualization of the nasal mucosa, turbinates, septum and mucosal blood vessels are recommended prior to start of treatment with Miacalcin nasal spray, periodically during the course of therapy, and at any time nasal symptoms occur.

Miacalcin nasal spray should be discontinued if severe ulceration of the nasal mucosa occurs, as indicated by ulcers greater than 1.5 mm in diameter or penetrating below the mucosa, or those associated with heavy bleeding. Although smaller ulcers often heal without withdrawal of Miacalcin nasal spray, medication should be discontinued temporarily until healing occurs [*see Adverse Reactions (6.1)*].

## 5.4 Malignancy

In a meta-analysis of 21 randomized, controlled clinical trials with calcitonin-salmon (nasal spray or investigational oral formulations), the overall incidence of malignancies reported was higher among calcitonin-salmon-treated patients (4.1%) compared with placebo-treated patients (2.9%). This suggests an increased risk of malignancies in calcitonin-salmon-treated patients compared to placebo-treated patients. The benefits for the individual patient should be carefully considered against possible risks [*see Adverse Reactions (6.1)*].

## 5.5 Antibody Formation

Circulating antibodies to calcitonin-salmon have been reported with Miacalcin nasal spray. The possibility of antibody formation should be considered in any patient with an initial response to Miacalcin nasal spray who later stops responding to treatment [*see Adverse Reactions (6.3)*].

## 5.6 Urine Sediment Abnormalities

Coarse granular casts and casts containing renal tubular epithelial cells were reported in young adult volunteers at bed rest who were given injectable calcitonin-salmon to study the effect of immobilization on osteoporosis. There was no other evidence of renal abnormality and the urine sediment normalized after calcitonin-salmon was stopped. Periodic examinations of urine sediment should be considered. Urine sediment abnormalities have not been reported in ambulatory volunteers treated with calcitonin-salmon nasal spray.

## 6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in greater detail in other sections of the label:

- Hypersensitivity Reactions, including anaphylaxis [see *Warnings and Precautions (5.1)*]
- Hypocalcemia [see *Warnings and Precautions (5.2)*]
- Nasal Adverse Reactions [see *Warnings and Precautions (5.3)*]
- Malignancy [see *Warnings and Precautions (5.4)*]

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of Miacalcin (calcitonin-salmon) nasal spray in the treatment of postmenopausal osteoporosis was assessed in 5 randomized, double-blind, placebo controlled trials that enrolled postmenopausal women, aged 45-75 years. The duration of the trials ranged from 1 to 2 years. The incidence of adverse reactions reported in studies involving postmenopausal osteoporotic patients chronically exposed to Miacalcin nasal spray (N=341) and to placebo nasal spray (N=131), and reported in greater than 3% of Miacalcin treated patients are presented in the following table. Other than flushing, nausea, possible allergic reactions, and possible local irritative effects in the respiratory tract, a relationship to Miacalcin nasal spray has not been established.

**Table 1: Adverse Reactions Occurring in at Least 3% of Postmenopausal Patients Treated with Miacalcin Nasal Spray**

	<b>Miacalcin Nasal Spray</b>	<b>Placebo Nasal Spray</b>
<b>Adverse Reaction</b>	<b>N=341 % of Patients</b>	<b>N=131 % of Patients</b>
Rhinitis	12	7
Symptom of Nose†	11	16
Back Pain	5	2
Arthralgia	4	5
Epistaxis	4	5
Headache	3	5

†Symptom of nose includes: nasal crusts, dryness, redness or erythema, nasal sores, irritation, itching, thick feeling, soreness, pallor, infection, stenosis, runny/blocked, small wound, bleeding wound, tenderness, uncomfortable feeling and sore across bridge of nose.

Nasal Adverse Reactions: In all postmenopausal patients treated with Miacalcin nasal spray, the most commonly reported nasal adverse reactions included rhinitis (12%), epistaxis (4%), and sinusitis (2%). Smoking did not have a contributory effect on the occurrence of nasal adverse reactions.

Adverse reactions reported in 1-3% of patients treated with Miacalcin nasal spray include: influenza-like symptoms, erythematous rash, arthrosis, myalgia, sinusitis, upper respiratory tract infection, bronchospasm, abdominal pain, nausea, dizziness, paresthesia, abnormal lacrimation, conjunctivitis, lymphadenopathy, infection, and depression.

## **Malignancy**

A meta-analysis of 21 randomized, controlled clinical trials with calcitonin-salmon (nasal spray or investigational oral formulations) was conducted to assess the risk of malignancies in calcitonin-salmon-treated patients compared to placebo-treated patients. The trials in the meta-analysis ranged in duration from 6 months to 5 years and included a total of 10883 patients (6151 treated with calcitonin-salmon and 4732 treated with placebo). The overall incidence of malignancies reported in these 21 trials was higher among calcitonin-salmon-treated patients (254/6151 or 4.1%) compared with placebo-treated patients (137/4732 or 2.9%). Findings were similar when analyses were restricted to the 18 nasal spray only trials [calcitonin-salmon 122/2712 (4.5%); placebo 30/1309 (2.3%)].

The meta-analysis results suggest an increased risk of overall malignancies in calcitonin-salmon-treated patients compared to placebo-treated patients when all 21 trials are included and when the analysis is restricted to the 18 nasal spray only trials (see Table 2). It is not possible to exclude an increased risk when calcitonin-salmon is administered by the subcutaneous, intramuscular, or intravenous route because these routes of administration were not investigated in the meta-analysis. The increased malignancy risk seen with the meta-analysis was heavily influenced by a single large 5-year trial, which had an observed risk difference of 3.4% [95% CI (0.4%, 6.5%)]. Imbalances in risks were still observed when analyses excluded basal cell carcinoma (see Table 2); the data were not sufficient for further analyses by type of malignancy. A mechanism for these observations has not been identified. Although a definitive causal relationship between calcitonin-salmon use and malignancies cannot be established from this meta-analysis, the benefits for the individual patient should be carefully evaluated against all possible risks [see *Warnings and Precautions (5.4)*].

**Table 2: Risk Difference for Malignancies in Calcitonin-Salmon-Treated Patients Compared with Placebo-Treated Patients**

<b>Patients</b>	<b>Malignancies</b>	<b>Risk Difference<sup>1</sup> (%)</b>	<b>95% Confidence Interval<sup>2</sup> (%)</b>
All (nasal spray + oral)	All	1.0	(0.3, 1.6)
All (nasal spray + oral)	Excluding basal cell carcinoma	0.5	(-0.1, 1.2)
All (nasal spray only)	All	1.4	(0.3, 2.6)
All (nasal spray only)	Excluding basal cell carcinoma	0.8	(-0.2, 1.8)

<sup>1</sup> The overall adjusted risk difference is the difference between the percentage of patients who had any malignancy (or malignancy excluding basal cell carcinoma) in calcitonin-salmon and placebo treatment groups, using the Mantel-Haenszel (MH) fixed-effect method. A risk difference of 0 is suggestive of no difference in malignancy risks between the treatment groups.

<sup>2</sup> The corresponding 95% confidence interval for the overall adjusted risk difference also based on MH fixed-effect method.

## **6.2 Postmarketing Experience**

Because postmarketing adverse reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The following adverse reactions have been reported during post-approval use of Miacalcin nasal spray.

***Allergic / Hypersensitivity Reactions:*** Serious allergic reactions have been reported in patients receiving calcitonin-salmon nasal spray, including anaphylaxis and anaphylactic shock.

***Hypocalcemia:*** Hypocalcemia with paresthesia has been reported.

***Body as a whole:*** facial or peripheral edema

***Cardiovascular:*** hypertension, vasodilatation, syncope, chest pain

***Nervous system:*** dizziness, seizure, visual or hearing impairment, tinnitus

***Respiratory/ Special Senses:*** cough, bronchospasm, dyspnea, loss of taste/smell

***Skin:*** rash/dermatitis, pruritus, alopecia, increased sweating

***Gastrointestinal:*** diarrhea

***Nervous system disorders:*** tremor

### **6.3 Immunogenicity**

Consistent with the potentially immunogenic properties of medicinal products containing peptides, administration of Miacalcin may trigger the development of anti-calcitonin antibodies. In a two-year Miacalcin nasal spray clinical study that evaluated immunogenicity, a measurable antibody titer was found in 69% of patients treated with Miacalcin and 3% of placebo-treated patients. Antibody formation may be associated with a loss of response to treatment [see *Warnings and Precautions (5.5)*].

The incidence of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of a positive antibody test result may be influenced by several factors, including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of antibodies to Miacalcin nasal spray with the incidence of antibodies to other calcitonin-containing products may be misleading.

## **7 DRUG INTERACTIONS**

No formal drug interaction studies have been performed with Miacalcin nasal spray.

Concomitant use of calcitonin-salmon and lithium may lead to a reduction in plasma lithium concentrations due to increased urinary clearance of lithium. The dose of lithium may require adjustment.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

Pregnancy Category C:

#### **Risk Summary**

There are no adequate and well-controlled studies in pregnant women. Miacalcin nasal spray should be used during pregnancy only if the potential benefit justifies the use as compared with potential risks to the patient and fetus. Based on animal data, Miacalcin is predicted to have low probability of increasing the risk of adverse developmental outcomes above background risk.

#### **Animal Data**

Calcitonin-salmon has been shown to cause a decrease in fetal birth weights in rabbits when given by subcutaneous injection in doses 4-18 times the parenteral dose (of 54 International Units/m<sup>2</sup>) and 70-278 times the intranasal dose recommended for human use based on body surface area.

No embryo/fetal toxicities related to Miacalcin were reported from maternal subcutaneous daily doses in rats up to 80 International Units/kg/day from gestation day 6 to 15.

### 8.3 Nursing Mothers

It is not known whether this drug is excreted in human milk. No studies have been conducted to assess the impact of Miacalcin on milk production in humans, its presence in human breast milk, or its effects on the breast-fed child. Because many drugs are excreted in human milk, caution should be exercised when Miacalcin is administered to a nursing woman. Calcitonin has been shown to inhibit lactation in rats.

### 8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

### 8.5 Geriatric Use

In a multi-centered, double-blind, randomized clinical study of calcitonin-salmon nasal spray, 279 patients were less than 65 years old, while 467 patients were 65 to 74 years old and 196 patients were 75 years old and older. Compared to subjects less than 65 years old, the incidence of nasal adverse reactions (rhinitis, irritation, erythema, and excoriation) was higher in patients over the age of 65, particularly among those over the age of 75. Other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

## 10 OVERDOSAGE


The pharmacologic actions of Miacalcin nasal spray suggest that hypocalcemic tetany could occur in overdose. Therefore, provisions for parenteral administration of calcium should be available for the treatment of overdose.

Single doses of Miacalcin nasal spray up to 1600 International Units, doses up to 800 International Units per day for 3 days and chronic administration of doses up to 600 International Units per day have been studied without serious adverse effects.

## 11 DESCRIPTION

Calcitonin is a polypeptide hormone secreted by the parafollicular cells of the thyroid gland in mammals and by the ultimobranchial gland of birds and fish.

Miacalcin (calcitonin-salmon) nasal spray is a synthetic polypeptide of 32 amino acids in the same linear sequence that is found in calcitonin of salmon origin. This is shown by the following graphic formula:



It is provided in a 3.7 mL fill glass bottle as a solution for nasal administration. This is sufficient medication for 30 doses.

*Active Ingredient:* calcitonin-salmon 2200 International Units per mL (corresponding to 200 International Units per 0.09 mL actuation).

*Inactive Ingredients:* sodium chloride, benzalkonium chloride, hydrochloric acid (added as necessary to adjust pH) and purified water.

The activity of Miacalcin nasal spray is stated in International Units based on bioassay in comparison with the International Reference Preparation of calcitonin-salmon for Bioassay, distributed by the National Institute of Biological Standards and Control, Holly Hill, London.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Calcitonin-salmon is a calcitonin receptor agonist. Calcitonin-salmon acts primarily on bone, but direct renal effects and actions on the gastrointestinal tract are also recognized. Calcitonin-salmon appears to have actions essentially identical to calcitonins of mammalian origin, but its potency per mg is greater and it has a longer duration of action.

The actions of calcitonin on bone and its role in normal human bone physiology are still not completely elucidated, although calcitonin receptors have been discovered in osteoclasts and osteoblasts.

### 12.2 Pharmacodynamics

The information below, describing the clinical pharmacology of calcitonin, has been derived from studies with *injectable* calcitonin-salmon. The mean bioavailability of calcitonin-salmon nasal spray is approximately 3% of the injectable calcitonin-salmon in healthy subjects and, therefore, the conclusions concerning the clinical pharmacology of this preparation may be different.

#### Bone

Single injections of calcitonin-salmon caused a marked transient inhibition of the ongoing bone resorptive process. With prolonged use, there is a persistent, smaller decrease in the rate of bone resorption. Histologically, this is associated with a decreased number of osteoclasts and an apparent decrease in their resorptive activity.

In healthy adults, who have a relatively low rate of bone resorption, the administration of exogenous calcitonin-salmon results in decreases in serum calcium within the limits of the normal range. In healthy children and in patients whose bone resorption is more rapid, decreases in serum calcium are more pronounced in response to calcitonin-salmon.

#### Kidney

Studies with injectable calcitonin-salmon show increases in the excretion of filtered phosphate, calcium, and sodium by decreasing their tubular reabsorption. Comparable studies have not been conducted with Miacalcin nasal spray.

#### Gastrointestinal Tract

Some evidence from studies with injectable preparations suggests that calcitonin-salmon may have effects on the gastrointestinal tract. Short-term administration of injectable calcitonin-salmon results in marked transient decreases in the volume and acidity of gastric juice and in the volume and the trypsin and amylase content of

pancreatic juice. Whether these effects continue to be elicited after each injection of calcitonin-salmon during chronic therapy has not been investigated. These studies have not been conducted with Miacalcin nasal spray.

### Calcium Homeostasis

In two clinical studies designed to evaluate the pharmacodynamic response to calcitonin-salmon nasal spray, administration of calcitonin-salmon 100-1600 International Units to healthy volunteers resulted in rapid and sustained decreases within the normal range for both total serum calcium and serum ionized calcium. Single doses of calcitonin-salmon greater than 400 International Units did not produce any further biological response to the drug.

## **12.3 Pharmacokinetics**

The bioavailability of Miacalcin nasal spray relative to intramuscular administration in healthy volunteers is between 3 and 5%. Miacalcin nasal spray is absorbed rapidly by the nasal mucosa with a mean  $T_{max}$  of about 13 minutes. The terminal half-life of calcitonin-salmon has been calculated to be around 18 minutes and no evidence of accumulation was observed with multiple dosing.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### **Carcinogenicity**

The incidence of pituitary adenomas was increased in rats after one and two years of subcutaneous exposure to synthetic calcitonin-salmon. The significance of this finding to humans is unknown because pituitary adenomas are very common in rats as they age, the pituitary adenomas did not transform into metastatic tumors, there were no other clear treatment-related neoplasms, and synthetic calcitonin-salmon related neoplasms were not observed in mice after two years of dosing.

#### *Rat findings:*

The only clear neoplastic finding in rats dosed subcutaneously with synthetic calcitonin-salmon was an increase in the incidence of pituitary adenomas in male Fisher 344 rats and female Sprague Dawley rats after one year of dosing and male Sprague Dawley rats dosed for one and two years. In female Sprague Dawley rats, the incidence of pituitary adenomas after two years was high in all treatment groups (between 80% and 92% including the control groups) such that a treatment-related effect could not be distinguished from natural background incidence. The lowest dose in male Sprague Dawley rats that developed an increased incidence of pituitary adenomas after two years of dosing (1.7 International Units/kg/day) is approximately 2 times the maximum recommended intranasal dose in humans (200 International Units/day) based on body surface area conversion between rats and humans and a 20-fold conversion factor to account for decreased clinical exposure via the intranasal route. The findings suggest that calcitonin-salmon reduced the latency period for development of non-functioning pituitary adenomas.

#### *Mouse findings:*

No carcinogenicity potential was evident in male or female mice dosed subcutaneously for two years with synthetic calcitonin-salmon at doses up to 800 International Units/kg/day. The 800 International Units/kg/day dose is approximately 390 times the maximum recommended intranasal dose in humans (200 International Units) based on scaling for body surface area and a 20-fold conversion factor to account for low clinical exposure via the intranasal route.

#### **Mutagenesis**

Synthetic calcitonin-salmon tested negative for mutagenicity using *Salmonella typhimurium* (5 strains) and *Escherichia coli* (2 strains), with and without rat liver metabolic activation, and was not clastogenic in a

chromosome aberration test in Chinese Hamster V79 cells. There was no evidence that calcitonin-salmon was clastogenic in the in vivo mouse micronucleus test.

### Fertility

Effects of calcitonin-salmon on fertility have not been assessed in animals.

## 14 CLINICAL STUDIES

Two randomized, placebo-controlled, two-year trials were conducted in 266 postmenopausal women who were greater than 5 years postmenopause with spinal, forearm or femoral bone mineral density (BMD) at least one standard deviation below the normal value for healthy premenopausal women (T-score < -1). In both studies, a total of 144 patients received Miacalcin nasal spray 200 International Units or placebo daily. The intent-to-treat population comprised 139 patients who had at least one follow-up BMD measurement. In study 1, patients also received 500 mg daily calcium supplements, while in study 2, patients received no calcium supplementation. The primary endpoint for both studies was percent change in lumbar spine BMD at 2 years. Miacalcin nasal spray increased lumbar vertebral BMD relative to placebo in women with low bone mass who were greater than 5 years post menopause (see Table 3 below).

<b>Table 3: Miacalcin nasal spray: Lumbar Spine Bone Mineral Density In Women Greater Than 5 years Postmenopause With Low Bone Mass</b>		
	<b>Lumbar Spine Bone Mineral Density, Mean Change From Baseline (in %) at Month 24</b>	
	<b>Study 1 (with calcium supplement) n (ITT) = 100</b>	<b>Study 2 (no calcium supplement) n (ITT) = 39</b>
Miacalcin 200 IU NS daily	+1.56	+1.02
Placebo	+0.20	-1.85
Treatment Difference	+1.36	+2.87
p-value <sup>†</sup>	< 0.05	< 0.005
ITT: Intent To Treat IU: International Units NS: nasal spray <sup>†</sup> p-values by parametric testing (2-tailed 2-sample t-test)		

No effects of calcitonin-salmon nasal spray on cortical bone of the forearm or hip were demonstrated.

In clinical studies of postmenopausal osteoporosis, bone biopsy and radial bone mass assessments at baseline and after 26 months of daily injectable calcitonin-salmon indicate that calcitonin therapy results in the formation of normal bone.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

### How Supplied

Miacalcin Nasal Spray is available as a metered dose clear solution in a 3.7 mL fill clear glass bottle that contains 2200 International Units calcitonin-salmon per mL. A screw-on pump is provided. After priming, the

pump will deliver 200 International Units per activation (0.09 mL per spray)..... NDC 0078-0311-54

### **Storage and Handling**

Store unopened bottle in refrigerator between 2°C-8°C (36°F-46°F). Protect from freezing.

Store bottle in use at room temperature between 15°C-30°C (59°F-86°F) in an upright position, for up to 35 days. Each bottle contains at least 30 doses.

Discard bottle after 30 doses.

## **17 PATIENT COUNSELING INFORMATION**

See FDA-approved patient labeling (Patient Information and Instructions for Use).

- Instruct patients on pump assembly, priming of the pump, and nasal introduction of Miacalcin nasal spray. Although instructions for patients are supplied with the individual bottle, procedures for use should be demonstrated to each patient [*see Dosage and Administration (2.2)*]. Patients should notify their healthcare provider if they develop significant nasal irritation [*see Warnings and Precautions (5.3)*].
- Inform patients of the potential increase in risk of malignancy [*see Warnings and Precautions (5.4)*].
- Advise patients to maintain an adequate calcium (at least 1000 mg elemental calcium per day) and vitamin D (at least 400 International Units per day) intake [*see Dosage and Administration (2.3)*].
- Instruct patients to seek emergency medical help or go to the nearest hospital emergency room right away if they develop any signs or symptoms of a serious allergic reaction.
- Advise patients how to correctly store unopened and opened product [*see How Supplied/Storage and Handling (16)*]. Advise patients that the bottle should be discarded after 30 doses, because after 30 doses, each spray may not deliver the correct amount of medication even if the bottle is not completely empty.

T201X-XX  
March 2014

## Patient Information

### Miacalcin® (MEE-uh-KAL-sin) (calcitonin-salmon) nasal spray

Read this Patient Information before you start using Miacalcin and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

#### What is Miacalcin?

Miacalcin is a prescription medicine used to treat osteoporosis in women more than 5 years after menopause. Miacalcin should be used for women who cannot use other treatments or who choose not to use other treatments for osteoporosis.

It is not known if Miacalcin lowers the chance of having bone fractures.

Miacalcin has not been shown to be effective in women less than 5 years after menopause.

It is not known if Miacalcin is safe and effective in children under 18 years of age.

#### Who should not use Miacalcin?

Do not use Miacalcin if you:

- are allergic to calcitonin-salmon or any of the ingredients in Miacalcin. See the end of this leaflet for a complete list of ingredients in Miacalcin.

#### What should I tell my healthcare provider before using Miacalcin?

**Before you use Miacalcin, tell your healthcare provider if you:**

- have any other medical conditions
- have low calcium levels in your blood
- are pregnant or plan to become pregnant. It is not known if Miacalcin can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if Miacalcin passes into your breast milk. You and your healthcare provider should decide if you will use Miacalcin or breastfeed.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Especially tell your healthcare provider if you take:

- **lithium.** Your healthcare provider may need to change your dose of lithium while you use Miacalcin.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

#### How should I use Miacalcin?

- **For detailed instructions, see the Instructions for Use at the end of this Patient Information leaflet.**

- Use Miacalcin exactly as your healthcare provider tells you to use it.
- **Do not** use Miacalcin until your healthcare provider shows you and you understand how to use it correctly.
- Use 1 spray of Miacalcin, 1 time each day, in 1 nostril (inside your nose).
  - Start with 1 spray in your **left** nostril on your first day, followed by 1 spray in your **right** nostril on the second day.
  - Continue to switch nostrils for your dose each day.
- Your healthcare provider should check your nose before you start using Miacalcin and often while you are using it.
- Tell your healthcare provider if you start to have discomfort (irritation) in your nose that bothers you while you use Miacalcin.
- Your health care provider should prescribe calcium and vitamin D to help prevent low calcium levels in your blood while you use Miacalcin.
- Take your calcium and vitamin D as your healthcare provider tells you to.
- There are 30 doses (sprays) of Miacalcin in each bottle. After 30 doses, each spray may not give you the right amount of medicine, even if the bottle is not completely empty. Keep track of the number of doses of medicine used from your bottle.
- If you use too much Miacalcin, call your healthcare provider or go to the nearest hospital emergency room right away.

### What are the possible side effects of Miacalcin?

#### Miacalcin may cause serious side effects, including:

- **allergic reactions**

Some people have had an allergic reaction when using Miacalcin. Some reactions may be serious and can be life threatening. Call your healthcare provider or go to the nearest hospital emergency room right away if you have any of these symptoms of an allergic reaction.

- trouble breathing
- swelling of your face, throat or tongue
- fast heartbeat
- chest pain
- feel dizzy or faint

If you might be allergic to calcitonin-salmon, your healthcare provider should do a skin test before you use Miacalcin.

- **low calcium levels in your blood (hypocalcemia)**

Miacalcin may lower the calcium levels in your blood. If you have low blood calcium before you start using Miacalcin, it may get worse during treatment. Your low blood calcium must be treated before you use Miacalcin. Most people with low blood calcium levels do not have symptoms, but some people may have symptoms. Call your healthcare provider right away if you have any of these symptoms of low blood calcium:

- numbness or tingling in your fingers, toes, or around your mouth

Your healthcare provider should:

- do blood tests while you use Miacalcin
- prescribe calcium and vitamin D to help prevent low calcium levels in your blood while you use Miacalcin.

Take your calcium and vitamin D as your healthcare provider tells you to.

- **nose irritation**

Irritation of your nose can happen while you are using Miacalcin, especially if you are over 65 years of age. Call your healthcare provider right way if you have any of these symptoms of nose irritation:

- crusting
- dryness
- redness or swelling
- nose sores (ulcers)
- nose bleeds

Your healthcare provider may stop your treatment with Miacalcin until your nose irritation symptoms go away.

- **risk of cancer**

People who use calcitonin-salmon, the medicine in Miacalcin, may have an increased risk of cancer.

- **increase of certain cells (sediment) in your urine**

Your healthcare provider should test your urine often while you are using Miacalcin.

### **The most common side effects of Miacalcin include:**

- back pain
- muscle aches
- headache
- runny nose

These are not all the possible side effects of Miacalcin. For more information, ask your healthcare provider or pharmacist.

Tell your healthcare provider right away if you have any side effect that bothers you or does not go away.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How do I store Miacalcin?**

- Store open bottles of Miacalcin at room temperature between 59°F to 86°F (15°C to 30°C) for 35 days.
- Store unopened bottles of Miacalcin in the refrigerator between 36°F to 46°F (2°C to 8°C). Do not freeze.
- Store Miacalcin bottles in an upright position.
- Safely throw away Miacalcin in the trash after you have used 30 doses (sprays).

**Keep Miacalcin and all other medicines out of the reach of children.**

**General information about the safe and effective use of Miacalcin.**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use Miacalcin for a condition for which it was not prescribed. Do not give Miacalcin to other people, even if they have the same symptoms you have. It may harm them.

This Patient Information summarizes the most important information about Miacalcin. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about Miacalcin that is written for health professionals.

For more information, call 1-888-669-6682.

**What are the ingredients in Miacalcin?**

**Active Ingredients:** calcitonin-salmon

**Inactive Ingredients:** sodium chloride, benzalkonium chloride, hydrochloric acid and purified water.


**Instructions for Use**  
**Miacalcin® (MEE-uh-KAL-sin)**  
**(calcitonin-salmon)**  
**nasal spray**

**For Nasal Use Only.**

**Important information about your Miacalcin:**

- A single spray of MIACALCIN (calcitonin-salmon) nasal spray contains 1 daily dose of medicine.
- Each Miacalcin bottle contains the right amount of medicine. The bottle may not be completely filled to the top. This is normal.
- This package contains 1 bottle of Miacalcin and 1 screw-on pump. **See Figure A.**
- Store unopened bottles of Miacalcin in the refrigerator between 36°F to 46°F (2°C to 8°C). Do not freeze.
- After you open your bottle of Miacalcin, store it at room temperature between 59°F to 86°F (15°C to 30°C) in an upright position. Do not shake the bottle.

**Figure A**



**Preparing your Miacalcin:**

**Step 1.** Remove the bottle from your refrigerator and let it reach room temperature. Check the medicine in the bottle to make sure it is clear and colorless without particles.

**Important: If your Miacalcin bottle and pump has already been put together by your pharmacist or healthcare provider, go to Step 7.**



Figure B

**Step 2.** Lift up the blue plastic tab and carefully pull the metal safety seal off the bottle. **See Figure B.**




Figure C

**Step 3.** Keep the bottle upright and remove the rubber stopper from the bottle. **See Figure C.**



Figure D

**Step 4.** Hold the nose spray pump and gently remove the plastic protective cap from the bottom of the nose spray pump. **See Figure D.**

Do not push down on the pump when it is not attached to the bottle.




Figure E

**Step 5.** Hold the bottle upright and insert the nose spray pump into the bottle. Turn the pump clockwise to tighten it until it is securely attached to the bottle. **See Figure E.**




Figure F

**Step 6.** Gently pull the clear protective cap to remove it from the top of the nose spray pump. **See Figure F.**




Figure G

**Step 7.** Check to see if your Miacalcin has been primed.

- To make sure you get the right dose of Miacalcin medicine you must prime each new bottle and pump before you use it for the first time.
- If your pharmacist or healthcare provider puts the bottle and pump together for you, check to see if it has already been primed by pressing on the pump 1 time. **See Figure G.**
- If you see a full spray from the pump, the bottle and pump has already been primed for you.
- If you do not see a full spray, you must prime the bottle and pump.

**Step 8. Priming your Miacalcin:**

- Hold the bottle upright with your pointer finger and middle finger on the 2 side arms of the pump, and your thumb on the bottom of the bottle. Firmly press down on the arms of the pump and press down again if needed until you see a full spray of medicine. **See**

**Figure G.**

- Now your Miacalcin is ready for you to use.
- Do not prime the pump before you use it each day because this will waste your medicine.

**Giving your Miacalcin dose:**

**Step 9.** Insert the nasal spray pump in 1 side of your nose. **See Figure H.**

- Keep your head upright. Carefully tilt the bottle and place the nose spray pump into 1 side of your nose.



**Figure H**

**Step 10.** Firmly press down on the nose spray pump to release the medicine. **See Figure I.**

- Give 1 spray of Miacalcin, 1 time daily, in 1 side of your nose (nostril). Spray your medicine in a different side of your nose each day.
- You do not need to breathe in or inhale while you are giving your dose.
- You may not feel the spray inside your nose.
- Some of the medicine may drip out of your nose. This is normal, and you are still getting all of the medicine you need.



**Figure I**

**Cleaning your Miacalcin pump:**

**Step 11.** Wipe the nose spray pump with a clean, damp cloth 1 to 2 times a week. **See Figure J.**

- Dry the nose spray pump with a clean cloth.




Figure J




Figure K

**Storing your Miacalcin:**

**Step 12.** Gently put the protective cap back on the nasal spray pump.

- Hold the bottle with 2 fingers under the 2 side arms of the pump. **See Figure K.**
- Be careful not push down on the pump while you are putting the cap back on it.
- Do not refrigerate Miacalcin between doses.
- Store Miacalcin upright.
- Do not shake the bottle.

**When should I throw away Miacalcin?**

- Unopened, refrigerated bottles can be used until the expiration date stamped on the bottle and box.
- Throw away Miacalcin after you use 30 doses (sprays).
- Throw away Miacalcin bottles left at room temperature (opened or unopened) for more than 35 days.

For more information on MIACALCIN nasal spray and how to put it together, call Novartis Pharmaceuticals Corporation at 1-888-669-6682.

This Patient Information and Instructions for Use has been approved by the U.S. Food and Drug Administration.

Distributed by:

Novartis Pharmaceuticals Corporation  
East Hanover, New Jersey 07936

© Novartis

T201X-XX/T201X-XX

March 2014