

Trim Safe Area Bleed Score

w 3.9375"


**Convenient 2-PACK!**

NDC 50580-601-25  
For Ages 2 to 11

**Children's Motrin®**

contains ibuprofen  
**Oral Suspension (NSAID)**  
100 mg per 5 mL (teaspoon)

**Pain Reliever / Fever Reducer**



**Lasts up to 8 hours**

**Alcohol Free**

**4 fl oz (120 ml)**

NDC 50580-601-25  
For Ages 2 to 11

**Children's Motrin®**

contains ibuprofen  
**Oral Suspension (NSAID)**  
100 mg per 5 mL (teaspoon)

**Pain Reliever / Fever Reducer**



**Lasts up to 8 hours**

**Alcohol Free**

**4 fl oz (120 ml)**

Distributed by: McNeil Consumer Healthcare  
DIVISION OF MCNEIL – PPC, NC.  
FORT WASHINGTON, PA 19034 USA  
© MCN-PPC, NC. 2012 [www.motrin.com](http://www.motrin.com)  
McN-CM-0207



3 0045-0192-08 0


h 5.6562"

h 0.7"

Front

(b) (4)





146 mm

63.5 mm

**Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.**

**Active ingredient (in each 5 mL = 1 teaspoon)**      **Purpose**  
 ibuprofen 100 mg (NSAID)      Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Use:** temporarily relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache relieves fever

**Warnings**  
**Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling asthma (wheezing) shock skin redness rash blisters. If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child has had stomach ulcers or bleeding problems takes a blood thinning (anticoagulant) or steroid drug takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or other s) takes more or for a longer time than directed. Some throat warnings: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and/or mild may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. Do not use if the child has ever had an allergic reaction to

**EXP:** [blacked out]

**LOT:** [blacked out]

981801

NDC 50580-601-20

## Children's Motrin®

contains **Ibuprofen**  
**Oral Suspension (NSAID)**  
**100 mg per 5 mL (teaspoon)**

4 fl oz (120 mL)

**Original Berry Flavor**

any other pain reliever/fever reducer right before or after heart surgery  
**Ask a doctor before use if** stomach bleeding warning applies to your child  
 child has a history of stomach problems, such as heartburn child has problems or serious side effects from taking pain relievers or fever reducers  
 child has not been drinking fluids child has lost a lot of fluid due to vomiting or diarrhea child has high blood pressure, heart disease, liver cirrhosis, or kidney disease child has asthma child is taking a diuretic  
**Ask a doctor or pharmacist before use if the child is** under a doctor's care for any serious condition taking any other drug. **When using this product** take with food or milk if stomach upset occurs the risk of heart attack or stroke may increase if you use more than directed or for longer than directed. **Stop use and ask a doctor if** child experiences any of the following

signs of stomach bleeding feels faint vomit blood has bloody or black stools has stomach pain that does not get better the child does not get any relief within first day (24 hours) of treatment  
 fever or pain gets worse or lasts more than 3 days redness or swelling is present in the painful area any new symptoms appear. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** this product does not contain directions or complete warnings for adult use  
 do not give more than directed shake well before using find right dose on chart below. If possible, use weight to dose; otherwise use age. use only enclosed measuring cup. if needed, repeat dose every 6-8 hours do not use more than 4 times a day replace original bottle cap to maintain child resistance

Dosing Chart		
Weight (lb)	Age (yr)	Dose (tsp or mL)
under 2 years		
24-35 lbs	2-3 years	ask a doctor
36-47 lbs	4-5 years	1 1/2 tsp or 7.5 mL
48-59 lbs	6-8 years	2 tsp or 10 mL
60-71 lbs	9-10 years	2 1/2 tsp or 12.5 mL
72-95 lbs	11 years	3 tsp or 15 mL

**Other information** each teaspoon contains sodium 1 mg store between 20-25°C (68-77°F) do not use if bottle wrap imprinted "SAFETY SEAL" is broken or missing  
**Inactive ingredients:** acacia, fennel, polyethylene glycol, purified water, sodium benzoate, sucrose, xanthan gum  
**Questions or comments?** Call 1-877-895-3468: weekdays 8:00 AM to 6:00 PM EST  
**Distributed by:**  
 McNeil Consumer Healthcare, DIVISION OF MCNEIL-PPC, INC., FORT WASHINGTON, PA 19034 USA  
 MCNEIL-PPC, INC. 2011      www.motrin.com

(b) (4)

-----  
**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**  
-----

/s/  
-----

KAREN M MAHONEY  
12/11/2014