


Artwork No. 6-000449, ChloroPrep One-Step Sepp Cylrel

(b) (4)

(b) (4)

t'



(b) (4)

Dimensions: 7.5" x 2.5"

2.5"

ChloroPrep® One-Step
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
Non-sterile Solution
 Applicator is sterile if package is intact
 0.67ml SEPP® APPLICATOR

Drug Facts	
Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v.....	Antiseptic
Isopropyl alcohol 70% v/v.....	Antiseptic

Use for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only
Flammable, keep away from fire or flame
 ■ do not use with electrocautery procedures

Do not use
 ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
 ■ for lumbar puncture or in contact with the meninges
 ■ on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
 ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
 ■ maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm)
 ■ remove applicator from package; do not touch applicator tip
 ■ hold the sponge tip down and pinch the barrel of the applicator only once to activate the ampule and release the antiseptic
 ■ wet the sponge tip by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
 ■ completely wet the treatment area with antiseptic
 ■ **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds.
 ■ **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute.
 ■ do not blot or wipe away
 ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F)
 ■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients USP purified water

Questions? ■ www.chloroprep.com
 ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

See other side for additional information

7.5"

Front

ChloroPrep® One-Step
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
Non-sterile Solution
 Applicator is sterile if package is intact

0.67 ml SEPP® APPLICATOR

Cat. No. 260441NSB / 260449NSB
 NDC 054365-400-03

See Drug Facts for Full Labeling Information

 Do Not Reuse

 Not made with natural rubber latex

External Use Only Professional Use Only


CareFusion, El Paso, TX 79912

Artwork 6-775449F · NXXXX
 Made in the USA of US and imported parts

Back

(b) (4)

Artwork No. 6-824479, Chloraprep One-Step 1ml Intermediate Carton




NDC 054365-400-15
1 ml
 APPLICATORS
 Clear

Do Not Reuse
 Not made with natural rubber latex
 STERILE EO

Chloraprep®
One-Step™

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
 Non-sterile Solution
 Applicator is sterile if package is intact



Professional Use Only
 Store between 15-30 °C (59-86 °F)
 Avoid freezing and excessive heat above 40 °C (104 °F)

Chloraprep®
One-Step™

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
1 ml
 APPLICATORS
 Clear

LOT XXXXX (printed in line) No Varnish EXP DATE XXXX
 Barcode Placeholder

Cat. No. 260479
 ©2012 CareFusion. All rights reserved. Chloraprep® is a registered trademark of CareFusion Corporation or one of its subsidiaries.

No Varnish / No printing area

Drug Facts		Drug Facts (continued)	
Active ingredients Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	Purposes Antiseptic Antiseptic	Directions ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²). ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings <u>only once</u> to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds. Allow the area to air dry for approximately 30 seconds. ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes. Allow the area to air dry for approximately 1 minute. ■ do not blot or wipe away. ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.	Other information ■ store between 15-30 °C (59-86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)
Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.	Warnings For external use only. Flammable, keep away from fire or flame. ■ do not use with electrocautery procedures	When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser	Inactive ingredients USP purified water
Do not use ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser	Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)	

Artwork No. 6-001479, ChloroPrep One-Step 1 mL Cyrel

(b) (4)

(b) (4)


(b) (4)



Trim Area

Expiry Date and Lot No. : reserved for manufacturing purpose only.
Reference ID: 3534450

(b) (4)



(b) (4)

4.75" width

Front

Chloraprep® One-Step
 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
Non-sterile Solution
 Applicator is sterile if package is intact
1 ml APPLICATOR

Not made with natural rubber latex
 Do Not Reuse
Professional Use Only
External Use Only

Cat. No. 260479 / 260479NS / 260479NSB NDC 054365-400-15

Drug Facts	Purposes
Active ingredients Chlorhexidine gluconate 2% w/v.....	Antiseptic
Isopropyl alcohol 70% v/v.....	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only. Flammable, keep away from fire or flame.
 ■ do not use with electrocautery procedures

Do not use
 ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
 ■ for lumbar puncture or in contact with the meninges
 ■ on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶

3.8" length


Chloraprep® One-Step
 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm ²) ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic. ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds. ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute. ■ do not blot or wipe away. ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.
Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)
Inactive ingredients USP purified water
Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, El Paso, TX 79912 · Artwork 6-775479 · DXXXX · Made in the USA of US and imported parts


Back

Artwork No. 6-824299, ChloraPrep One-Step Frepp Intermediate Carton



NDC 054365-400-02
1.5ml FREPP[®] APPLICATORS
 Clear
 Do Not Reuse
 Not made with natural rubber latex
 STERILE EO
 20 applicators
 0.05 fl. oz (1.5 ml) each

ChloraPrep[®] One-Step
 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
Non-sterile Solution
 Applicator is sterile if package is intact



ChloraPrep[®] One-Step
 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
1.5ml FREPP[®] APPLICATORS
 Clear

LOT XXXXXX EXP DATE: XXXX
 No Varnish
 (01) 20354365400024

Professional Use Only
 Store between 15-30 °C (59-86 °F)
 Avoid freezing and excessive heat above 40 °C (104 °F)

Drug Facts		Drug Facts (continued)	
Active ingredients Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	Purposes Antiseptic Antiseptic	Directions Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm ²) remove applicator from package; do not touch sponge hold the applicator horizontally with the sponge down. Pinch wings <u>only once</u> to activate the ampule and release the antiseptic. wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin completely wet the treatment area with antiseptic dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds. Allow the area to air dry for approximately 30 seconds. moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes. Allow the area to air dry for approximately 1 minute. do not blot or wipe away discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.	Warnings For external use only. Flammable, keep away from fire or flame. do not use with electrocautery procedures
Do not use on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol for lumbar puncture or in contact with the meninges on open skin wounds or as a general skin cleanser	When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	Other information store between 15-30 °C (59-86 °F) avoid freezing and excessive heat above 40 °C (104 °F)	Inactive ingredients USP purified water
Questions? www.chloraprep.com call 1-800-523-6902 (M-F 8 a.m.-5 p.m. CST)			

BACK PANEL
 'ChloraPrep[®] & 'One-Step'
 Interstate Bold (VECTOR)
DRUG FACTS BOX
 'Drug Facts'
 14 pt Helvetica Bold Oblique
Headings
 8 pt Helvetica Bold Oblique
Body Text
 6 pt Helvetica Regular/Bold
Hairline - 0.5 pt
Box Barline - 2.5 pt
Bullets - 5 pt zapf dingbats

Artwork No. 6-000299, ChloroPrep One-Step Frepp Cyrel

(b) (4)

(b) (4)

72 mm

40 mm

Exp. Date xx/xx
Lot No. xxxxx

Cat. No. 260299 NDC 054365-400-02

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

Non-sterile Solution
Applicator is sterile if package is intact

1.5 ml FREPP® APPLICATOR

WARNING. FLAMMABLE.
KEEP AWAY FROM FIRE OR FLAME.

CareFusion, El Paso, TX 79912
1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST)
Artwork No. 6-000299 - MXXXX
Made in the USA of US and imported parts

Do Not Reuse

Bend and Peel

Trim Area

Artwork No. 6-001299, ChloroPrep One-Step Frepp Cyrel (No Lot/Exp. Date)

(b) (4)



(b) (4)



(b) (4)

66 mm

40 mm

Placeholder for lot and expiry data

Cat. No. 260299 NDC 054365-400-02

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

Non-sterile Solution
Applicator is sterile if package is intact

1.5 ml FREPP® APPLICATOR

 **WARNING. FLAMMABLE.**
KEEP AWAY FROM FIRE OR FLAME.

CareFusion, El Paso, TX 79912
1-800-523-0502 (M-F 8 a.m.- 5 p.m. CST)

Artwork No. 6-001299 · FXXXX
Made in the USA of US and imported parts

Do Not Reuse 



Bend and Peel

(b) (4)

Dimensions: 7.5" x 2.5"

← 2.5" →

ChloroPrep® One-Step
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact
 1.5 ml FREPP® APPLICATOR

Drug Facts				
<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%;">Active ingredients</td> <td style="width: 50%;">Purposes</td> </tr> <tr> <td>Chlorhexidine gluconate 2% w/v.....Antiseptic</td> <td>Isopropyl alcohol 70% v/v.....Antiseptic</td> </tr> </table>	Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v.....Antiseptic	Isopropyl alcohol 70% v/v.....Antiseptic
Active ingredients	Purposes			
Chlorhexidine gluconate 2% w/v.....Antiseptic	Isopropyl alcohol 70% v/v.....Antiseptic			
Use For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.				
Warnings For external use only Flammable, keep away from fire or flame. <ul style="list-style-type: none"> ■ do not use with electrocautery procedures Do not use <ul style="list-style-type: none"> ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general cleanser When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.				
Directions <ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²) ■ remove applicator from package; do not touch sponge ■ hold the applicator horizontally with the sponge down. Pinch wings <u>only once</u> to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds. ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute. ■ do not blot or wipe away ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)				
Inactive ingredients USP purified water				
Questions? ■ www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)				

See other side for additional information

7.5"

Front

ChloroPrep® One-Step
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact
 1.5 ml FREPP® APPLICATOR

Cat. No. 260299NSB / 260599NSB
 NDC 054365-400-02

See Drug Facts for Full Labeling Information

 Do Not Reuse
 External Use Only

 Not made with natural rubber latex
 Professional Use Only

CareFusion, El Paso, TX 79912

Artwork 6-775403F · NXXXX
 Made in the USA of US and imported parts

Back


(b) (4)

(1/8" from outline to text for cutting process)

A k S

$1\frac{15}{32}$

ChloroPrep®
One-Step
2% w/v povidone-iodine (PVP-I) solution
ad 0.05% w/v chlorhexidine gluconate (CHG)
antiseptic solution
N t t i s i t
P t t k P t
A l t t l f k t t



5 a p i a o s

3 ml
A P L C A O R S

ChloroPrep®
One-Step
d h d l l ()

3 ml
A P P I C T O R S

ChloroPrep®
One-Step
d h d l l ()

3 ml
A P P I A T O R S



3 ml
A P P I A T O R S

STERILE (EO)
b l h l

25 p p c t r s

25 p p c t r s

ChloroPrep®
One-Step

2% w/v povidone-iodine (PVP-I) solution
ad 0.05% w/v chlorhexidine gluconate (CHG)
antiseptic solution
N t t i s i t
P t t k P t
A l t t l f k t t



ChloroPrep®
One-Step
d h d l l ()

3 ml
P P L C A O R S



$3\frac{1}{4}$

$1\frac{7}{4}$

$6\frac{1}{2}$

$2\frac{5}{6}\frac{1}{4}$

$3\frac{1}{16}$

$\frac{1}{16}$

$1\frac{2}{8}$

$\frac{5}{8}$

$1\frac{2}{8}$


$5\frac{9}{32}$

$3\frac{7}{32}$

A k

$1\frac{15}{32}$

ChloroPrep®
With Tint
2% w/v hexachlorophene (HCl) solution
in alcohol (90% v/v)
Antiseptic
Nontoxic
Airtight



3 ml
APPLICATORS
STERILE

ChloroPrep®
With Tint
3 ml
APPLICATORS
STERILE

ChloroPrep®
With Tint
3 ml
APPLICATORS
STERILE

3 ml
APPLICATORS
STERILE

ChloroPrep®
With Tint
2% w/v hexachlorophene (HCl) solution
in alcohol (90% v/v)
Antiseptic
Nontoxic
Airtight

ChloroPrep®
With Tint
3 ml
APPLICATORS
STERILE

3 ml
APPLICATORS
STERILE



$3\frac{1}{16}$

$2\frac{5}{64}$

1

$12\frac{7}{8}$

$5\frac{5}{8}$

$12\frac{7}{8}$

$5\frac{19}{32}$

$37\frac{9}{32}$

$3\frac{1}{4}$

$6\frac{1}{2}$

$17\frac{1}{4}$

A k

$1\frac{15}{32}$

$3\frac{11}{16}$

1

$12\frac{7}{8}$

$5\frac{5}{8}$

$12\frac{7}{8}$

$5\frac{19}{32}$


$37\frac{9}{32}$


$3\frac{1}{4}$

$6\frac{1}{2}$

$2\frac{51}{64}$



$17\frac{1}{4}$






(b) (4)

Lot/Exp Placeholder - for reference only.





(b) (4)

(b) (4)

4.75" width

Front

<div style="background-color: #f00; color: white; text-align: center; padding: 2px;">WARNING</div> <div style="background-color: #f00; color: white; text-align: center; padding: 2px;">FLAMMABLE</div> <p>Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet materials from prep area 	<p>Cat. No. 260415 / 260415NS / 260415NSB NDC 054365 400-11 Cat. No. 260400 / 260400NS / 260500NS NDC 054365 400 01</p> <h2 style="margin: 0;">ChloraPrep® One-Step ChloraPrep® With Tint</h2> <p style="margin: 0;">2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation Non-sterile Solution Applicator is sterile if package is intact</p> <p style="text-align: center; font-weight: bold; font-size: 1.2em;">3 ml APPLICATOR</p> <p style="font-size: 0.8em;">Professional Use Only Do Not Reuse Not made with natural rubber latex</p>																				
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th colspan="2" style="text-align: left; padding: 2px;">Drug Facts</th> </tr> <tr> <td style="width: 60%; padding: 2px;">Active ingredients</td> <td style="width: 40%; padding: 2px;">Purposes</td> </tr> <tr> <td style="padding: 2px;">Chlorhexidine gluconate 2% w/v.....</td> <td style="padding: 2px;">.....Antiseptic</td> </tr> <tr> <td style="padding: 2px;">Isopropyl alcohol 70% v/v.....</td> <td style="padding: 2px;">.....Antiseptic</td> </tr> <tr> <td colspan="2" style="padding: 2px;">Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.</td> </tr> <tr> <td colspan="2" style="padding: 2px;">Warnings</td> </tr> <tr> <td colspan="2" style="padding: 2px;">For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</td> </tr> <tr> <td colspan="2" style="padding: 2px;"> <ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area </td> </tr> <tr> <td colspan="2" style="padding: 2px;">Do not use</td> </tr> <tr> <td colspan="2" style="padding: 2px;"> <ul style="list-style-type: none"> ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser </td> </tr> </table>		Drug Facts		Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v.....Antiseptic	Isopropyl alcohol 70% v/v.....Antiseptic	Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.		Warnings		For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:		<ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area 		Do not use		<ul style="list-style-type: none"> ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser 	
Drug Facts																					
Active ingredients	Purposes																				
Chlorhexidine gluconate 2% w/v.....Antiseptic																				
Isopropyl alcohol 70% v/v.....Antiseptic																				
Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.																					
Warnings																					
For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:																					
<ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area 																					
Do not use																					
<ul style="list-style-type: none"> ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser 																					

3.8" length


ChloraPrep® One-Step / ChloraPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)
When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions
<ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ use in a well ventilated area ■ maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²) ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for approximately 30 seconds. Allow solution to air dry for approximately 30 seconds. Do not blot or wipe away. ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes of the sponge for approximately 2 minutes. Allow solution to air dry for approximately 1 minute. Do not blot or wipe away. ■ if using an ignition source, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.
Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) ■ the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.
Inactive ingredients FD&C green #3 dye (Scrub-Teal applicators only), FD&C yellow #6 dye (Hi-Lite Orange applicators only), USP purified water
Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, El Paso, TX 79912 · Artwork 6-775400 · GXXX · Made in the USA of US and imported parts

Back



A k # 0 % S

CloraPrep®
 One-Step
 % / clreine lcaec G
 nd 0% v v a o h l p)
 pt t p t k p t
 N t i s t k t t
 A l t t i f k t t

10.5 ml
 PPL CATO S

CloraPrep®
 One-Step
 % h h d i l ()
 d % i l l i c)
 10.5 ml
 APP IC TORS

CloraPrep®
 One-Step
 % h h d i l ()
 d % i l l i c)
 10.5 ml
 PPL CATO S

10.5 ml
 APP IC TORS

b d i h l
 (STERILE EO)

25 p l c t r s

25 p l c t r s

CloraPrep®
 One-Step
 2% w v h o h x d n e u o n t (H G)
 a n 7 % / i o r p l i o o (A)
 P t t P t t S k P t
 N t i s l t
 A l t t i f k t t



WARNING
 FLAMMABLE

CloraPrep®
 One-Step
 % h h d i l ()
 d % i l l i c)
 10.5 ml
 APP I ATORS

B

A k 0

1 15/32



25 p p c t r s
S b T I
10.5 ml
APPL CA ORS

Chloraprep®
With Tint
10.5 ml
APPL CA ORS
S b T I

Chloraprep®
With Tint
10.5 ml
A PL CATO S
S T I

10.5 ml
APP ICA ORS
S b T I
STERILE EO

Chloraprep®
With Tint
2% w/v Chlorhexidine (G)
and 0.1% Povidone-iodine (P)
Solution for Skin Preparation
Net Weight 10.5 ml (0.35 fl. oz.)
Net Content 10.5 ml (0.35 fl. oz.)
Net Volume 10.5 ml (0.35 fl. oz.)



Chloraprep®
With Tint
10.5 ml
APP IC TORS
S b T I



3 1/16

1 1/16

1 2 7/8

5 5/8

1 2 7/8

5 1 9/32

3 7 1 9/32

3 1/4


6 1/2

2 5 1/6 4






1 7 1/4

A k 0

$1\frac{15}{32}$



2% wvc o h e d n e l c n t e (HG)
and 0% / i o r p l l o o (P)
P t t P t S k P t
N t l S l t
A l t t l f k t t



$3\frac{1}{16}$

$\frac{11}{1}$

$12\frac{7}{8}$

$5\frac{5}{8}$

$12\frac{7}{8}$

$5\frac{19}{32}$

$37\frac{9}{32}$

$3\frac{1}{4}$

$17\frac{1}{4}$


$6\frac{1}{2}$

$2\frac{5}{64}$

(b) (4)


Part No. 6-773700
10.5-ml ChloroPrep One-Step
In-Line Mold Labels

(b) (4)




(b) (4)

Part No. 6-773725
10.5-ml ChloroPrep With Tint
In-Line Mold Labels




Part No. 6-773715
10.5-ml ChloroPrep With Tint
In-Line Mold Labels



Text starts 3mm from top of package die and ends 15mm from bottom of package die
 Text (Drug Facts Box) is centered 10.25mm from either side




(b) (4)



Text starts 3mm from top of package die and ends 15mm from bottom of package die
Text is centered 10.25mm from either side



(b) (4)



4.75" width

Front

⚠ WARNING FLAMMABLE	<p>ChloraPrep® One-Step ChloraPrep® With Tint</p> <p>2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation Non-sterile Solution Applicator is sterile if package is intact</p> <p>10.5 ml APPLICATOR </p> <p>Professional Use Only Do Not Reuse Not made with natural rubber latex</p> <p>Cat. No. 260700 / 260700NS / 260700NSB NDC 054365-400-04 Cat. No. 260715 / 260715NS / 260715NSB NDC 054365-400-12 Cat. No. 260725 / 260725NS / 260725NSB NDC 054365-400-09</p>						
<p>Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet materials from prep area 	<p>Drug Facts</p> <table border="0" style="width: 100%;"> <tr> <td style="width: 60%;">Active ingredients</td> <td style="width: 40%;">Purposes</td> </tr> <tr> <td>Chlorhexidine gluconate 2% w/v.....</td> <td>Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v.....</td> <td>Antiseptic</td> </tr> </table> <p>Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.</p> <p>Warnings</p> <p>For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY</p> <ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area <p>Do not use</p> <ul style="list-style-type: none"> ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser 	Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v.....	Antiseptic	Isopropyl alcohol 70% v/v.....	Antiseptic
Active ingredients	Purposes						
Chlorhexidine gluconate 2% w/v.....	Antiseptic						
Isopropyl alcohol 70% v/v.....	Antiseptic						


3.8" length

<p>ChloraPrep® One-Step / ChloraPrep® With Tint 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)</p>
<p>Drug Facts (continued)</p> <p>When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ use in a well ventilated area ■ maximal treatment area for one applicator is approximately 8.4 in. x 8.4 in. (457 cm²) ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes ■ do not allow solution to pool; tuck prep towels to absorb solution, and then remove ■ allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. <p>Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)</p> <p>■ the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.</p> <p>Inactive ingredients FD&C green #3 dye (Scrub-Teal applicators only), FD&C yellow #6 dye (Hi-Lite Orange applicators only), USP purified water</p> <p>Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)</p>
<p>CareFusion, El Paso, TX 79912 · Artwork No. 6-640248 · KXXXX · Made in the USA of US and imported parts</p>

Back

(b) (4)

Part No. 6-773800-L
26-ml ChloraPrep One-Step
In-Line Mold Labels



(b) (4)


(b) (4)

(b) (4)

(b) (4)

Part No. 6-773825-L
26-ml ChloroPrep with tint
In-Line Mold Labels

(b) (4)



(b) (4)

(b) (4)


Part No. 6-773815-L
26-ml ChloroPrep with tint
In-Line Mold Labels

(b) (4)

(b) (4)


(b) (4)

(b) (4)



(b) (4)

Part No. 6-773800-R
26-ml ChloroPrep One-Step
In-Line Mold Labels



(b) (4)

(b) (4)


(b) (4)

(b) (4)

(b) (4)


Part No. 6-773825-R
26-ml ChloroPrep with tint
In-Line Mold Labels

(b) (4)



(b) (4)

Part No. 6-773815-R
26-ml ChloraPrep with tint
In-Line Mold Labels




(b) (4)


(b) (4)

(b) (4)

Artwork No. 6-787800, 26-mL ChloroPrep One-Step Applicator Lidding




(b) (4)



(b) (4)

Lot # and Exp. Date are displayed at this end of the package.



5" width Front

**⚠ WARNING
FLAMMABLE**

Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

**ChloroPrep® One-Step
ChloroPrep® With Tint**

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is sterile if package is intact
26 ml APPLICATOR

Professional Use Only Do Not Reuse Not made with natural rubber latex

Cat. No. 260800 / 260800NS / 260800NSB NDC 054365 400-14
Cat. No. 260815 / 260815NS / 260815NSB NDC 054365 400-13
Cat. No. 260825 / 260825NS / 260825NSB NDC 054365 400-05

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v.....	Antiseptic
Isopropyl alcohol 70% v/v.....	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

4" length

ChloroPrep® One-Step / ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm²).
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings **only once** to activate the ampules and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.


Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) ■ the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.

Inactive ingredients FD&C green #3 dye (Scrub-Teal applicators only), FD&C yellow #6 dye (Hi-Lite Orange applicators only), USP purified water

Questions? ■ www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, El Paso, TX 79912 · Artwork No. 6-775815 · JXXXX · Made in the USA of US and imported parts

Back



(b) (4)

5.5"

4.5"

1.25"

1.625"

Chloraprep® Single Swabstick
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is non-sterile
1.75 ml APPLICATOR
See Drug Facts for full labeling information
Made in the USA of US and imported parts

Do Not Reuse

Cat. No. 260100 NDC 054365-400-07

This panel shows the top portion of the packaging with dimensions. A pink double-headed arrow indicates a width of 5.5 inches. A blue double-headed arrow indicates an inner width of 4.5 inches. On the right side, a blue double-headed arrow indicates a height of 1.25 inches, and a pink double-headed arrow indicates a total height of 1.625 inches. The central area contains a product illustration of a swabstick, the product name, active ingredients, and usage instructions.

Manufactured by:
CareFusion
EL Paso, TX 79912
1-800-523-0502 (CST)
(M-F 8 a.m. - 5 p.m. CST)
www.chloraprep.com
Artwork No. 6-800101
JXXX

Tear Here

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Directions:

- Tear pouch at notch.
- Pull across package to expose the end of the swabstick.
- Do not touch foam when removing swabstick from pouch.
- Place foam flat side down on treatment area.

Lot # and Exp. Date are displayed on package.

This panel shows the bottom portion of the packaging. On the left, it lists manufacturing information. A vertical dashed arrow labeled "Tear Here" points to a notch. A warning about flammability is present. A list of directions for use is provided. On the right, a red-bordered box states that the lot number and expiration date are displayed on the package.

.95" From left edge of live area to point of arrow for tear perforation

A dimension line with an arrowhead pointing to the right, indicating a distance of .95 inches from the left edge of the live area to the point of the tear perforation arrow.

4.75" width
Front

Cat. No. 260100 / 260100B NDC 054365-400-07
 Cat. No. 260103 / 260103B NDC 054365-400-08

Chloraprep® Single Swabstick/Chloraprep® Triple Swabstick
 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is non-sterile
 1.75 ml APPLICATOR / 5.25 ml APPLICATOR

Not made with natural rubber latex Do Not Reuse Professional Use Only
 External Use Only

Drug Facts	
Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic
Use for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infection.	
Warnings	
For external use only. Flammable, keep away from fire or flame.	
<ul style="list-style-type: none"> do not use with electrocautery procedures 	
Do not use	
<ul style="list-style-type: none"> on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol for lumbar puncture or in contact with the meninges on open skin wounds or as a general skin cleanser 	
When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	


3.8" length

Chloraprep® Single Swabstick/Chloraprep® Triple Swabstick
 2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (cont nued)
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions
<ul style="list-style-type: none"> use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. maximal treatment area using <ul style="list-style-type: none"> one single swabstick applicator is approximately 2.5 in. x 2.5 in. (42 cm²) three swabsticks sequentially is approximately 5 in. x 5 in. (160 cm²) tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area. completely wet the treatment area with antiseptic dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds (when using the triple swabstick applicators, use each swabstick sequentially within the 30 seconds). Allow the area to air dry for approximately 30 seconds. Do not blot or wipe away. moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes (when using the triple swabstick applicators, use each swabstick sequentially within the 2 minutes). Allow the area to air dry for approximately 1 minute. Do not blot or wipe away. discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.
Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)
Inactive ingredients USP purified water
Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, El Paso, TX 79912 · Artwork No. 6-656900 · NXXXX · Made in the USA of US and imported parts

Back



(b) (4)

5.5"

5"

2"

2.375"

Chloraprep® Triple Swabstick
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
Non-sterile Solution
Applicator is non-sterile
5.25 ml APPLICATOR
See Drug Facts for full labeling information
Made in the USA of US and imported parts

Do Not Reuse

Car. No. 260103 NDC 054365-400-08

Illustration of three swabsticks.

Manufactured by:
CareFusion
EL Paso, TX 79912
1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST)
www.chloraprep.com

Artwork No. 6-800104 IXXXX

Tear Here

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Directions:

- Tear pouch at notch.
- Pull across package to expose the ends of the swabsticks.
- Do not touch foam when removing swabsticks from pouch.
- Place foam flat side down on treatment area.

Lot # and Exp. Date are embossed on package film

95" From left edge of live area to point of arrow for tear perforation