




(b) (4)



**Drug Facts (continued)**  
 Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.  
 If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

adults and children 6 years and over	chew 2 tablets daily; not more than 2 tablets in 24 hours
children 2 to under 6 years of age	chew 1 tablet daily; not more than 1 tablet in 24 hours
children under 2 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Drug Facts (continued)**  
**Other information**  
 ■ phenylethanolamines: contains phenylalanine 1.4 mg per tablet  
 ■ safety sealed: do not use if the individual blister unit imprinted with Children's Claritin® is open or torn  
 ■ store between 20° to 25°C (68° to 77°F)

**Inactive ingredients**  
 aspartame, citric acid anhydrous, colloidal silicon dioxide, D&C red No. 27 aluminum lake, FD&C blue No. 2 aluminum lake, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid

**Questions or comments?**  
 1-800-CLARITIN (1-800-252-7484) or www.claritin.com

© Copyright & Distributed by MSD Consumer Care, Inc., PO Box 377, Memphis, TN 38151 USA, a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA. All rights reserved. Product of Ireland. 27780-06

Sample - Not for Sale NDC 11523-4328-4

**Children's Claritin Chewables**  
 loratadine 5 mg/antihistamine  
 Indoor & Outdoor Allergies

**Non-Drowsy\* 24 Hour Relief of:**  
 • Sneezing  
 • Runny Nose  
 • Itchy, Watery Eyes  
 • Itchy Throat or Nose

**Grape Flavored**  
 2 CHEWABLE TABLETS

\* When taken as directed. See Drug Facts Panel.

**Drug Facts**

Active ingredient (in each tablet)	Purpose
Loratadine 5 mg	Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  
 ■ runny nose  
 ■ itchy, watery eyes  
 ■ sneezing  
 ■ itching of the nose or throat

**Warnings**  
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients.  
 Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

**When using this product** do not take more than directed. Taking more than directed may cause drowsiness.

2.81466" CUT OFF

ages 2 years and older  
**Children's Claritin Chewables**  
 loratadine 5 mg/antihistamine

**Drug Facts**  
**Active ingredient (in each tablet) Purpose**  
 Loratadine 5mg ..... Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  
 ■ runny nose ■ itchy, watery eyes  
 ■ sneezing ■ itching of the nose or throat

**Warnings**  
**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.  
**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.  
**When using this product** do not take more than directed. Taking more than directed may cause drowsiness.  
**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.  
**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

adults and children 6 years and over	chew 2 tablets daily; not more than 2 tablets in 24 hours
children 2 to under 6 years of age	chew 1 tablet daily; not more than 1 tablet in 24 hours
children under 2 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Drug Facts (continued)**  
**Other information**  
 ■ phenylethanolamines: contains phenylethylamine 1.4 mg per tablet  
 ■ safety sealed: do not use if the individual blister unit imprinted with Children's Claritin® is open or torn  
 ■ store between 20° to 25°C (68° to 77°F)

**Inactive ingredients**  
 aspartame, citric acid anhydrous, colloidal silicon dioxide, D&C red No. 27 aluminum lake, FD&C blue No. 2 aluminum lake, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid

**Questions or comments?**  
 1-800-CLARITIN (1-800-252-7484) or www.claritin.com

The graphics on this product constitute trademarks of MSD Consumer Care, Inc.  
 © Copyright & Distributed by MSD Consumer Care, Inc., PO Box 377, Memphis, TN 38151 USA, a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA. All rights reserved. Product of Ireland. 36505-04

LOT & EXP

ages 2 years and older  
**Children's Claritin Chewables**  
 loratadine 5 mg/antihistamine

**Non-Drowsy\* Allergy Relief**

**Indoor & Outdoor Allergies**

50 2-Count Samples  
 Total 100 Chewable Tablets

\*When taken as directed. See Drug Facts Panel.

PROFESSIONAL SAMPLES - NOT FOR SALE

ages 2 years and older  
**Children's Claritin Chewables**  
 loratadine 5 mg/antihistamine

**Non-Drowsy\* Allergy Relief**

**Indoor & Outdoor Allergies**

50 2-Count Samples  
 Total 100 Chewable Tablets

\*When taken as directed. See Drug Facts Panel.

PROFESSIONAL SAMPLES - NOT FOR SALE

ages 2 years and older  
**Children's Claritin Chewables**  
 loratadine 5 mg/antihistamine

**Non-Drowsy\* Allergy Relief**

**Indoor & Outdoor Allergies**

50 2-Count Samples  
 Total 100 Chewable Tablets

\*When taken as directed. See Drug Facts Panel.

PROFESSIONAL SAMPLES - NOT FOR SALE

ages 2 years and older  
**Children's Claritin Chewables**  
 loratadine 5 mg/antihistamine

**Non-Drowsy\* Allergy Relief**

**Indoor & Outdoor Allergies**

50 2-Count Samples  
 Total 100 Chewable Tablets

\*When taken as directed. See Drug Facts Panel.

PROFESSIONAL SAMPLES - NOT FOR SALE

ages 2 years and older  
**Children's Claritin Chewables**  
 loratadine 5 mg/antihistamine

**Non-Drowsy\* Allergy Relief**

**Indoor & Outdoor Allergies**


50 2-Count Samples  
 Total 100 Chewable Tablets

\*When taken as directed. See Drug Facts Panel.

PROFESSIONAL SAMPLES - NOT FOR SALE



(b) (4)





ITM# 809810  
4 1100 80981 0

The graphics on the front panel constitute trademarks of MSD Consumer Care, Inc. © Copyright & Distributed by MSD Consumer Care, Inc., PO Box 377, Memphis, TN 38151 USA, a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA. All rights reserved. Product of Ireland. 32435-00

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each tablet)</b> Purpose Loratadine 5 mg.....Antihistamine</p> <p><b>Uses</b> temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  <ul style="list-style-type: none"> <li>■ runny nose</li> <li>■ itchy, watery eyes</li> <li>■ sneezing</li> </ul> </p> <p><b>Warnings</b>                  Do not use if you have ever had an allergic reaction to this product or any of its ingredients.                  Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.                  When using this product do not take more than directed. Taking more than directed may cause drowsiness.                  Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.                  If pregnant or breast-feeding, ask a health professional before use.                  Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p><b>Questions or comments?</b> 1-800-CLARITIN (1-800-252-7484) or www.claritin.com</p> <p><b>Inactive ingredients</b> D&amp;C red No. 27 aluminum lake, FD&amp;C blue No. 2 aluminum lake, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid</p> <p><b>Other information</b>  <ul style="list-style-type: none"> <li>■ phenylephrine: contains phenylephrine 1.4 mg per tablet</li> <li>■ safety sealed: do not use if the individual blister unit is opened or torn</li> <li>■ store between 20° to 25°C (68° to 77°F)</li> </ul> </p> <p><b>Directions</b>                  adults and children                  chew 2 tablets daily; not more than 2 tablets in 24 hours                  6 years and over                  chew 1 tablet daily; not more than 1 tablet in 24 hours                  6 years of age                  children under 6 years of age                  ask a doctor                  2 years of age                  consumers with liver or kidney disease                  ask a doctor</p>
--	--



**50**  
CHEWABLE TABLETS



**Children's Claritin Chewables**  
2 years and older  
loratadine 5 mg/antihistamine



**Children's Claritin Chewables**  
ages 2 years and older  
loratadine 5 mg/antihistamine  
**50 CHEWABLE TABLETS**

**Indoor & Outdoor Allergies**




**50**  
CHEWABLE TABLETS

6.0625

6



07.14.14 mpls



(b) (4)