

Labeling Format Information:			
Fonts : Helvetica Roman , Bold , Bold Oblique			
Drug Facts :	9,5pt	Leading :	6,5pt
Header :	8pt	Bullets :	5pt
Subheader :	6pt	Barlines :	2,5pt
Body Text :	6pt	Hairlines :	0,5pt
Horizontal Scale: 80%-100%		Average Kerning: 0	

100%出力

**6/5** Salonpas® Pain Relief Patch  
 Menthol 3%, Methyl Salicylate 10%

Manufactured by  
 Hisamitsu Pharmaceutical Co., Inc.  
 JAPAN SAGA-TOSU  
 MADE IN JAPAN

Distributed by  
 Hisamitsu America, Inc.  
 Hoffman Park, NJ 07932  
 ® : Registered Trade Mark

# Salonpas Pain Relief Patch

Menthol 3%, Methyl Salicylate 10%

**6/5 patches**

for temporary relief of mild to moderate pain

**BONUS PACK! ONE FREE PATCH**

# Salonpas Pain Relief Patch

**APPLY FOR 8-12 HOURS**

**Menthol 3% Methyl Salicylate 10% Topical Analgesic**

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH  
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

**6/5 patches**  
2 1/4" x 3 15/16" (7cm x 10cm)

Package not child resistant.  
Keep out of reach of children.

**DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS**

LOT \_\_\_\_\_  
EXP. \_\_\_\_\_

**Salonpas Pain Relief Patch**  
Menthol 3%, Methyl Salicylate 10%

**6/5 patches**

**press open**

**Salonpas Pain Relief Patch**  
Menthol 3%, Methyl Salicylate 10%

**minty scent**

**Drug Facts**

**Active ingredients (in each patch)**

Active ingredients (in each patch)	Purpose
Menthol 3%	Topical analgesic
Methyl salicylate 10%(NSAID : nonsteroidal anti-inflammatory drug)	Topical analgesic

**Uses**  
Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:  
 ■ strains ■ sprains ■ simple backache ■ arthritis ■ bruises

**Warnings**  
For external use only

**Stomach bleeding warning:** This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:  
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]  
 ■ have 3 or more alcoholic drinks every day while using this product  
 ■ take more or for a longer time than directed

**Do not use**  
 ■ on the face or rashes ■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs  
 ■ with a heating pad ■ when sweating (such as from exercise or heat)  
 ■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

**Ask a doctor before use if**  
 ■ you are allergic to topical products ■ the stomach bleeding warning applies to you  
 ■ you are taking a diuretic ■ you have high blood pressure, heart disease, or kidney disease

**When using this product**  
 ■ wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.  
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**  
 ■ stomach pain or upset gets worse or lasts  
 ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.  
 ■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days  
 ■ symptoms clear up and occur again within a few days

**If pregnant or breast-feeding,** ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

**Directions**

**Adults 18 years and older:**  
 ■ clean and dry affected area ■ only use one patch at a time  
 ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours ■ remove patch from backing film and apply to skin (see illustration)  
 ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours  
 ■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row  
 ■ the used patch should be removed from the skin when a new one is applied

**Children under 18 years of age:** do not use; this product has not been shown to work in children

**Other information**  
 ■ store at 20-25°C (68-77°F) ■ avoid storing product in direct sunlight  
 ■ protect product from excessive moisture  
 ■ some individuals may not experience pain relief until several hours after applying the patch

**Inactive ingredients**  
 alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

**Questions or comments?**  
 Toll free 1-87-SALONPAS [www.salonpas.us](http://www.salonpas.us)

Labeling Format Information:  
 Fonts : Helvetica BoldOblique  
 Helvetica 77 Bold Condensed Oblique  
 Helvetica 57 Condensed  
 Helvetica Condensed Bold  
 Drug Facts : 9.5pt Leading : 2pt  
 Header : 8pt Bullets : 5pt  
 Subheader : 6.5pt Barlines : 2pt  
 Body Text : 6pt Hairlines : 0.4pt  
 Drug Facts (Continued) NA  
 Horizontal Scale: 80%-100% Average Kerning: 0

100%出力  
 FS67\_3s\_f\_130606

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

FOLD TWICE AFTER OPENING

Discard unused patches 14 days after opening

FOLD TWICE AFTER OPENING



**Menthol 3%  
 Methyl Salicylate 10%  
 Topical Analgesic**

**3 patches**  
 2 3/4" x 3 1/4"  
 (7cm x 10cm)

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH  
**EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL**

※ DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

**Hisamitsu®**

Manufactured by  
**Hisamitsu Pharmaceutical Co., Inc.**  
 JAPAN SAGA TOSU  
**MADE IN JAPAN**

Distributed by  
**Hisamitsu America, Inc.**  
 Florham Park, NJ 07932

Package not child resistant.  
 Keep out of reach of children.

LOT

EXP.

© : Registered Trade Mark

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂  
**Salonpas® Pain Relief Patch** Menthol 3%, Methyl Salicylate 10% minty scent

**Drug Facts**

Active ingredients (in each patch)	Purpose
Menthol 3%	Topical analgesic
Methyl salicylate 10%(NSAID : nonsteroidal anti-inflammatory drug)	Topical analgesic

**Uses** Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:  
 ■ strains ■ sprains ■ simple backache ■ arthritis ■ bruises

**Warnings**

For external use only

**Stomach bleeding warning:**

This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:  
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)  
 ■ have 3 or more alcoholic drinks every day while using this product  
 ■ take more or for a longer time than directed

**Do not use**

■ on the face or rashes ■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs  
 ■ with a heating pad ■ when sweating (such as from exercise or heat)  
 ■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

**Ask a doctor before use if**

■ you are allergic to topical products ■ the stomach bleeding warning applies to you  
 ■ you are taking a diuretic ■ you have high blood pressure, heart disease, or kidney disease

**When using this product**

■ wash hands after applying or removing patch. Avoid contact with eyes.  
 If eye contact occurs, rinse thoroughly with water.  
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

■ stomach pain or upset gets worse or lasts  
 ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.  
 ■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days  
 ■ symptoms clear up and occur again within a few days

**If pregnant or breast-feeding**, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

**Directions**

**Adults 18 years and older:**

■ only use one patch at a time  
 ■ clean and dry affected area ■ remove patch from backing film and apply to skin (see illustration)  
 ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours  
 ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours  
 ■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row  
 ■ the used patch should be removed from the skin when a new one is applied  
**Children under 18 years of age:** do not use; this product has not been shown to work in children

**Other information**

■ store at 20-25°C (68-77°F) ■ avoid storing product in direct sunlight  
 ■ protect product from excessive moisture  
 ■ some individuals may not experience pain relief until several hours after applying the patch

**Inactive ingredients** alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

**Questions or comments?**

Toll free 1-87-SALONPAS

[www.salonpas.us](http://www.salonpas.us)

1L835T-0

-----  
**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**  
-----

/s/  
-----

THERESA M MICHELE  
01/31/2014