

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other antihistamines
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better

Drug Facts (continued)

- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours
- if taken with food, this product may take longer to work

Other information



- read all warnings and directions before use. Keep out of carton.
- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients camauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments? 1-800-395-0689 (Mon - Fri 9AM - 5PM EST)

XXXXXXXX

0000



Drug Facts

Active ingredients (in each caplet)

Diphenhydramine hydrochloride 25 mg.....Nighttime sleep-aid
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*.....Pain reliever
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash

● blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed


(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts (continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39


Actual Size 100%




Actual Size 100%



BASE



FRONT COVER




INSIDE FRONT COVER


Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	-
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	-
Leading:	N/A	+0.5 pt	6.5/6.1 pt
Characters Per Inch:	N/A	No more than 39	39



Actual Size 100%



BASE

Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.7 pt
Characters Per Inch:	N/A	No more than 39	39

Drug Facts (continued)

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- your pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours
- if taken with food, this product may take longer to work

Other Information

- read all warnings and directions before use.
- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive Ingredients: carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments? 1-800-366-0880 (Mon - Fri 9AM - 5PM EST)

Bayer

Do not use this product if printed safety seal bearing "Bayer HealthCare" under cap is torn or missing.

0 00000 00000 0

Drug Facts

Active Ingredients (in each caplet)	Purpose
Diphenhydramine hydrochloride 25 mg	Nighttime sleep-aid
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever

*nonsteroidal anti-inflammatory drug

FRONT COVER

Drug Facts (continued)

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin redness
- facial swelling
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if:

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are:

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other antihistamines
- taking any other drug

When taking this product:

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk, if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

INSIDE FRONT COVER

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

KAREN M MAHONEY
11/20/2014