

- After REYATAZ oral powder is mixed with food or liquid it may be kept at room temperature 68°F to 86°F (20°C to 30°C) for up to 1 hour. Take REYATAZ oral powder within 1 hour after mixing with food or liquid.

Keep REYATAZ and all medicines out of the reach of children.

General information about REYATAZ

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use REYATAZ for a condition for which it was not prescribed. Do not give REYATAZ to other people, even if they have the same symptoms that you have. It may harm them. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about REYATAZ that is written for health professionals.

For more information, go to www.reyataz.com or call 1-800-321-1335.

What are the ingredients in REYATAZ?

Active ingredient: atazanavir sulfate

Inactive ingredients:

REYATAZ capsules: crospovidone, lactose monohydrate, and magnesium stearate. The capsule shells contain gelatin, FD&C Blue No. 2, titanium dioxide, black iron oxide, red iron oxide, and yellow iron oxide. The capsules are printed with ink containing shellac, titanium dioxide, FD&C Blue No. 2, isopropyl alcohol, ammonium hydroxide, propylene glycol, n-butyl alcohol, simethicone, and dehydrated alcohol.

REYATAZ oral powder: aspartame, sucrose, and orange-vanilla flavor.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Distributed by:

Bristol-Myers Squibb Company

Princeton, NJ 08543 USA

Revised: June 2014

VIDEX[®] and REYATAZ[®] are registered trademarks of Bristol-Myers Squibb Company. COUMADIN[®] and SUSTIVA[®] are registered trademarks of Bristol-Myers Squibb

