

Drug Facts (continued)

Active Ingredients (in each caplet)

Loperamide HCl 2 mg
Simethicone 125 mg

Purposes

Anti-diarrhea
Anti-gas

Uses

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- if you have difficulty swallowing
- if you have a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Other Information

Each caplet contains: loperamide 165 mg, sodium 5 mg

■ Tamper evident statement

Children under 6 years (up to 47 lbs)

ask a doctor

children 6-8 years (48-59 lbs)

1 caplet after the first loose stool, but no more than 2 caplets in 24 hours

children 9-11 years (60-95 lbs)

1 caplet after the first loose stool, but no more than 3 caplets in 24 hours

children 12 years and over

2 caplets after the first loose stool, but no more than 4 caplets in 24 hours

adults and children

2 caplets after the first loose stool, but no more than 4 caplets in 24 hours

Directions

Drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Take only on an empty stomach (1 hour before or 2 hours after a meal)

Take with a full glass of water

Do not take more than the dose on chart below. If possible, use weight to dose.

otherwise, use age.

Drug Facts (continued)

Active Ingredients (in each caplet)

Loperamide HCl 2 mg
Simethicone 125 mg

Purposes

Anti-diarrhea
Anti-gas

Uses

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- if you have difficulty swallowing
- if you have a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Other Information

Each caplet contains: loperamide 165 mg, sodium 5 mg

■ Tamper evident statement

Children under 6 years (up to 47 lbs)

ask a doctor

children 6-8 years (48-59 lbs)

1 caplet after the first loose stool, but no more than 2 caplets in 24 hours

children 9-11 years (60-95 lbs)

1 caplet after the first loose stool, but no more than 3 caplets in 24 hours

children 12 years and over

2 caplets after the first loose stool, but no more than 4 caplets in 24 hours

adults and children

2 caplets after the first loose stool, but no more than 4 caplets in 24 hours

Directions

Drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Take only on an empty stomach (1 hour before or 2 hours after a meal)

Take with a full glass of water

Do not take more than the dose on chart below. If possible, use weight to dose.

otherwise, use age.

Imodium®

How can we help?

1-877-895-3665

See New Warnings and Directions

NDC 50580-338-12

Imodium®

Loperamide HCl, 2 mg/Simethicone, 125 mg Anti-diarrhea /Anti-gas

Multi-Symptom Relief

Caplet

Relieves symptoms of diarrhea *plus*

- Cramps & Pressure
- Bloating
- Gas

12 Caplets*

*capsule-shaped tablets

Actual Size

Imodium®

Loperamide HCl, 2 mg/Simethicone, 125 mg Anti-diarrhea /Anti-gas

Multi-Symptom Relief

Caplet

Relieves symptoms of diarrhea *plus*

- Cramps & Pressure
- Bloating
- Gas

12 Caplets*

*capsule-shaped tablets

Actual Size


Loperamide HCl 2 mg/Simethicone, 125 mg Anti-diarrhea /Anti-gas
Loperamide HCl 2 mg/Simethicone, 125 mg Anti-diarrhea /Anti-gas
Loperamide HCl 2 mg/Simethicone, 125 mg Anti-diarrhea /Anti-gas
Loperamide HCl 2 mg/Simethicone, 125 mg Anti-diarrhea /Anti-gas

PDP = 10.8675 square inches

Multi-Symptom Relief

Caplet

Relieves symptoms of diarrhea *plus*



Imodium



Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas
Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas
Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas
Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas

PDP = 10.33 square inches

Relieves symptoms
of diarrhea **plus**

- Cramps & Pressure
- Bloating



Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas
Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas
Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas
Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas

Reference ID: 3718183

Fraction - Decimal Conversion Chart	
Inches	
$\frac{1}{16}$.0625
$\frac{1}{8}$.125
$\frac{3}{16}$.1875

PDP = 9.34 square inches

Relieves symptoms of diarrhea **plus**

- Cramps & Pressure
- Bloating
- Gas

42+12=54 Caplets*

*capsule-shaped tablets

Actual Size

[illegible]

Active ingredients (in each caplet) Purposes
Loperamide HCl 2 mg Anti diarrhea
Simethicone 125 mg Anti gas

Uses relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Imodium[®]
Loperamide HCl 2 mg/Simethicone 125 mg
Antidiarrheal/Anti gas

Multi-Symptom Relief

30 Caplets*
*capsule-shaped tablets

Warnings: Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.
Do not use • if you have bloody or black stool • if you have difficulty swallowing. **Ask a doctor before use if you have •** fever • mucus in the stool • a history of liver disease. **Ask a doctor or pharmacist before**

Use if you are taking antibiotics. When using this product If heartiness, drowsiness or dizziness may occur. Be careful when driving or operating machinery. **Stop use and ask a doctor if •** symptoms get worse • diarrhea lasts for more than 2 days • you get abdominal swelling or bulging. These may be signs of a serious condition. • you have difficulty swallowing the caplet. If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1 800 222 1222). **Directions •** drink plenty of clear fluids to help prevent dehydration caused by diarrhea • take only on an empty stomach (1 hour before or 2 hours after a meal) • take with a full (8 oz.) glass of water • find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9 to 11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6 to 8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours

Children under 6 years (up to 47 lbs) ask a doctor.

Other information • each caplet contains: calcium 165 mg, sodium 3 mg • store between 20-25°C (68-77°F). Protect from light • tamper evident statement. **Questions or comments?** call 1-877-895-3665 (toll free) or 215-273-8755 (in PA).
Distributed by McKE L PPC, Inc., Fort Washington, PA 19034 USA. ©MCKE L PPC, INC. 2015 US6 103,260

(b) (4)