

Drug Facts (continued)

Inactive Ingredients
 acetylsalicylic acid, croscarmellose sodium, dibasic calcium phosphate, flavor, microcrystalline cellulose, stearic acid

Questions or comments?
 call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

0045-0212-122

TEAR OFF SECTION

FOLD AND TEAR AT NOTCH

Other Information

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

If pregnant or breast-feeding, ask a health professional before use. You have difficulty swallowing the caplet.

Do not use if you have bloody or black stool.

Do not use if you have difficulty swallowing.

Ask a doctor before use if you have:

- ever had a history of liver disease
- ever had a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics.

When using this product (drowsiness, dizziness or lightheadedness may occur). Be careful when driving or operating machinery.

Stop use and ask a doctor if:

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bloating. These may be signs of a serious condition.

Do not use if you have bloody or black stool.

Do not use if you have difficulty swallowing.

Ask a doctor before use if you have:

- ever had a history of liver disease
- ever had a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics.

When using this product (drowsiness, dizziness or lightheadedness may occur). Be careful when driving or operating machinery.

Stop use and ask a doctor if:

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bloating. These may be signs of a serious condition.

Drug Facts (continued)

Directions

caused by diarrhea

drink plenty of clear fluids to help prevent dehydration

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

Do not use if you have bloody or black stool.

Do not use if you have difficulty swallowing.

Ask a doctor before use if you have:

- ever had a history of liver disease
- ever had a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics.

When using this product (drowsiness, dizziness or lightheadedness may occur). Be careful when driving or operating machinery.

Stop use and ask a doctor if:

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bloating. These may be signs of a serious condition.

Do not use if you have bloody or black stool.

Do not use if you have difficulty swallowing.

Ask a doctor before use if you have:

- ever had a history of liver disease
- ever had a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics.

When using this product (drowsiness, dizziness or lightheadedness may occur). Be careful when driving or operating machinery.

Stop use and ask a doctor if:

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bloating. These may be signs of a serious condition.

0000000

EXP

LOT

Distributed by:
 McKesson Healthcare
 Distribution
 Fort Washington, PA 19034 USA
 ©MCK PCC, INC. 2015
 www.imodium.com
 U.S. Patent No. 6,109,260

Imodium
How can we help?
1-877-895-3665

See New Warnings and Directions NDC 50580-338-12

Imodium®

Loperamide HC, 2 mg/Simethicone, 125 mg Antidiarrhea /Anti-gas

Multi-Symptom Relief Caplet

Relieves symptoms of diarrhea **plus**

- Cramps & Pressure
- Bloating
- Gas

12 Caplets*

*capsule-shaped tablets Actual Size

00000000

Loperamide HC 2 mg/Simethicone, 125 mg Antidiarrhea /Anti-gas
 Loperamide HC 2 mg/Simethicone, 125 mg Antidiarrhea /Anti-gas
 Loperamide HC 2 mg/Simethicone, 125 mg Antidiarrhea /Anti-gas
 Loperamide HC 2 mg/Simethicone, 125 mg Antidiarrhea /Anti-gas


Imodium®

PDP = 10.8675 square inches

Multi-Symptom Relief Caplet

Relieves symptoms of diarrhea **plus**

Fraction - Decimal Conversion Chart	
1	Inches
0005	Inches



Imodium

Imodium®

Loperamide HC, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas

Multi-Symptom Relief Caplet

Relieves symptoms
of diarrhea **plus**

- Cramps & Pressure
- Bloating
- Gas

30 Caplets*

*capsule-shaped tablets




Actual Size


Multi-Symptom Relief 30 Caplets

Loperamide HC, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas

Imodium®



00000000



00000000

Imodium®
How can we help?
1-877-895-3665

Imodium®
Multi-Symptom Relief 30 Caplets

Care To Recycle™

Distributed by:
McNeil Consumer Healthcare
Division of McNeil, P.C., Inc.
Fort Washington, PA 19034 USA
© MCN P.C., INC., 2015
www.imodium.com
U.S. Patent No. 6,103,260

Drug Facts (continued)

Other information
■ each caplet contains calcium 165 mg,
sodium 3 mg
■ store between 20-25°C (68-77°F). Protect
from light.
■ tamper evident statement

Inactive ingredients
acetylsalicylic acid, croscarmellose
sodium, dibasic calcium phosphate, flavor,
microcrystalline cellulose, stearic acid

Questions or comments?
call **1-877-895-3665** (toll-free) or
215-273-8755 (collect)

3 0045-0212-30 6

Drug Facts

Active ingredients (in each caplet)
Loperamide HCl 2 mg
Simethicone 125 mg
Anti-diarrheal
Anti-gas

Purposes
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
In case of overdose, get medical help or contact Poison Control Center right away. (1-800-222-1222)

Directions
Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.
Take only on an empty stomach (1 hour before or 2 hours after a meal).
Take with a full (8 oz.) glass of water.
Find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children	2 caplets a ter the first loose stool, 1 caplet a ter each subsequent loose stool, but no more than 4 caplets in 24 hours
children 12 years and over	1 caplet a ter the first loose stool, 1/2 caplet a ter each subsequent loose stool, but no more than 3 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet a ter the first loose stool, 1/2 caplet a ter each subsequent loose stool, but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet a ter the first loose stool, 1/2 caplet a ter each subsequent loose stool, but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide (HCl).
Do not use if you have bloody or black stool.
Do not use if you have difficulty swallowing.

Ask a doctor before use if
■ you are taking antibiotics
■ you have difficulty swallowing
■ a history of liver disease
■ mucus in the stool
■ fever

Ask a doctor or pharmacist before use if
you are taking antibiotics

When using this product
drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if
■ symptoms get worse
■ diarrhea lasts for more than 2 days
■ you get abdominal swelling or bulging.
These may be signs of a serious condition.
You have difficulty swallowing the caplet.

00000000

Loperamide HC, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas
Loperamide HC, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas
Loperamide HC, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas
Loperamide HC, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas

Imodium®

PDP = 10.33 square inches

Reference ID: 3718183

Fraction - Decimal Conversion Chart

Inches

1/2	0.5
1/4	0.25
1/8	0.125
3/8	0.375
1/2	0.5
5/8	0.625
3/4	0.75
7/8	0.875

Relieves symptoms
of diarrhea **plus**

- Cramps & Pressure
- Bloating

BONUS! 12 More Caplets!
See New Warnings and Directions NDC 50580-338-43

Imodium®

Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas

Multi-Symptom Relief
Caplet

Relieves symptoms of diarrhea **plus**

- Cramps & Pressure
- Bloating
- Gas

42+12=
54 Caplets*

*capsule-shaped tablets



Actual Size



00000000

How can we help?
Imodium
(1-877-895-3665)

Drug Facts

Active ingredients (in each caplet)
Loperamide HCl 2 mg..... Anti-diarrheal
Simethicone 125 mg..... Anti-gas

Purposes

Uses relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use if you have bloody or black stool or if you have difficulty swallowing

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Drug Facts (continued)

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Other information

- each caplet contains: calcium 165 mg, sodium 3 mg
- store between 20-25°C (68-77°F). Protect from light.
- tamper evident statement

Drug Facts (continued)

Inactive ingredients acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, flavor, microcrystalline cellulose, stearic acid

Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)



Care To Recycle™

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
© MCN-PPC, INC. 2015
www.imodium.com
U.S. Patent No. 6,103,260



3 0045-02 12-54 2

00000000



Imodium
Loperamide HCl, 2 mg/Simethicone, 125 mg
Antidiarrheal/Anti-gas

Multi-Symptom Relief 42+12 Caplets

LOT:

EXP:

0000000

Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas
Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas
Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas
Loperamide HCl, 2 mg/Simethicone, 125mg Antidiarrheal/Anti-gas

PDP = 9.34 square inches

Reference ID: 3718183

Fraction	Decimal
1/16	.0625
1/8	.125
3/16	.1875

Relieves symptoms of diarrhea **plus**

- Cramps & Pressure
- Bloating
- Gas

42+12=
54 Caplets*

*capsule-shaped tablets Actual Size

Active ingredients in each caplet: Loperamide HCl 2 mg, Simethicone 125 mg. **Purposes:** Anti-diarrheal, Anti-gas.

Uses: relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas.

Warnings: Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl. **Do not use** if you have bloody or black stool or if you have difficulty swallowing. **Ask a doctor before use** if you have a fever, mucus in the stool, a history of liver disease, or if you are taking antibiotics. **Ask a doctor or pharmacist before use** if you are taking antibiotics. When using this product, drowsiness or dizziness may occur. Be careful when driving or operating machinery. **Stop use and ask a doctor** if symptoms get worse, diarrhea lasts for more than 2 days, you get abdominal swelling or bulging. These may be signs of a serious condition if you have difficulty swallowing the caplet. **If pregnant or breast-feeding, ask a health professional before use.** Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions:** drink plenty of clear fluids to help prevent dehydration caused by diarrhea. Take only on an empty stomach (1 hour before or 2 hours after a meal). Take with a full 8 oz glass of water. Find right dose on chart below. If possible, use weight to dose; otherwise, use age.

Other information: each caplet contains calcium 165 mg, sodium 3 mg + store between 20-25°C (68-77°F). Protect from light. Tamper evident seal. Questions or comments? call 1-877-895-3665 (ask for) or 215-273-4735 (collect). US# 103,260. Dist by MCNE LPPC, Inc., Fort Washington, PA 19054 USA. © MCN LPPC, INC 2015.

Multi-Symptom Relief
30 Caplets*
 *capsule shaped tablets

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

0000000

(b) (4)