

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use NovoLog Mix 70/30 safely and effectively. See full prescribing information for NovoLog Mix 70/30.

NovoLog® Mix 70/30 (70% insulin aspart protamine suspension and 30% insulin aspart injection, [rDNA origin])
Suspension for subcutaneous injection
Initial U.S. Approval: 2001

-----RECENT MAJOR CHANGES-----

- Warnings and Precautions (5.1) 02/2015

-----INDICATIONS AND USAGE-----

NovoLog Mix 70/30 is an insulin analog indicated to improve glycemic control in patients with diabetes mellitus.

Important Limitations of Use: In premix insulins, such as NovoLog Mix 70/30, the proportions of rapid acting and long acting insulins are fixed and do not allow for basal versus prandial dose adjustments (1).

-----DOSAGE AND ADMINISTRATION-----

- Only for subcutaneous injection (2.1).
Type 1 DM: dose within 15 minutes before meal initiation.
Type 2 DM: dose within 15 minutes before or after starting a meal.
- Do not administer intravenously (2.1).
- Do not use in insulin infusion pumps (2.1).
- Must be resuspended immediately before use (2.2).

-----DOSAGE FORMS AND STRENGTHS-----

Each presentation contains 100 Units of insulin aspart per mL (U-100) (3)

- 10 mL vials
- 3 mL NovoLog Mix 70/30 FlexPen

-----CONTRAINDICATIONS-----

- Do not use during episodes of hypoglycemia (4).
- Do not use in patients with hypersensitivity to NovoLog Mix 70/30 or one of its excipients (4).

-----WARNINGS AND PRECAUTIONS-----

- Never share a NovoLog Mix 70/30 FlexPen between patients, even if the needle is changed (5.1).
- NovoLog Mix 70/30 should not be mixed with any other insulin product (5.2).
- Hypoglycemia is the most common adverse effect of insulin therapy. Glucose monitoring is recommended for all patients with diabetes. Any change of insulin dose should be made cautiously and only under medical supervision (5.2, 5.3).

- Insulin, particularly when given in settings of poor glycemic control, can cause hypokalemia. Use caution in patients predisposed to hypokalemia (5.4).
- Like all insulins, NovoLog Mix 70/30 requirements may be reduced in patients with renal impairment or hepatic impairment (5.5, 5.6).
- Severe, life-threatening, generalized allergy, including anaphylaxis, may occur with insulin products, including NovoLog Mix 70/30 (5.7).
- Fluid retention and heart failure can occur with concomitant use of thiazolidinediones (TZDs), which are PPAR-gamma agonists, and insulin, including NovoLog Mix 70/30 (5.9).

-----ADVERSE REACTIONS-----

Adverse reactions observed with insulin therapy include hypoglycemia, allergic reactions, local injection site reactions, lipodystrophy, rash and pruritus (6).

To report SUSPECTED ADVERSE REACTIONS, contact Novo Nordisk Inc. at 1-800-727-6500 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----DRUG INTERACTIONS-----

- The following may increase the blood-glucose-lowering effect and susceptibility to hypoglycemia: oral antidiabetic products, pramlintide, ACE inhibitors, disopyramide, fibrates, fluoxetine, monoamine oxidase (MAO) inhibitors, propoxyphene, salicylates, somatostatin analog (e.g. octreotide), sulfonamide antibiotics (7).
- The following may reduce the blood-glucose-lowering effect: corticosteroids, niacin, danazol, diuretics, sympathomimetic agents (e.g., epinephrine, salbutamol, terbutaline), isoniazid, phenothiazine derivatives, somatropin, thyroid hormones, estrogens, progestogens (e.g., in oral contraceptives), atypical antipsychotics (7).
- Beta-blockers, clonidine, lithium salts, and alcohol may either potentiate or weaken the blood-glucose-lowering effect of insulin (7).
- Pentamidine may cause hypoglycemia, which may be followed by hyperglycemia (7).
- The signs of hypoglycemia may be reduced or absent in patients taking sympatholytic products such as beta-blockers, clonidine, guanethidine, and reserpine (7).

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 02/2015

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosing
- 2.2 Resuspension

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Never Share a NovoLog Mix 70/30 FlexPen Between Patients
- 5.2 Administration
- 5.3 Hypoglycemia
- 5.4 Hypokalemia
- 5.5 Renal Impairment
- 5.6 Hepatic Impairment
- 5.7 Hypersensitivity and Allergic Reactions
- 5.8 Antibody Production
- 5.9 Fluid retention and heart failure can occur with concomitant use of PPAR- gamma agonists

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use

8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES

- 14.1 NovoLog Mix 70/30 versus Novolin 70/30
- 14.2 Combination Therapy: Insulin and Oral Agents in Patients with Type 2 Diabetes

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Recommended Storage

17 PATIENT COUNSELING INFORMATION

- 17.1 Never Share a NovoLog Mix 70/30 FlexPen Between Patients
- 17.2 Physician Instructions

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

NovoLog Mix 70/30 is an insulin analog indicated to improve glycemic control in patients with diabetes mellitus.

Important Limitations of Use:

In premix insulins, such as NovoLog Mix 70/30, the proportions of rapid acting and long acting insulins are fixed and do not allow for basal versus prandial dose adjustments.

2 DOSAGE AND ADMINISTRATION

2.1 Dosing

NovoLog Mix 70/30 is an insulin analog with an earlier onset and intermediate duration of action in comparison to the basal human insulin premix. The addition of protamine to the rapid-acting aspart insulin analog (NovoLog) results in insulin activity that is 30% short-acting and 70% long-acting. NovoLog Mix 70/30 is typically dosed on a twice-daily basis (with each dose intended to cover 2 meals or a meal and a snack). The dosage of NovoLog Mix 70/30 must be individualized. The written prescription for NovoLog Mix 70/30 should include the full name, to avoid confusion with NovoLog (insulin aspart) and Novolin 70/30 (human premix).

NovoLog Mix 70/30 should appear uniformly white and cloudy. Do not use it if it looks clear or if it contains solid particles. NovoLog Mix 70/30 should not be used after the printed expiration date.

NovoLog Mix 70/30 should be administered by subcutaneous injection in the abdominal region, buttocks, thigh, or upper arm. NovoLog Mix 70/30 has a faster onset of action than human insulin premix 70/30 and should be dosed within 15 minutes before meal initiation for patients with type 1 diabetes. For patients with type 2 diabetes, dosing should occur within 15 minutes before or after meal initiation. Injection sites should be rotated within the same region to reduce the risk of lipodystrophy. As with all insulins, the duration of action may vary according to the dose, injection site, blood flow, temperature, and level of physical activity.

NovoLog Mix 70/30 should not be administered intravenously or used in insulin infusion pumps. Dose regimens of NovoLog Mix 70/30 will vary among patients and should be determined by the health care professional familiar with the patient's recommended glucose treatment goals, metabolic needs, eating habits, and other lifestyle variables.

2.2 Resuspension

NovoLog Mix 70/30 is a suspension that must be visually inspected and resuspended immediately before use.

The NovoLog Mix 70/30 vial should be rolled gently in your hands in a horizontal position 10 times to mix it. The rolling procedure must be repeated until the suspension appears uniformly white and cloudy. Inject immediately. Resuspension is easier when the insulin has reached room temperature.

The NovoLog Mix 70/30 FlexPen should be rolled 10 times gently between your hands in a horizontal position. Thereafter, turn the NovoLog Mix 70/30 FlexPen upside down so that the glass ball moves from one end of the reservoir to the other. Do this at least 10 times. The rolling and turning procedure must be repeated until the suspension appears uniformly white and cloudy. Inject immediately. Before each subsequent injection, turn the disposable NovoLog Mix 70/30 FlexPen upside down so that the glass ball moves from one end of the reservoir to the other at least 10 times and until the suspension appears uniformly white and cloudy. Inject immediately.

3 DOSAGE FORMS AND STRENGTHS

NovoLog Mix 70/30 is available in the following package sizes: each presentation contains 100 units of insulin aspart per mL (U-100).

- 10 mL vials
- 3 mL NovoLog Mix 70/30 FlexPen

4 CONTRAINDICATIONS

NovoLog Mix 70/30 is contraindicated

- during episodes of hypoglycemia
- in patients with hypersensitivity to NovoLog Mix 70/30 or one of its excipients.

5 WARNINGS AND PRECAUTIONS

5.1 Never Share a NovoLog Mix 70/30 FlexPen Between Patients

NovoLog Mix 70/30 FlexPens must never be shared between patients, even if the needle is changed. Pen-sharing poses a risk for transmission of blood-borne pathogens.

5.2 Administration

The short and long-acting components of insulin mixes, including NovoLog Mix 70/30, cannot be titrated independently. Because NovoLog Mix 70/30 has peak pharmacodynamic activity between 1-4 hours after injection, it should be administered within 15 minutes of meal initiation [see *Clinical Pharmacology* (12)]. The dose of insulin required to provide adequate glycemic control for one of the meals may result in hyper- or hypoglycemia for the other meal. The pharmacodynamic profile may also be inadequate for patients who require more frequent meals.

NovoLog Mix 70/30 should not be mixed with any other insulin product.

NovoLog Mix 70/30 should not be used intravenously.

NovoLog Mix 70/30 should not be used in insulin infusion pumps.

Glucose monitoring is recommended for all patients with diabetes. Any change of insulin dose should be made cautiously and only under medical supervision. Changing from one insulin product to another or changing the insulin strength may result in the need for a change in dosage. Changes may also be necessary during illness, emotional stress, and other physiologic stress in addition to changes in meals and exercise.

The pharmacokinetic and pharmacodynamic profiles of all insulins may be altered by the site used for injection and the degree of vascularization of the site. Smoking, temperature,

and exercise contribute to variations in blood flow and insulin absorption. These and other factors contribute to inter- and intra-patient variability.

5.3 Hypoglycemia

Hypoglycemia is the most common adverse effect of insulin therapy, including NovoLog Mix 70/30. Severe hypoglycemia may lead to unconsciousness and/or convulsions and may result in temporary or permanent impairment of brain function or even death. Severe hypoglycemia requiring the assistance of another person and/or parenteral glucose infusion or glucagon administration has been observed in clinical trials with insulin, including trials with NovoLog Mix 70/30.

The timing of hypoglycemia may reflect the time-action profile of the insulin formulation [see *Clinical Pharmacology* (12)]. Other factors, such as changes in dietary intake (e.g., amount of food or timing of meals), injection site, exercise, and concomitant medications may also alter the risk of hypoglycemia [see *Drug Interactions* (7)]. As with all insulins, use caution in patients with hypoglycemia unawareness and in patients who may be predisposed to hypoglycemia (e.g. patients who are fasting or have erratic food intake). The patient's ability to concentrate and react may be impaired as a result of hypoglycemia. This may present a risk in situations where these abilities are especially important, such as driving or operating machinery.

Rapid changes in serum glucose levels may induce symptoms of hypoglycemia in persons with diabetes, regardless of the glucose value. Early warning symptoms of hypoglycemia may be different or less pronounced under certain conditions, such as long duration of diabetes, diabetic nerve disease, use of medications such as beta-blockers, or intensified diabetes control [see *Drug Interactions* (7)].

5.4 Hypokalemia

All insulin products, including NovoLog Mix 70/30, cause a shift in potassium from the extracellular to intracellular space, possibly leading to hypokalemia that, if left untreated, may cause respiratory paralysis, ventricular arrhythmia, and death. Use caution in patients who may be at risk for hypokalemia (e.g. patients using potassium-lowering medications or patients taking medications sensitive to potassium concentrations).

5.5 Renal Impairment

Clinical or pharmacology studies with NovoLog Mix 70/30 in diabetic patients with various degrees of renal impairment have not been conducted. As with other insulins, the requirements for NovoLog Mix 70/30 may be reduced in patients with renal impairment [see *Clinical Pharmacology* (12.3)].

5.6 Hepatic Impairment

Clinical or pharmacology studies with NovoLog Mix 70/30 in diabetic patients with various degrees of hepatic impairment have not been conducted. As with other insulins, the requirements for NovoLog Mix 70/30 may be reduced in patients with hepatic impairment [see *Clinical Pharmacology* (12.3)].

5.7 Hypersensitivity and Allergic Reactions

Local Reactions- As with other insulin therapy, patients may experience reactions such as erythema, edema or pruritus at the site of NovoLog Mix 70/30 injection. These reactions usually resolve in a few days to a few weeks, but in some occasions, may require discontinuation of NovoLog Mix 70/30. In some instances, these reactions may be related to the insulin molecule, other components in the insulin preparation including protamine and cresol, components in skin cleansing agents, or injection techniques. Localized reactions and generalized myalgias have been reported with the use of cresol as an injectable excipient.

Systemic Reactions- Less common, but potentially more serious, is generalized allergy to insulin, which may cause rash (including pruritus) over the whole body, shortness of breath, wheezing, reduction in blood pressure, rapid pulse, or sweating. Severe cases of generalized allergy, including anaphylactic reaction, may be life threatening.

5.8 Antibody Production

Specific anti-insulin antibodies as well as cross-reacting anti-insulin antibodies were monitored in a 3-month, open-label comparator trial as well as in a long-term extension trial. Changes in cross-reactive antibodies were more common after NovoLog Mix 70/30 than with Novolin 70/30 but these changes did not correlate with change in HbA1c or increase in insulin dose. The clinical significance of these antibodies has not been established. Antibodies did not increase further after long-term exposure (>6 months) to NovoLog Mix 70/30.

5.9 Fluid retention and heart failure can occur with concomitant use of PPAR-gamma agonists

Thiazolidinediones (TZDs), which are peroxisome proliferator-activated receptor (PPAR)-gamma agonists, can cause dose-related fluid retention, particularly when used in combination with insulin. Fluid retention may lead to or exacerbate heart failure. Patients treated with insulin, including NovoLog Mix 70/30, and a PPAR-gamma agonist should be observed for signs and symptoms of heart failure. If heart failure develops, it should be managed according to current standards of care, and discontinuation or dose reduction of the PPAR-gamma agonist must be considered.

6 ADVERSE REACTIONS

Clinical Trial Experience

Clinical trials are conducted under widely varying designs, therefore, the adverse reaction rates reported in one clinical trial may not be easily compared to those rates reported in another clinical trial, and may not reflect the rates actually observed in clinical practice.

- Hypoglycemia
Hypoglycemia is the most commonly observed adverse reaction in patients using insulin, including NovoLog Mix 70/30 [see *Warnings and Precautions* (5.3)]. NovoLog Mix 70/30 should not be used during episodes of hypoglycemia [see *Contraindications* (4) and *Warnings and Precautions* (5)].
- Insulin initiation and glucose control intensification
Intensification or rapid improvement in glucose control has been associated with transitory, reversible ophthalmologic refraction disorder, worsening of diabetic retinopathy, and acute painful peripheral neuropathy. However, long-term glycemic control decreases the risk of diabetic retinopathy and neuropathy.

- *Lipodystrophy*
Long-term use of insulin, including NovoLog Mix 70/30, can cause lipodystrophy at the site of repeated insulin injections. Lipodystrophy includes lipohypertrophy (thickening of adipose tissue) and lipoatrophy (thinning of adipose tissue), and may affect insulin absorption. Rotate insulin injection sites within the same region to reduce the risk of lipodystrophy.
- *Weight gain*
Weight gain can occur with some insulin therapies, including NovoLog Mix 70/30, and has been attributed to the anabolic effects of insulin and the decrease in glycosuria.
- *Peripheral Edema*
Insulin may cause sodium retention and edema, particularly if previously poor metabolic control is improved by intensified insulin therapy.
- *Frequencies of adverse drug reactions*
The frequencies of adverse drug reactions during a clinical trial with NovoLog Mix 70/30 in patients with type 1 diabetes mellitus and type 2 diabetes mellitus are listed in the tables below. The trial was a three-month, open-label trial in patients with type 1 or type 2 diabetes who were treated twice daily (before breakfast and before supper) with NovoLog Mix 70/30.

Table 1: Treatment-Emergent Adverse Events in Patients with Type 1 diabetes mellitus (Adverse events with frequency \geq 5% are included.)

	NovoLog Mix 70/30 (N=55)		Novolin 70/30 (N=49)	
Preferred Term	N	%	N	%
Hypoglycemia	38	69	37	76
Headache	19	35	6	12
Influenza-like symptoms	7	13	1	2
Dyspepsia	5	9	3	6
Back pain	4	7	2	4
Diarrhea	4	7	3	6
Pharyngitis	4	7	1	2
Rhinitis	3	5	6	12
Skeletal pain	3	5	2	4
Upper respiratory tract infection	3	5	1	2

Table 2: Treatment-Emergent Adverse Events in Patients with Type 2 diabetes mellitus (Adverse events with frequency \geq 5% are included.)

	NovoLog Mix 70/30 (N=85)		Novolin 70/30 (N=102)	
Preferred Term	N	%	N	%
Hypoglycemia	40	47	51	50
Upper respiratory tract infection	10	12	6	6
Headache	8	9	8	8
Diarrhea	7	8	2	2
Neuropathy	7	8	2	2
Pharyngitis	5	6	4	4

Abdominal pain	4	5	0	0
Rhinitis	4	5	2	2

Postmarketing Data

Additional adverse reactions have been identified during post-approval use of NovoLog Mix 70/30. Because these adverse reactions are reported voluntarily from a population of uncertain size, it is generally not possible to reliably estimate their frequency. They include medication errors in which other insulins have been accidentally substituted for NovoLog Mix 70/30 [see *Patient Counseling Information* (17)].

7 DRUG INTERACTIONS

A number of substances affect glucose metabolism and may require insulin dose adjustment and particularly close monitoring.

- The following are examples of substances that may increase the blood-glucose-lowering effect and susceptibility to hypoglycemia: oral antidiabetic products, pramlintide, ACE inhibitors, disopyramide, fibrates, fluoxetine, monoamine oxidase (MAO) inhibitors, propoxyphene, salicylates, somatostatin analog (e.g. octreotide), sulfonamide antibiotics.
- The following are examples of substances that may reduce the blood-glucose-lowering effect: corticosteroids, niacin, danazol, diuretics, sympathomimetic agents (e.g., epinephrine, salbutamol, terbutaline), isoniazid, phenothiazine derivatives, somatropin, thyroid hormones, estrogens, progestogens (e.g., in oral contraceptives), atypical antipsychotics.
- Beta-blockers, clonidine, lithium salts, and alcohol may either potentiate or weaken the blood-glucose-lowering effect of insulin.
- Pentamidine may cause hypoglycemia, which may sometimes be followed by hyperglycemia.
- The signs of hypoglycemia may be reduced or absent in patients taking sympatholytic products such as beta-blockers, clonidine, guanethidine, and reserpine.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category B.

All pregnancies have a background risk of birth defects, loss, or other adverse outcome regardless of drug exposure. This background risk is increased in pregnancies complicated by hyperglycemia and may be decreased with good metabolic control. It is essential for patients with diabetes or history of gestational diabetes to maintain good metabolic control before conception and throughout pregnancy. Insulin requirements may decrease during the first trimester, generally increase during the second and third trimesters, and rapidly decline after delivery. Careful monitoring of glucose control is essential in such patients.

An open-label, randomized study compared the safety and efficacy of NovoLog (the rapid-acting component of NovoLog Mix 70/30) versus human insulin in the treatment of

pregnant women with type 1 diabetes (322 exposed pregnancies (NovoLog: 157, human insulin: 165)). Two-thirds of the enrolled patients were already pregnant when they entered the study. Since only one-third of the patients enrolled before conception, the study was not large enough to evaluate the risk of congenital malformations. Mean HbA_{1c} of ~ 6% was observed in both groups during pregnancy, and there was no significant difference in the incidence of maternal hypoglycemia.

Animal reproduction studies have not been conducted with NovoLog Mix 70/30. However, subcutaneous reproduction and teratology studies have been performed with NovoLog (the rapid-acting component of NovoLog Mix 70/30) and regular human insulin in rats and rabbits. In these studies, NovoLog was given to female rats before mating, during mating, and throughout pregnancy, and to rabbits during organogenesis. The effects of NovoLog did not differ from those observed with subcutaneous regular human insulin. NovoLog, like human insulin, caused pre- and post-implantation losses and visceral/skeletal abnormalities in rats at a dose of 200 U/kg/day (approximately 32-times the human subcutaneous dose of 1.0 U/kg/day, based on U/body surface area), and in rabbits at a dose of 10 U/kg/day (approximately three times the human subcutaneous dose of 1.0 U/kg/day, based on U/body surface area). The effects are probably secondary to maternal hypoglycemia at high doses. No significant effects were observed in rats at a dose of 50 U/kg/day and rabbits at a dose of 3 U/kg/day. These doses are approximately 8 times the human subcutaneous dose of 1.0 U/kg/day for rats and equal to the human subcutaneous dose of 1.0 U/kg/day for rabbits based on U/body surface area.

Female patients should be advised to discuss with their physician if they intend to, or if they become pregnant. There are no adequate and well-controlled studies of the use of NovoLog Mix 70/30 in pregnant women.

8.3 Nursing Mothers

It is unknown whether insulin aspart is excreted in human milk as occurs with human insulin. There are no adequate and well-controlled studies of the use of NovoLog Mix 70/30 or NovoLog in lactating women. Women with diabetes who are lactating may require adjustments of their insulin doses.

8.4 Pediatric Use

Safety and effectiveness of NovoLog Mix 70/30 have not been established in pediatric patients.

8.5 Geriatric Use

Clinical studies of NovoLog Mix 70/30 did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently than younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy in this population.

10 OVERDOSAGE

Hypoglycemia may occur as a result of an excess of insulin relative to food intake, energy expenditure, or both. Mild episodes of hypoglycemia usually can be treated with oral glucose. Adjustments in drug dosage, meal patterns, or exercise, may be needed. More severe

episodes with coma, seizure, or neurologic impairment may be treated with intramuscular/subcutaneous glucagon or concentrated intravenous glucose. Sustained carbohydrate intake and observation may be necessary because hypoglycemia may recur after apparent clinical recovery.

11 DESCRIPTION

NovoLog Mix 70/30 (70% insulin aspart protamine suspension and 30% insulin aspart injection, [rDNA origin]) is a human insulin analog suspension containing 70% insulin aspart protamine crystals and 30% soluble insulin aspart. NovoLog Mix 70/30 is a blood-glucose-lowering agent with an earlier onset and an intermediate duration of action. Insulin aspart is homologous with regular human insulin with the exception of a single substitution of the amino acid proline by aspartic acid in position B28, and is produced by recombinant DNA technology utilizing *Saccharomyces cerevisiae* (baker's yeast). Insulin aspart (NovoLog) has the empirical formula $C_{256}H_{381}N_{65}O_{79}S_6$ and a molecular weight of 5825.8 Da.




Figure 1. Structural formula of insulin aspart

NovoLog Mix 70/30 is a uniform, white, sterile suspension that contains insulin aspart 100 Units/mL.

Inactive ingredients are glycerol 16.0 mg/mL, phenol 1.50 mg/mL, metacresol 1.72 mg/mL, zinc 19.6 µg/mL, disodium hydrogen phosphate dihydrate 1.25 mg/mL, sodium chloride 0.877 mg/mL, and protamine sulfate 0.32 mg/mL. NovoLog Mix 70/30 has a pH of 7.20 - 7.44. Hydrochloric acid or sodium hydroxide may be added to adjust pH.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The primary activity of NovoLog Mix 70/30 is the regulation of glucose metabolism. Insulins, including NovoLog Mix 70/30, bind to the insulin receptors on muscle, liver and fat cells and lower blood glucose by facilitating the cellular uptake of glucose and simultaneously inhibiting the output of glucose from the liver.

12.2 Pharmacodynamics

The two euglycemic clamp studies described below [see *Clinical Pharmacology* (12.3)] assessed glucose utilization after dosing of healthy volunteers. NovoLog Mix 70/30 has an earlier onset of action than human premix 70/30 in studies of normal volunteers and patients with diabetes. The onset of action is between 10-20 minutes for NovoLog Mix 70/30 compared

to 30 minutes for Novolin 70/30. The mean \pm SD time to peak activity for NovoLog Mix 70/30 is 2.4 hr \pm 0.8 hr compared to 4.2 hr \pm 0.4 hr for Novolin 70/30. The duration of action may be as long as 24 hours (see Figure 2).




Figure 2. Pharmacodynamic Activity Profile of NovoLog Mix 70/30 and Novolin 70/30 in healthy subjects.

12.3 Pharmacokinetics

The single substitution of the amino acid proline with aspartic acid at position B28 in insulin aspart (NovoLog) reduces the molecule's tendency to form hexamers as observed with regular human insulin. The rapid absorption characteristics of NovoLog are maintained by NovoLog Mix 70/30. The insulin aspart in the soluble component of NovoLog Mix 70/30 is absorbed more rapidly from the subcutaneous layer than regular human insulin. The remaining 70% is in crystalline form as insulin aspart protamine which has a prolonged absorption profile after subcutaneous injection.

Bioavailability and Absorption- The relative bioavailability of NovoLog Mix 70/30 compared to NovoLog and Novolin 70/30 indicates that the insulins are absorbed to similar extent. In euglycemic clamp studies in healthy volunteers (n=23) after dosing with NovoLog Mix 70/30 (0.2 U/kg), a mean maximum serum concentration (C_{max}) of 23.4 ± 5.3 mU/L was reached after 60 minutes. The mean half-life ($t_{1/2}$) of NovoLog Mix 70/30 was about 8 to 9 hours. Serum insulin levels returned to baseline 15 to 18 hours after a subcutaneous dose of NovoLog Mix 70/30. Similar data were seen in a separate euglycemic clamp study in healthy volunteers (n=24) after dosing with NovoLog Mix 70/30 (0.3 U/kg). A C_{max} of 61.3 ± 20.1 mU/L was reached after 85 minutes. Serum insulin levels returned to baseline 12 hours after a subcutaneous dose.

The C_{\max} and the area under the insulin concentration-time curve (AUC) after administration of NovoLog Mix 70/30 was approximately 20% greater than those after administration of Novolin 70/30, (see Fig. 3 for pharmacokinetic profiles).




Figure 3. Pharmacokinetic Profiles of NovoLog Mix 70/30 and Novolin 70/30

Distribution and Elimination- NovoLog has a low binding to plasma proteins, 0 to 9%, similar to regular human insulin. After subcutaneous administration in normal male volunteers (n=24), NovoLog was more rapidly eliminated than regular human insulin with an average apparent half-life of 81 minutes compared to 141 minutes for regular human insulin.

The effect of sex, age, obesity, ethnic origin, renal and hepatic impairment, pregnancy, or smoking, on the pharmacodynamics and pharmacokinetics of NovoLog Mix 70/30 has not been studied.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Standard 2-year carcinogenicity studies in animals have not been performed to evaluate the carcinogenic potential of NovoLog Mix 70/30. In 52-week studies, Sprague-Dawley rats were dosed subcutaneously with NovoLog, the rapid-acting component of NovoLog Mix 70/30, at 10, 50, and 200 U/kg/day (approximately 2, 8, and 32 times the human subcutaneous dose of 1.0 U/kg/day, based on U/body surface area, respectively). At a dose of 200 U/kg/day, NovoLog increased the incidence of mammary gland tumors in females when compared to untreated controls. The incidence of mammary tumors found with NovoLog was not

significantly different from that found with regular human insulin. The relevance of these findings to humans is not known.

NovoLog was not genotoxic in the following tests: Ames test, mouse lymphoma cell forward gene mutation test, human peripheral blood lymphocyte chromosome aberration test, *in vivo* micronucleus test in mice, and in *ex vivo* UDS test in rat liver hepatocytes.

In fertility studies in male and female rats, NovoLog at subcutaneous doses up to 200 U/kg/day (approximately 32 times the human subcutaneous dose, based on U/body surface area) had no direct adverse effects on male and female fertility, or on general reproductive performance of animals.

13.2 Animal Toxicology and/or Pharmacology

In standard biological assays in mice and rabbits, one unit of NovoLog has the same glucose-lowering effect as one unit of regular human insulin. However, the effect of NovoLog Mix 70/30 is more rapid in onset compared to Novolin (human insulin) 70/30 due to its faster absorption after subcutaneous injection.

14 CLINICAL STUDIES

14.1 NovoLog Mix 70/30 versus Novolin 70/30

In a three-month, open-label trial, patients with type 1 (n=104) or type 2 (n=187) diabetes were treated twice daily (before breakfast and before supper) with NovoLog Mix 70/30 or Novolin 70/30. Patients had received insulin for at least 24 months before the study. Oral hypoglycemic agents were not allowed within 1 month prior to the study or during the study. The small changes in HbA_{1c} were comparable across the treatment groups (see Table 3).

Table 3: Glycemic Parameters at the End of Treatment [Mean ± SD (N subjects)]

	NovoLog Mix 70/30	Novolin 70/30
Type 1, N=104		
Fasting Blood Glucose (mg/dL)	174 ± 64 (48)	142 ± 59 (44)
1.5 Hour Post Breakfast (mg/dL)	187 ± 82 (48)	200 ± 82 (42)
1.5 Hour Post Dinner (mg/dL)	162 ± 77 (47)	171 ± 66 (41)
HbA _{1c} (%) Baseline	8.4 ± 1.2 (51)	8.5 ± 1.1 (46)
HbA _{1c} (%) Week 12	8.4 ± 1.1 (51)	8.3 ± 1.0 (47)
Type 2, N=187		
Fasting Blood Glucose (mg/dL)	153 ± 40 (76)	152 ± 69 (93)
1.5 Hour Post Breakfast (mg/dL)	182 ± 65 (75)	200 ± 80 (92)
1.5 Hour Post Dinner (mg/dL)	168 ± 51 (75)	191 ± 65 (93)
HbA _{1c} (%) Baseline	8.1 ± 1.2 (82)	8.2 ± 1.3 (98)
HbA _{1c} (%) Week 12	7.9 ± 1.0 (81)	8.1 ± 1.1 (96)

The significance, with respect to the long-term clinical sequelae of diabetes, of the differences in postprandial hyperglycemia between treatment groups has not been established.

Specific anti-insulin antibodies as well as cross-reacting anti-insulin antibodies were monitored in the 3-month, open-label comparator trial as well as in a long-term extension trial.

14.2 Combination Therapy: Insulin and Oral Agents in Patients with Type 2 Diabetes

Trial 1:

In a 34-week, open-label trial, insulin-naïve patients with type 2 diabetes currently treated with 2 oral antidiabetic agents were switched to treatment with metformin and pioglitazone. During an 8-week optimization period metformin and pioglitazone were increased to 2500 mg per day and 30 or 45 mg per day, respectively. After the optimization period, subjects were randomized to receive either NovoLog Mix 70/30 twice daily added on to the metformin and pioglitazone regimen or continue the current optimized metformin and pioglitazone therapy. NovoLog Mix 70/30 was started at a dose of 6 IU twice daily (before breakfast and before supper). Insulin doses were titrated to a pre-meal glucose goal of 80-110 mg/dL. The total daily insulin dose at the end of the study was 56.9 ± 30.5 IU.

Table 4: Combination Therapy with Oral Agents and Insulin in Patients with Type 2 Diabetes Mellitus [Mean (SD)]

Treatment duration 24-weeks	NovoLog Mix 70/30 + Metformin + Pioglitazone	Metformin + Pioglitazone
HbA _{1c}		
Baseline mean \pm SD (n)	8.1 ± 1.0 (102)	8.1 ± 1.0 (98)
End-of-study mean \pm SD (n) - LOCF	6.6 ± 1.0 (93)	7.8 ± 1.2 (87)
Adjusted Mean change from baseline \pm SE (n)*	-1.6 ± 0.1 (93)	-0.3 ± 0.1 (87)
Treatment difference mean \pm SE* 95% CI*	-1.3 ± 0.1 (-1.6, -1.0)	
Percentage of subjects reaching HbA _{1c} <7.0%	76%	24%
Percentage of subjects reaching HbA _{1c} \leq 6.5%	59%	12%
Fasting Blood Glucose (mg/dL)		
Baseline Mean \pm SD (n)	173 ± 39.8 (93)	163 ± 35.4 (88)
End of Study Mean \pm SD (n) – LOCF	130 ± 50.0 (90)	162 ± 40.8 (84)
Adjusted Mean change from baseline \pm SE (n)*	-43.0 ± 5.3 (90)	-3.9 ± 5.3 (84)
End-of-Study Blood Glucose (Plasma) (mg/dL)		
2 Hour Post Breakfast	138 ± 42.8 (86)	188 ± 57.7 (74)
2 Hour Post Lunch	150 ± 41.5 (86)	176 ± 56.5 (74)
2 Hour Post Dinner	141 ± 57.8 (86)	195 ± 60.1 (74)
% of patients with severe hypoglycemia**	3	0
% of patients with minor hypoglycemia**	52	3
Weight gain at end of study (kg)**	4.6 ± 4.3 (92)	0.8 ± 3.2 (86)

*Adjusted mean per group, treatment difference, and 95% CI were obtained based on an ANCOVA model with treatment, FPG stratum, and secretagogue stratum as fixed factors and baseline HbA_{1c} as the covariate.

**If metabolic control is improved by intensified insulin therapy, an increased risk of hypoglycemia and weight gain may occur.

Trial 2:

In a 28-week, open-label trial, insulin-naïve patients with type 2 diabetes with fasting plasma glucose above 140 mg/dL currently treated with metformin \pm thiazolidinedione therapy were randomized to receive either NovoLog Mix 70/30 twice daily [before breakfast and before supper] or insulin glargine once daily¹ (see Table 5). NovoLog Mix 70/30 was started at an

average dose of 5-6 IU (0.07 ± 0.03 IU/kg) twice daily (before breakfast and before supper), and bedtime insulin glargine was started at 10-12 IU (0.13 ± 0.03 IU/kg). Insulin doses were titrated weekly by decrements or increments of -2 to +6 units per injection to a pre-meal glucose goal of 80-110 mg/dL. The metformin dose was adjusted to 2550 mg/day. Approximately one-third of the patients in each group were also treated with pioglitazone (30 mg/day). Insulin secretagogues were discontinued in order to reduce the risk of hypoglycemia. Most patients were Caucasian (53%), and the mean initial weight was 90 kg.

Table 5: Combination Therapy with Oral Agents and Two Types of Insulin in Patients with Type 2 Diabetes Mellitus [Mean (SD)]

Treatment duration 28-weeks	NovoLog Mix 70/30 + Metformin ± Pioglitazone	Insulin Glargine + Metformin ± Pioglitazone
Number of patients	117	116
HbA _{1c}		
Baseline mean (%)	9.7 ± 1.5 (117)	9.8 ± 1.4 (114)
End-of-study mean (\pm SD)	6.9 ± 1.2 (108)	7.4 ± 1.2 (114)
Mean change from baseline	-2.7 ± 1.6 (108)	-2.4 ± 1.5 (114)
Percentage of subjects reaching HbA _{1c} <7.0%	66%	40%
Total Daily Insulin Dose at end of study (U)	78 ± 40 (117)	51 ± 27 (116)
% of patients with severe hypoglycemia	0	0
% of minor hypoglycemia	43	16
Weight gain at end of study	5.4 ± 4.8 (117)	3.5 ± 4.5 (116)

15 REFERENCES

1. Raskin R, Allen E, Hollander P, et al. Initiating insulin therapy in type 2 diabetes: a comparison of biphasic and basal insulin analogs. *Diabetes Care*. 2005; 28:260-265.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

NovoLog Mix 70/30 is available in the following package sizes: each presentation contains 100 Units of insulin aspart per mL (U-100).

10 mL vials NDC 0169-3685-12
3 mL NovoLog Mix 70/30 FlexPen NDC 0169-3696-19

NovoLog Mix 70/30 vials and NovoLog Mix 70/30 FlexPen are latex free. Novolog Mix 70/30 FlexPens must never be shared between patients, even if the needle is changed.

16.2 Recommended Storage

Unused NovoLog Mix 70/30 should be stored in a refrigerator between 2°C and 8°C (36°F to 46°F). Do not store in the freezer or directly adjacent to the refrigerator cooling element. **Do not freeze NovoLog Mix 70/30 or use NovoLog Mix 70/30 if it has been frozen.**

Vials: After initial use, a vial may be kept at temperatures below 30°C (86°F) for up to 28 days, but should not be exposed to excessive heat or sunlight. Open vials may be refrigerated.

Unpunctured vials can be used until the expiration date printed on the label if they are stored in a refrigerator. Keep unused vials in the carton so they will stay clean and protected from light.

NovoLog Mix 70/30 FlexPen: Once a NovoLog Mix 70/30 FlexPen is punctured, it should be kept at temperatures below 30°C (86°F) for up to 14 days, but should not be exposed to excessive heat or sunlight. A NovoLog Mix 70/30 FlexPen in use must NOT be stored in the refrigerator. Keep the disposable NovoLog Mix 70/30 FlexPen away from direct heat and sunlight. An unpunctured NovoLog Mix 70/30 FlexPen can be used until the expiration date printed on the label if they are stored in a refrigerator. Keep any unused NovoLog Mix 70/30 FlexPen in the carton so it will stay clean and protected from light.

Always remove the needle after each injection and store NovoLogMix 70/30 FlexPen without a needle attached. This prevents contamination and/or infection, or leakage of insulin, and will ensure accurate dosing. Always use a new needle for each injection to prevent contamination.

These storage conditions are summarized in the following table:

	Not in-use (unopened) Room Temperature (below 30° C [86° F])	Not in-use (unopened) Refrigerated (2° C - 8° C [36° F - 46° F])	In-use (opened) Room Temperature (below 30° C [86° F])
10 mL vial	28 days	Until expiration date	28 days (refrigerated/room temperature)
3 mL NovoLog Mix 70/30 FlexPen	14 days	Until expiration date	14 days (Do not refrigerate)

17 PATIENT COUNSELING INFORMATION

[see FDA-Approved Patient Labeling]

17.1 Never Share a NovoLog Mix 70/30 FlexPen Between Patients

Advise patients that they must never share a NovoLog Mix 70/30 FlexPen with another person, even if the needle is changed, because doing so carries a risk for transmission of bloodborne pathogens.

17.2 Physician Instructions

Maintenance of normal or near-normal glucose control is a treatment goal in diabetes mellitus and has been associated with a reduction in diabetic complications. Patients should be informed about potential risks and advantages of NovoLog Mix 70/30 therapy including the possible adverse reactions. Patients should also be offered continued education and advice on insulin therapies, injection technique, life-style management, regular glucose monitoring, periodic glycosylated hemoglobin testing, recognition and management of hypo- and hyperglycemia, adherence to meal planning, complications of insulin therapy, timing of dose, instruction for use of injection devices, and proper storage of insulin. See Patient Information supplied with the product. Patients should be informed that frequent, patient-performed blood glucose measurements are needed to achieve optimal glycemic control and avoid both hyper- and hypoglycemia, and diabetic ketoacidosis.

The patient's ability to concentrate and react may be impaired as a result of hypoglycemia. This may present a risk in situations where these abilities are especially important, such as driving or operating other machinery. Patients who have frequent hypoglycemia or reduced or absent warning signs of hypoglycemia should be advised to use caution when driving or operating machinery.

Accidental substitutions between NovoLog Mix 70/30 and other insulin products have been reported. Patients should be instructed to always carefully check that they are administering the appropriate insulin to avoid medication errors between NovoLog Mix 70/30 and any other insulin. **The prescription for NovoLog Mix 70/30 should be written clearly in order to avoid confusion with other insulin products, for example, NovoLog or Novolin 70/30.** In addition, the written prescription should clearly indicate the presentation, for example FlexPen or vial.

Rx only

Date of Issue: February 2015

Version: XX

Novo Nordisk[®], NovoLog[®], FlexPen[®], and Novolin[®] are registered trademarks of Novo Nordisk[®] A/S.

NovoLog[®] Mix 70/30 is covered by US Patent Nos. 5,547,930, 5,618,913, 5,834,422, 5,840,680, 5,866,538 and other patents pending.

FlexPen[®] is covered by US Patent Nos. 6,004,297, RE 43,834, RE 41,956 and other patents pending.

© 2002 – 2015 Novo Nordisk

Manufactured by:
Novo Nordisk A/S
DK-2880 Bagsvaerd, Denmark

For information about NovoLog Mix 70/30 contact:
Novo Nordisk Inc.
800 Scudders Mill Road
Plainsboro, New Jersey 08536
1-800-727-6500

www.novonordisk-us.com

Patient Information

NovoLog® Mix 70/30 (NŌ-vō-log-MIX-SEV-en-tee-THIR-tee)

(70% insulin aspart protamine suspension and 30% insulin aspart injection, [rDNA origin])

Do not share your NovoLog Mix 70/30 FlexPen with other people, even if the needle has been changed. You may give other people a serious infection, or get a serious infection from them.

What is NovoLog Mix 70/30?

- NovoLog Mix 70/30 is a man-made insulin that is used to control high blood sugar in people with diabetes mellitus.
- It is not known if NovoLog Mix 70/30 is safe and effective in children.

Who should not take NovoLog Mix 70/30?

Do not take NovoLog Mix 70/30 if you:

- are having an episode of low blood sugar (hypoglycemia).
- have an allergy to NovoLog Mix 70/30 or any of the ingredients in NovoLog Mix 70/30.

Before taking NovoLog Mix 70/30, tell your healthcare provider about all your medical conditions including, if you are:

- pregnant, planning to become pregnant, or are breastfeeding.
- taking new prescription or over-the-counter medicines, vitamins, or herbal supplements.

Before you start taking NovoLog Mix 70/30, talk to your healthcare provider about low blood sugar and how to manage it.

How should I take NovoLog Mix 70/30?

- **Read the Instructions for Use** that come with your NovoLog Mix 70/30.
- Take NovoLog Mix 70/30 exactly as your healthcare provider tells you to.
- **NovoLog Mix 70/30 starts acting fast. If you have Type 1 diabetes, inject it up to 15 minutes before you eat a meal. Do not inject NovoLog Mix 70/30 if you are not planning to eat within 15 minutes. If you have Type 2 diabetes, you may inject NovoLog Mix 70/30 up to 15 minutes before or after starting your meal.**
- **Do not mix** NovoLog Mix 70/30 with other insulin products **or** use in an insulin pump.
- Know the type and strength of insulin you take. **Do not** change the type of insulin you take unless your healthcare provider tells you to. The amount of insulin and the best time for you to take your insulin may need to change if you take different types of insulin.
- **Check your blood sugar levels.** Ask your healthcare provider what your blood sugars should be and when you should check your blood sugar levels.
- **Do not reuse or share your needles or syringes with other people.** You may give other people a serious infection or get a serious infection from them.

What should I avoid while taking NovoLog Mix 70/30?

While taking NovoLog Mix 70/30 do not:

- Drive or operate heavy machinery, until you know how NovoLog Mix 70/30 affects you.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol.

What are the possible side effects of NovoLog Mix 70/30?

NovoLog Mix 70/30 may cause serious side effects that can lead to death, including:

Low blood sugar (hypoglycemia). Signs and symptoms that may indicate low blood sugar include:

- | | | |
|---------------------------------|-------------------|--|
| • dizziness or light-headedness | • blurred vision | • anxiety, irritability, or mood changes |
| • sweating | • slurred speech | • hunger |
| • confusion | • shakiness | |
| • headache | • fast heart beat | |

Your insulin dose may need to change because of:

- | | | |
|--|--------------------|------------------|
| • change in level of physical activity or exercise | • increased stress | • change in diet |
| • weight gain or loss | • illness | |

Other common side effects of NovoLog Mix 70/30 may include:

- low potassium in your blood (hypokalemia), reactions at the injection site, itching, rash, serious allergic reactions (whole body reactions), skin thickening or pits at the injection site (lipodystrophy), weight gain, and swelling of your hands and feet.

Get emergency medical help if you have:

- trouble breathing, shortness of breath, fast heartbeat, swelling of your face, tongue, or throat, sweating, extreme drowsiness, dizziness, confusion.

These are not all the possible side effects of NovoLog Mix 70/30. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of NovoLog Mix 70/30.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. You can ask your pharmacist or healthcare provider for information about NovoLog Mix 70/30 that is written for health professionals. Do not use NovoLog Mix 70/30 for a condition for which it was not prescribed. Do not give NovoLog Mix 70/30 to other people, even if they have the same symptoms that you have. It may harm them.

What are the ingredients in NovoLog Mix 70/30?

Active Ingredient: 70% insulin aspart protamine suspension and 30% insulin aspart (rDNA origin).

Inactive Ingredients: glycerol, phenol, metacresol, zinc, disodium hydrogen phosphate dihydrate, sodium chloride, protamine sulfate, water for injection, hydrochloric acid or sodium hydroxide.

Manufactured by: Novo Nordisk A/S; DK-2880 Bagsvaerd, Denmark

For more information, go to www.novonordisk-us.com or call 1-800-727-6500.

Instructions For Use

NovoLog[®] Mix 70/30 FlexPen[®]

Read the following instructions carefully before you start using your NovoLog[®] Mix 70/30 FlexPen[®] and each time you get a refill. There may be new information. You should read the instructions even if you have used NovoLog Mix 70/30 FlexPen before.

Do not share your NovoLog Mix 70/30 FlexPen with other people, even if the needle has been changed. You may give other people a serious infection, or get a serious infection from them.

NovoLog Mix 70/30 FlexPen is a disposable dial-a-dose insulin pen. You can select doses from 1 to 60 units in increments of 1 unit. NovoLog Mix 70/30 FlexPen is designed to be used with NovoFine[®], NovoFine[®] Plus or NovoTwist[®] needles.


NovoLog Mix 70/30 FlexPen should not be used by people who are blind or have severe visual problems without the help of a person who has good eyesight and who is trained to use the NovoLog Mix 70/30 FlexPen the right way.

Getting ready


Make sure you have the following items:

- NovoLog Mix 70/30 FlexPen
- New NovoFine, NovoFine Plus or NovoTwist needle
- Alcohol swab


NovoLog[®] Mix 70/30 FlexPen[®]




NovoFine[®]



NovoFine[®] Plus



NovoTwist[®]




Preparing your NovoLog Mix 70/30 FlexPen

- Wash your hands with soap and water.
- Before you start to prepare your injection, check the label to make sure that you are taking the right type of insulin. This is especially important if you take more than 1 type of insulin. NovoLog Mix 70/30 should look cloudy after mixing.

Before your first injection with a new NovoLog Mix 70/30 FlexPen you must mix the insulin:

- A. Let the insulin reach room temperature before you use it. This makes it easier to mix.

Pull off the pen cap (see diagram A).




- B. Roll the pen between your palms 10 times – it is important that the pen is kept horizontal (see diagram B).



- C. Then gently move the pen up and down ten times between position 1 and 2 as shown, so the glass ball moves from one end of the cartridge to the other (see diagram C).


Repeat rolling and moving the pen until the liquid appears white and cloudy.



For every following injection move the pen up and down between positions 1 and 2 at least ten times until the liquid appears white and cloudy.

After mixing, complete all the following steps of the injection right away. If there is a delay, the insulin will need to be mixed again.

Wipe the rubber stopper with an alcohol swab.


 Before you inject, there must be at least 12 units of insulin left in the cartridge to make sure the remaining insulin is evenly mixed. If there are less than 12 units left, use a new NovoLog Mix 70/30 FlexPen.

Attaching the needle


- D. Remove the protective tab from a disposable needle.

Screw the needle tightly onto your NovoLog Mix 70/30 FlexPen. It is important that the needle is put on straight (see diagram D).


Never place a disposable needle on your NovoLog Mix 70/30 FlexPen until you are ready to take your injection.



E. Pull off the big outer needle cap (see diagram E).



F. Pull off the inner needle cap and dispose of it (see diagram F).



- ⚠ Always use a new needle for each injection to help ensure sterility and prevent blocked needles. Do not reuse or share your needles or syringes with other people. You may give other people a serious infection, or get a serious infection from them.
- ⚠ Be careful not to bend or damage the needle before use.
- ⚠ To reduce the risk of a needle stick, **never put the inner needle cap back on the needle.**

Giving the airshot before each injection

Before each injection small amounts of air may collect in the cartridge during normal use. **To avoid injecting air and to make sure you take the right dose of insulin:**



G. Turn the dose selector to select 2 units (see diagram G).

H. Hold your NovoLog Mix 70/30 FlexPen with the needle pointing up. Tap the cartridge gently with your finger a few times to make any air bubbles collect at the top of the cartridge (see diagram H).



- I. Keep the needle pointing upwards, press the push-button all the way in (see diagram I). The dose selector returns to 0.

A drop of insulin should appear at the needle tip. If not, change the needle and repeat the procedure no more than 6 times.

If you do not see a drop of insulin after 6 times, do not use the NovoLog Mix 70/30 FlexPen and contact Novo Nordisk at 1-800-727-6500.

A small air bubble may remain at the needle tip, but it will not be injected.



Selecting your dose

Check and make sure that the dose selector is set at 0.

- J. Turn the dose selector to the number of units you need to inject. The pointer should line up with your dose.

The dose can be corrected either up or down by turning the dose selector in either direction until the correct dose lines up with the pointer (see diagram J). When turning the dose selector, be careful not to press the push-button as insulin will come out.

You cannot select a dose larger than the number of units left in the cartridge.

You will hear a click for every single unit dialed. Do not set the dose by counting the number of clicks you hear.

- ⚠ Do not use the cartridge scale printed on the cartridge to measure your dose of insulin.



Giving the injection

- Do the injection exactly as shown to you by your healthcare provider. Your healthcare provider should tell you if you need to pinch the skin before injecting. Wipe the skin with an alcohol swab and let the area dry.
- Novolog Mix 70/30 can be injected under the skin (subcutaneously) or your stomach area, buttocks, upper legs (thighs), or upper arms.
- For each injection, change (rotate) your injection site within the area of skin that you use. **Do not** use the same injection site for each injection.

- K. Insert the needle into your skin.

Inject the dose by pressing the push-button all the way in until the 0 lines up with the pointer (see diagram K). Be careful only to push the button when injecting.

Turning the dose selector will not inject insulin.



- L. Keep the needle in the skin for at least 6 seconds, and keep the push-button pressed all the way in until the needle has been pulled out from the skin (see diagram L). This will make sure that the full dose has been

given.

You may see a drop of insulin at the needle tip. This is normal and has no effect on the dose you just received. If blood appears after you take the needle out of your skin, press the injection site lightly with an alcohol swab. **Do not rub the area.**



After the injection

Do not recap the needle. Recapping can lead to a needle stick injury. Remove the needle from the NovoLog Mix 70/30 FlexPen after each injection and dispose of it. This helps to prevent infection, leakage of insulin, and will help to make sure you inject the right dose of insulin. If you do not have a sharps container, carefully slip the needle into the outer needle cap. Safely remove the needle and throw it away as soon as you can.

- Put your used NovoLog Mix 70/30 FlexPen and needles in a FDA-cleared sharps disposal container right away after use. Do not throw away (dispose of) loose needles and Pens in your household trash.
- If you do not have a FDA-cleared sharps disposal container, you may use a household container that is:
 - made of a heavy-duty plastic
 - can be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out
 - upright and stable during use
 - leak-resistant
 - properly labeled to warn of hazardous waste inside the container
- When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container. There may be state or local laws about how you should throw away used needles and syringes. For more information about the safe sharps disposal, and for specific information about sharps disposal in the state that you live in, go to the FDA's website at:
<http://www.fda.gov/safesharpsdisposal>.

Do not dispose of your used sharps disposal container in your household trash unless your community guidelines permit this. Do not recycle your used sharps disposal container.

The NovoLog Mix 70/30 FlexPen prevents the cartridge from being completely emptied. It is designed to deliver 300 units.

- M. Put the pen cap on the NovoLog Mix 70/30 FlexPen and store the NovoLog Mix 70/30 FlexPen without the needle attached (see diagram M). Storing without the needle attached helps prevent leaking, blocking of the needle, and air from entering the Pen.



How should I store NovoLog Mix 70/30 FlexPen?

- Store unused NovoLog Mix 70/30 FlexPen in the refrigerator at 36°F to 46°F (2°C to 8°C).
- Store the FlexPen you are currently using out of the refrigerator below 86°F (30°C) for up to 14 days.
- **Do not** freeze NovoLog Mix 70/30. **Do not** use NovoLog Mix 70/30 if it has been frozen.
- Keep NovoLog Mix 70/30 away from heat or light.
- Unused NovoLog Mix 70/30 FlexPen may be used until the expiration date printed on the label, if kept in the refrigerator.

- The NovoLog Mix 70/30 FlexPen you are using should be thrown away after 14 days, even if it still has insulin left in it.
- Store the NovoLog Mix 70/30 FlexPen without the needle attached.

Maintenance

For the safe and proper use of your NovoLog Mix 70/30 FlexPen be sure to handle it with care.. Avoid dropping your NovoLog Mix 70/30 FlexPen as it may damage it. If you are concerned that your NovoLog Mix 70/30 FlexPen is damaged, use a new one. You can clean the outside of your NovoLog Mix 70/30 FlexPen by wiping it with a damp cloth. Do not soak or wash your NovoLog Mix 70/30 FlexPen as it may damage it. Do not refill your NovoLog Mix 70/30 FlexPen.

- ▲ Remove the needle from the NovoLog Mix 70/30 FlexPen after each injection. This helps to ensure sterility, prevent leakage of insulin, and will help to make sure you inject the right dose of insulin for future injections.
- ▲ Be careful when handling used needles to avoid needle sticks and transfer of infectious diseases.
- ▲ Keep your NovoLog Mix 70/30 FlexPen and needles out of the reach of children.
- ▲ Use NovoLog Mix 70/30 FlexPen as directed to treat your diabetes.
- ▲ **Do not** share your NovoLog Mix 70/30 FlexPen or needles with other people. You may give other people a serious infection, or get a serious infection from them.
- ▲ Always use a new needle for each injection.
- ▲ Novo Nordisk is not responsible for harm due to using this insulin pen with products not recommended by Novo Nordisk.
- ▲ As a precautionary measure, always carry a spare insulin delivery device in case your NovoLog Mix 70/30 FlexPen is lost or damaged.
- ▲ Remember to keep the disposable NovoLog Mix 70/30 FlexPen with you. Do not leave it in a car or other location where it can get too hot or too cold.

This Instructions for Use has been approved by the U.S. Food and Drug Administration

Revised: 04/2015

Instructions for Use

NovoLog® Mix 70/30 (NŌvōLog-MIX-SEV-en-tee-THIR-tee)


(70% insulin aspart protamine suspension and 30% insulin aspart [rDNA origin] injection)

10 mL vial (100 Units/mL, U-100)

Read this Instructions for Use before you start taking NovoLog® Mix 70/30 and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

Supplies you will need to give your NovoLog® Mix 70/30 injection:

- 10 mL NovoLog® Mix 70/30 vial
- insulin syringe and needle
- alcohol swab




Preparing your NovoLog® Mix 70/30 dose:

- Wash your hands with soap and water.
- Before you start to prepare your injection, check the NovoLog® Mix 70/30 label to make sure that you are taking the right type of insulin. This is especially important if you use more than 1 type of insulin.
- NovoLog® Mix 70/30 should look white and cloudy after mixing. **Do not** use NovoLog Mix 70/30 if it looks clear or contains any lumps or particles.
- NovoLog® Mix 70/30 is easier to mix when it is at room temperature.
- After mixing NovoLog® Mix 70/30, inject your dose right away. If you wait to inject your dose, the insulin will need to be mixed again.
- **Do not** use NovoLog® Mix 70/30 past the expiration date printed on the label.




Step 1: If you are using a new vial, pull off the tamper-resistant cap (See Figure A).

Step 2: Wipe the rubber stopper with an alcohol swab (See Figure B).




(Figure A Figure B)

Step 3: Roll the NovoLog Mix 70/30 vial between your hands 10 times. Keep the vial in a horizontal (flat) position (See Figure C). Roll the vial between your hands until the Novolog[®] Mix 70/30 looks white and cloudy. Do not shake the vial.









(Figure C)

Step 4: Hold the syringe with the needle pointing up. Pull down on the plunger until the black tip reaches the line for the number of units for your prescribed dose (See Figure D).





(Figure D)

<p>Step 5: Push the needle through the rubber stopper of the NovoLog® Mix 70/30 vial (See Figure E).</p>	 <p>(Figure E)</p>
<p>Step 6: Push the plunger all the way in. This puts air into the NovoLog® Mix 70/30 vial (See Figure F).</p>	 <p>(Figure F)</p>
<p>Step 7: Turn the NovoLog® Mix 70/30 vial and syringe upside down and slowly pull the plunger down until the black tip is a few units past the line for your dose (See Figure G).</p> <ul style="list-style-type: none"> • If there are air bubbles, tap the syringe gently a few times to let any air bubbles rise to the top (See Figure H). 	 <p>(Figure G)</p>  <p>(Figure H)</p>
<p>Step 8: Slowly push the plunger up until the black tip reaches the line for your NovoLog® Mix 70/30 dose (See Figure I).</p>	 <p>(Figure I)</p>

Step 9: Check the syringe to make sure you have the right dose of NovoLog® Mix 70/30.	
Step 10: Pull the syringe out of the vial's rubber stopper (See Figure J).	
	(Figure J)


Giving your injection:

- Inject your NovoLog® Mix 70/30 exactly as your healthcare provider has shown you. Your healthcare provider should tell you if you need to pinch the skin before injecting.
- NovoLog® Mix 70/30 is injected under the skin (subcutaneously) of your stomach area, buttocks, upper legs, or upper arms.
- Change (rotate) your injection sites within the area you choose for each dose.
Do not use the same injection site for each injection.

Step 11: Choose your injection site and wipe the skin with an alcohol swab. Let the injection site dry before you inject your dose (See Figure K).	
	(Figure K)
Step 12: Insert the needle into your skin. Push down on the plunger to inject your dose (See Figure L). Needle should remain in the skin for at least 6 seconds to make sure you have injected all the insulin.	
	(Figure L)

Step 13: Pull the needle out of your skin. After that, you may see a drop of NovoLog[®] Mix 70/30 at the needle tip. This is normal and does not affect the dose you just received (See Figure M).

- If you see blood after you take the needle out of your skin, press the injection site lightly with a piece of gauze or an alcohol swab. **Do not** rub the area.



(Figure M)

After your injection:

- **Do not** recap the needle. Recapping the needle can lead to a needle stick injury.
- Throw away empty insulin vials, used syringes and needles in a sharps container or some type of hard plastic or metal container with a screw on cap such as a detergent bottle or coffee can. Check with your healthcare provider about the right way to throw away the container. There may be local or state laws about how to throw away used syringes and needles. **Do not** throw away used syringes and needles in household trash or recycling bins.

How should I store NovoLog[®] Mix 70/30?

- **Do not** freeze NovoLog[®] Mix 70/30. **Do not** use NovoLog[®] Mix 70/30 if it has been frozen.
- Keep NovoLog[®] Mix 70/30 away from heat or light.
- Store opened and unopened NovoLog[®] Mix 70/30 vials in the refrigerator at 36°F to 46°F (2°C to 8°C). Opened NovoLog[®] Mix 70/30 vials can also be stored out of the refrigerator below 86°F (30°C).
- Unopened vials may be used until the expiration date printed on the label, if they are kept in the refrigerator.
- Opened NovoLog[®] Mix 70/30 vials should be thrown away after 28 days, even if they still have insulin left in them.

General information about the safe and effective use of NovoLog[®] Mix 70/30

- Always use a new syringe and needle for each injection.
- Do not share syringes or needles.
- Keep NovoLog[®] Mix 70/30 vials, syringes, and needles out of the reach of children.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Manufactured by:
Novo Nordisk A/S
DK-2880 Bagsvaerd, Denmark

Revised: December 2012

NovoLog[®] is a registered trademark of Novo Nordisk A/S.

NovoLog[®] Mix 70/30 is covered by US Patent Nos. 5,547,930, 5,618,913, 5,834,422, 5,840,680, 5,866,538 and other patents pending.

© 2002-2012 Novo Nordisk A/S

For information about NovoLog[®] Mix 70/30 contact:
Novo Nordisk Inc.
100 College Road West
Princeton, New Jersey 08540
1-800-727-6500
www.novonordisk-us.com