

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Adult Patients

PREZISTA[®], co-administered with ritonavir (PREZISTA/ritonavir), and with other antiretroviral agents, is indicated for the treatment of human immunodeficiency virus (HIV-1) infection.

This indication is based on analyses of plasma HIV-1 RNA levels and CD4+ cell counts from 2 controlled Phase 3 trials of 48 weeks duration in antiretroviral treatment-naïve and treatment-experienced patients and 2 controlled Phase 2 trials of 96 weeks duration in clinically advanced, treatment-experienced adult patients.

1.2 Pediatric Patients

PREZISTA, co-administered with ritonavir (PREZISTA/ritonavir), and with other antiretroviral agents, is indicated for the treatment of HIV-1 infection in pediatric patients 3 years of age and older [see *Use in Specific Populations (8.4)*].

The indication for treatment-experienced pediatric patients 3 to less than 18 years of age is based on analyses of plasma HIV-1 RNA levels and CD4+ cell counts from two open-label Phase 2 trials in antiretroviral treatment-experienced pediatric subjects (24-week analysis for one trial in patients 6 to less than 18 years of age; 48-week analysis for one trial in patients 3 to less than 6 years of age). The indication for treatment-naïve pediatric patients or antiretroviral treatment-experienced patients with no darunavir resistance associated substitutions is based on one open-label Phase 2 trial of 48 weeks duration in antiretroviral treatment-naïve subjects 12 to less than 18 years of age and pharmacokinetic modeling and simulation for patients 3 to less than 12 years of age.

In treatment-experienced adult and pediatric patients, the following points should be considered when initiating therapy with PREZISTA/ritonavir:

- Treatment history and, when available, genotypic or phenotypic testing should guide the use of PREZISTA/ritonavir [see *Microbiology (12.4)*].
- The use of other active agents with PREZISTA/ritonavir is associated with a greater likelihood of treatment response [see *Microbiology (12.4)* and *Clinical Studies (14.3)*].

2 DOSAGE AND ADMINISTRATION

2.1 Adult Patients

PREZISTA must be co-administered with ritonavir to exert its therapeutic effect. Failure to correctly co-administer PREZISTA with ritonavir will result in plasma levels of darunavir that will be insufficient to achieve the desired antiviral effect and will alter some drug interactions.

Patients who have difficulty swallowing PREZISTA tablets can use the 100 mg/mL PREZISTA oral suspension.

Treatment-Naïve Adult Patients

The recommended oral dose of PREZISTA is 800 mg (one 800 mg tablet or 8 mL of the oral suspension) taken with ritonavir 100 mg (one 100 mg tablet/capsule or 1.25 mL of a 80 mg/mL ritonavir oral solution) once daily and with food. An 8 mL darunavir dose should be taken as two 4 mL administrations with the included oral dosing syringe.

Treatment-Experienced Adult Patients

Table 1: Treatment-Experienced Adult Patients □	
With no darunavir resistance associated substitutions*	With at least one darunavir resistance associated substitution*
Formulations: PREZISTA tablets or oral suspension (100 mg/mL) and ritonavir tablets/capsules (100 mg) or solution (80 mg/mL)	
PREZISTA 800 mg (one 800 mg tablet) or 8 mL [†] (oral suspension) once daily with ritonavir 100 mg (one 100 mg tablet/capsule or 1.25 mL (oral solution) once daily and with food	PREZISTA 600 mg (e.g. one 600 mg tablet) or 6 mL (oral suspension) twice daily with ritonavir 100 mg (one 100 mg tablet/capsule or 1.25 mL (oral solution) twice daily and with food

* V11I, V32I, L33F, I47V, I50V, I54L, I54M, T74P, L76V, I84V and L89V

[†] An 8 mL darunavir dose should be taken as two 4 mL administrations with the included oral dosing syringe

For antiretroviral treatment-experienced patients genotypic testing is recommended. However, when genotypic testing is not feasible, PREZISTA/ritonavir 600/100 mg twice daily dosing is recommended.

2.2 Pediatric Patients (age 3 to less than 18 years)

Healthcare professionals should pay special attention to accurate dose selection of PREZISTA, transcription of the medication order, dispensing information and dosing instruction to minimize risk for medication errors, overdose, and underdose.

Prescribers should select the appropriate dose of PREZISTA/ritonavir for each individual child based on body weight (kg) and should not exceed the recommended dose for adults.

Before prescribing PREZISTA, children weighing greater than or equal to 15 kg should be assessed for the ability to swallow tablets. If a child is unable to reliably swallow a tablet, the use of PREZISTA oral suspension should be considered.

The recommended dose of PREZISTA/ritonavir for pediatric patients (3 to less than 18 years of age and weighing at least 10 kg is based on body weight (see Tables 2, 3, 4, and 5) and should not exceed the recommended adult dose. PREZISTA should be taken with ritonavir and with food.

The recommendations for the PREZISTA/ritonavir dosage regimens were based on the following:

Twice daily dosing

- Results from two trials in treatment-experienced pediatric subjects 3 to less than 18 years of age demonstrating similar darunavir plasma exposures, virologic response rate and safety profile compared to treatment-experienced adults.

Once daily dosing

- Results from one trial in treatment-naïve pediatric subjects 12 to less than 18 years of age demonstrating similar darunavir plasma exposures, virologic response rate and safety profile compared to treatment-naïve adults.
- Results from population pharmacokinetic modeling and simulation in children 3 to less than 12 years of age predicting similar darunavir plasma exposures compared to treatment-naïve adults. Although no clinical trial was conducted to collect exposure-safety data, the predicted exposures from the once daily dosing is supported by exposures observed in a pediatric clinical trial where twice-daily dosing was administered.

Dosing recommendations for treatment-naïve pediatric patients or antiretroviral treatment-experienced pediatric patients with no darunavir resistance associated substitutions

- *Pediatric patients weighing at least 10 kg but less than 15 kg*

The weight-based dose in antiretroviral treatment-naïve pediatric patients or antiretroviral treatment-experienced pediatric patients with no darunavir resistance associated substitutions is PREZISTA 35 mg/kg once daily with ritonavir 7 mg/kg once daily using the following table.

Table 2: Recommended dose for pediatric patients weighing 10 kg to less than 15 kg who are treatment-naïve or treatment-experienced with no darunavir resistance associated substitutions*	
Body weight (kg)	Formulation: PREZISTA oral suspension (100 mg/mL) and Ritonavir oral solution (80 mg/mL)
Dose: once daily with food	
Greater than or equal to 10 kg to less than 11 kg	PREZISTA 3.6 mL [‡] (350 mg) with ritonavir 0.8 mL (64 mg)
Greater than or equal to 11 kg to less than 12 kg	PREZISTA 4 mL [‡] (385 mg) with ritonavir 0.8 mL (64 mg)
Greater than or equal to 12 kg to less than 13 kg	PREZISTA 4.2 mL (420 mg) with ritonavir 1 mL (80 mg)
Greater than or equal to 13 kg to less than 14 kg	PREZISTA 4.6 mL [‡] (455 mg) with ritonavir 1 mL (80 mg)
Greater than or equal to 14 kg to less than 15 kg	PREZISTA 5 mL [‡] (490 mg) with ritonavir 1.2 mL (96 mg)

* darunavir resistance associated substitutions: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

‡ The 350 mg, 385 mg, 455 mg and 490 mg darunavir dose for the specified weight groups were rounded up for suspension dosing convenience to 3.6 mL, 4 mL, 4.6 mL and 5 mL, respectively.

- *Pediatric patients weighing at least 15 kg*

Pediatric patients weighing at least 15 kg can be dosed with PREZISTA oral tablet(s) or suspension using the following table:

Table 3: Recommended dose for pediatric patients weighing at least 15 kg who are treatment-naïve or treatment-experienced with no darunavir resistance associated substitutions*		
Body Weight (kg)	Formulation: PREZISTA tablet(s) and ritonavir capsules or tablets (100 mg)	Formulation: PREZISTA oral suspension (100 mg/mL) and ritonavir oral solution (80 mg/mL)
	Dose: once daily with food	Dose: once daily with food
Greater than or equal to 15 kg to less than 30 kg	PREZISTA 600 mg with ritonavir 100 mg	PREZISTA 6 mL (600 mg) with ritonavir 1.25 mL (100 mg)
Greater than or equal to 30 kg to less than 40 kg	PREZISTA 675 mg with ritonavir 100 mg	PREZISTA 6.8 mL [§] (675 mg) with ritonavir 1.25 mL (100 mg)
Greater than or equal to 40 kg	PREZISTA 800 mg with ritonavir 100 mg	PREZISTA 8 mL [‡] (800 mg) with ritonavir 1.25 mL (100 mg)

* darunavir resistance associated substitutions: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

§ The 675 mg dose using darunavir tablets for this weight group is rounded up to 6.8 mL for suspension dosing convenience.

‡ The 6.8 mL and 8 mL darunavir dose should be taken as two (3.4 mL or 4 mL respectively) administrations with the included oral dosing syringe

Dosing recommendations for treatment-experienced pediatric patients with at least one darunavir resistance associated substitutions

- *Pediatric patients weighing at least 10 kg but less than 15 kg*

The weight-based dose in antiretroviral treatment-experienced pediatric patients with at least one darunavir resistance associated substitution is PREZISTA 20 mg/kg twice daily with ritonavir 3 mg/kg twice daily using the following table:

Table 4: Recommended dose for pediatric patients weighing 10 kg to less than 15 kg who are treatment-experienced with at least one darunavir resistance associated substitution*	
Body weight (kg)	Formulation: PREZISTA oral suspension (100 mg/mL) and Ritonavir oral solution (80 mg/mL)
	Dose: twice daily with food
Greater than or equal to 10 kg to less than 11 kg	PREZISTA 2 mL (200 mg) with ritonavir 0.4 mL (32 mg)
Greater than or equal to 11 kg to less than 12 kg	PREZISTA 2.2 mL (220 mg) with ritonavir 0.4 mL (32 mg)
Greater than or equal to 12 kg to less than 13 kg	PREZISTA 2.4 mL (240 mg) with ritonavir 0.5 mL (40 mg)
Greater than or equal to 13 kg to less than 14 kg	PREZISTA 2.6 mL (260 mg) with ritonavir 0.5 mL (40 mg)
Greater than or equal to 14 kg to less than 15 kg	PREZISTA 2.8 mL (280 mg) with ritonavir 0.6 mL (48 mg)

* darunavir resistance associated substitutions: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

- *Pediatric patients weighing at least 15 kg*

Pediatric patients weighing at least 15 kg can be dosed with PREZISTA oral tablet(s) or suspension using the following table:

Table 5: Recommended dose for pediatric patients weighing at least 15 kg who are treatment-experienced with at least one darunavir resistance associated substitution*		
Body Weight (kg)	Formulation: PREZISTA tablet(s) and ritonavir tablets, capsules (100 mg) or oral solution (80 mg/mL)	Formulation: PREZISTA oral suspension (100 mg/mL) and ritonavir oral solution (80 mg/mL)
	Dose: twice daily with food	Dose: twice daily with food
Greater than or equal to 15 kg to less than 30 kg	PREZISTA 375 mg with ritonavir 0.6 mL (48 mg)	PREZISTA 3.8 mL (375 mg) with ritonavir 0.6 mL (48 mg)
Greater than or equal to 30 kg to less than 40 kg	PREZISTA 450 mg with ritonavir 0.75 mL (60 mg)	PREZISTA 4.6 mL (450 mg) [§] with ritonavir 0.75 mL (60 mg)
Greater than or equal to 40 kg	PREZISTA 600 mg with ritonavir 100 mg	PREZISTA 6 mL (600 mg) with ritonavir 1.25 mL (100 mg)

* darunavir resistance associated substitutions: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

§ The 375 mg and 450 mg dose using darunavir tablets for this weight group is rounded up to 3.8 mL and 4.6 mL for suspension dosing convenience.

Do not use PREZISTA/ritonavir in pediatric patients below 3 years of age [see *Warnings and Precautions (5.11)* and *Nonclinical Toxicology (13.2)*].

2.3 Patients with Hepatic Impairment

No dose adjustment is required in patients with mild or moderate hepatic impairment. No data are available regarding the use of PREZISTA/ritonavir when co-administered to subjects with severe hepatic impairment; therefore, PREZISTA/ritonavir is not recommended for use in patients with severe hepatic impairment [see *Use in Specific Populations (8.6)* and *Clinical Pharmacology (12.3)*].

3 DOSAGE FORMS AND STRENGTHS

3.1 PREZISTA 100 mg/mL Oral Suspension

PREZISTA (darunavir) 100 mg/mL oral suspension is supplied as a white to off-white opaque suspension for oral use, containing darunavir ethanolate equivalent to 100 mg of darunavir per mL of suspension.

3.2 PREZISTA 75 mg Tablets

PREZISTA (darunavir) 75 mg tablets are supplied as white, caplet-shaped, film-coated tablets containing darunavir ethanolate equivalent to 75 mg of darunavir per tablet. Each tablet is debossed with “75” on one side and “TMC” on the other side.

3.3 PREZISTA 150 mg Tablets

PREZISTA (darunavir) 150 mg tablets are supplied as white, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 150 mg of darunavir per tablet. Each tablet is debossed with “150” on one side and “TMC” on the other side.

3.4 PREZISTA 600 mg Tablets

PREZISTA (darunavir) 600 mg tablets are supplied as orange, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 600 mg of darunavir per tablet. Each tablet is debossed with “600MG” on one side and “TMC” on the other side.

3.5 PREZISTA 800 mg Tablets

PREZISTA (darunavir) 800 mg tablets are supplied as dark red, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 800 mg of darunavir per tablet. Each tablet is debossed with “800” on one side and “T” on the other side.

4 CONTRAINDICATIONS

Co-administration of PREZISTA/ritonavir is contraindicated with drugs that are highly dependent on CYP3A for clearance and for which elevated plasma concentrations are associated with serious and/or life-threatening events (narrow therapeutic index). These drugs and other contraindicated drugs (which may lead to reduced efficacy of darunavir) are listed in Table 6 [also see *Drug Interactions* (7.3), Table 11].

Drug Class	Drugs Within Class That Are Contraindicated With PREZISTA/ritonavir	Clinical Comment
Alpha 1-adrenoreceptor antagonist	Alfuzosin	Potential for serious and/or life-threatening reactions such as hypotension.
Antiarrhythmic	Dronedarone	Potential for serious and/or life-threatening reactions such as cardiac arrhythmias.
Anti-gout	Colchicine	Potential for serious and/or life-threatening reactions in patients with renal and/or hepatic impairment.
Anti-anginal	Ranolazine	Potential for serious and/or life-threatening reactions.
Antipsychotic/Neuroleptic	Pimozide	Potential for serious and/or life-threatening reactions such as cardiac arrhythmias.
Ergot Derivative	Dihydroergotamine, Ergotamine, Methylergonovine	Potential for serious and/or life-threatening events such as acute ergot toxicity characterized by peripheral vasospasm and ischemia of the extremities and other tissues.
GI Motility Agent	Cisapride	Potential for serious and/or life-threatening reactions such as cardiac arrhythmias.
Sedative/hypnotic	Orally administered Midazolam, Triazolam	Triazolam and orally administered midazolam are extensively metabolized by CYP3A. Co-administration of triazolam or orally administered midazolam with PREZISTA/ritonavir may cause large increases in the concentrations of these benzodiazepines. Potential for serious and/or life-threatening events such as prolonged or increased sedation or respiratory depression.
Herbal Product	St. John’s Wort (<i>Hypericum</i>)	Patients taking PREZISTA/ritonavir should

	<i>perforatum</i>)	not use products containing St. John's wort because co-administration may result in reduced plasma concentrations of darunavir. This may result in loss of therapeutic effect and development of resistance.
HMG-CoA Reductase Inhibitor	Lovastatin, Simvastatin	Potential for serious reactions such as myopathy including rhabdomyolysis. For dosing recommendation regarding atorvastatin, pravastatin and rosuvastatin, see Table 11: Established and Other Potentially Significant Drug Interactions: Alterations in Dose or Regimen May Be Recommended Based on Drug Interaction Studies or Predicted Interaction.
Antimycobacterial	Rifampin	Rifampin is a potent inducer of CYP450 metabolism. PREZISTA/ritonavir should not be used in combination with rifampin, as this may cause significant decreases in darunavir plasma concentrations. This may result in loss of therapeutic effect to PREZISTA.
PDE-5 inhibitor	Sildenafil for treatment of pulmonary arterial hypertension	A safe and effective dose for the treatment of pulmonary arterial hypertension has not been established with PREZISTA/ritonavir. There is an increased potential for sildenafil-associated adverse events (which include visual disturbances, hypotension, prolonged erection, and syncope).

Due to the need for co-administration of PREZISTA with ritonavir, please refer to ritonavir prescribing information for a description of ritonavir contraindications.

5 WARNINGS AND PRECAUTIONS

5.1 General

PREZISTA must be co-administered with ritonavir and food to achieve the desired antiviral effect. Failure to administer PREZISTA with ritonavir and food may result in a loss of efficacy of darunavir.

Please refer to ritonavir prescribing information for additional information on precautionary measures.

5.2 Hepatotoxicity

Drug-induced hepatitis (e.g., acute hepatitis, cytolytic hepatitis) has been reported with PREZISTA/ritonavir. During the clinical development program (N=3063), hepatitis was reported in 0.5% of patients receiving combination therapy with PREZISTA/ritonavir. Patients with pre-existing liver dysfunction, including chronic active hepatitis B or C, have an increased risk for liver function abnormalities including severe hepatic adverse events.

Post-marketing cases of liver injury, including some fatalities, have been reported. These have generally occurred in patients with advanced HIV-1 disease taking multiple concomitant medications, having co-morbidities including hepatitis B or C co-infection, and/or developing immune reconstitution syndrome. A causal relationship with PREZISTA/ritonavir therapy has not been established.

Appropriate laboratory testing should be conducted prior to initiating therapy with PREZISTA/ritonavir and patients should be monitored during treatment. Increased AST/ALT monitoring should be considered in patients with underlying chronic hepatitis, cirrhosis, or in patients who have pre-treatment elevations of transaminases, especially during the first several months of PREZISTA/ritonavir treatment.

Evidence of new or worsening liver dysfunction (including clinically significant elevation of liver enzymes and/or symptoms such as fatigue, anorexia, nausea, jaundice, dark urine, liver tenderness, hepatomegaly) in patients on PREZISTA/ritonavir should prompt consideration of interruption or discontinuation of treatment.

5.3 Severe Skin Reactions

During the clinical development program (n=3063), severe skin reactions, accompanied by fever and/or elevations of transaminases in some cases, have been reported in 0.4% of subjects. Stevens-Johnson Syndrome was rarely (less than 0.1%) reported during the clinical development program. During post-marketing experience toxic epidermal necrolysis, drug rash with eosinophilia and systemic symptoms, and acute generalized exanthematous pustulosis have been reported. Discontinue PREZISTA/ritonavir immediately if signs or symptoms of severe skin reactions develop. These can include but are not limited to severe rash or rash accompanied with fever, general malaise, fatigue, muscle or joint aches, blisters, oral lesions, conjunctivitis, hepatitis and/or eosinophilia.

Rash (all grades, regardless of causality) occurred in 10.3% of subjects treated with PREZISTA/ritonavir [also see *Adverse Reactions (6)*]. Rash was mostly mild-to-moderate, often occurring within the first four weeks of treatment and resolving with continued dosing. The discontinuation rate due to rash in subjects using PREZISTA/ritonavir was 0.5%.

Rash occurred more commonly in treatment-experienced subjects receiving regimens containing PREZISTA/ritonavir + raltegravir compared to subjects receiving PREZISTA/ritonavir without raltegravir or raltegravir without PREZISTA/ritonavir. However, rash that was considered drug related occurred at similar rates for all three groups. These rashes were mild to moderate in severity and did not limit therapy; there were no discontinuations due to rash.

5.4 Sulfa Allergy

Darunavir contains a sulfonamide moiety. PREZISTA should be used with caution in patients with a known sulfonamide allergy. In clinical studies with PREZISTA/ritonavir, the incidence and severity of rash were similar in subjects with or without a history of sulfonamide allergy.

5.5 Risk of Serious Adverse Reactions due to Drug Interactions

Initiation of PREZISTA/ritonavir, a CYP3A inhibitor, in patients receiving medications metabolized by CYP3A or initiation of medications metabolized by CYP3A in patients already receiving PREZISTA/ritonavir, may increase plasma concentrations of medications metabolized by CYP3A. Initiation of medications that inhibit or induce CYP3A may increase or decrease concentrations of PREZISTA/ritonavir, respectively. These interactions may lead to:

- Clinically significant adverse reactions, potentially leading to severe, life threatening, or fatal events from greater exposures of concomitant medications.
- Clinically significant adverse reactions from greater exposures of PREZISTA/ritonavir.
- Loss of therapeutic effect of PREZISTA/ritonavir and possible development of resistance.

See Table 11 for steps to prevent or manage these possible and known significant drug interactions, including dosing recommendations [see *Drug Interactions (7)*]. Consider the potential for drug interactions prior to and during PREZISTA/ritonavir therapy; review concomitant medications during PREZISTA/ritonavir therapy; and monitor for the adverse reactions associated with the concomitant drugs [see *Contraindications (4)* and *Drug Interactions (7)*].

5.6 Diabetes Mellitus / Hyperglycemia

New onset diabetes mellitus, exacerbation of pre-existing diabetes mellitus, and hyperglycemia have been reported during postmarketing surveillance in HIV-infected patients receiving protease inhibitor (PI) therapy. Some patients required either initiation or dose adjustments of insulin or oral hypoglycemic agents for treatment of these events. In some cases, diabetic ketoacidosis has occurred. In those patients who discontinued PI therapy, hyperglycemia persisted in some cases. Because these events have been reported voluntarily during clinical practice, estimates of frequency cannot be made and causal relationships between PI therapy and these events have not been established.

5.7 Fat Redistribution

Redistribution/accumulation of body fat, including central obesity, dorsocervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and

“cushingoid appearance” have been observed in patients receiving antiretroviral therapy. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.

5.8 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in patients treated with combination antiretroviral therapy, including PREZISTA. During the initial phase of combination antiretroviral treatment, patients whose immune systems respond may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* infection, cytomegalovirus, *Pneumocystis jirovecii* pneumonia [PCP], or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves’ disease, polymyositis, and Guillain-Barré syndrome) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable, and can occur many months after initiation of antiretroviral treatment.

5.9 Hemophilia

There have been reports of increased bleeding, including spontaneous skin hematomas and hemarthrosis in patients with hemophilia type A and B treated with PIs. In some patients, additional factor VIII was given. In more than half of the reported cases, treatment with PIs was continued or reintroduced if treatment had been discontinued. A causal relationship between PI therapy and these episodes has not been established.

5.10 Resistance/Cross-Resistance

Because the potential for HIV cross-resistance among PIs has not been fully explored in PREZISTA/ritonavir treated patients, the effect therapy with PREZISTA will have on the activity of subsequently administered PIs is unknown [see *Microbiology (12.4)*].

5.11 Pediatric Patients

Do not administer PREZISTA/ritonavir in pediatric patients below 3 years of age in view of toxicity and mortality observed in juvenile rats dosed with darunavir (from 20 mg/kg to 1000 mg/kg) up to days 23 to 26 of age [see *Use in Specific Populations (8.1 and 8.4)*, *Clinical Pharmacology (12.3)*, and *Nonclinical Toxicology (13.2)*].

6 ADVERSE REACTIONS

The overall safety profile of PREZISTA/ritonavir 800/100 mg once daily and PREZISTA/ritonavir 600/100 mg twice daily is based on clinical trials and post-marketing data, and is consistent with the data presented below.

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Due to the need for co-administration of PREZISTA with ritonavir, please refer to ritonavir prescribing information for ritonavir-associated adverse reactions.

6.1 Clinical Trials Experience: Treatment-Naïve Adults

Study TMC114-C211

The safety assessment is based on all safety data from the Phase 3 trial TMC114-C211 comparing PREZISTA/ritonavir 800/100 mg once daily versus lopinavir/ritonavir 800/200 mg per day in 689 antiretroviral treatment-naïve HIV-1-infected adult subjects. The total mean exposure for subjects in the PREZISTA/ritonavir 800/100 mg once daily arm and in the lopinavir/ritonavir 800/200 mg per day arm was 162.5 and 153.5 weeks, respectively.

The majority of the adverse drug reactions (ADRs) reported during treatment with PREZISTA/ritonavir 800/100 mg once daily were mild in severity. The most common clinical ADRs to PREZISTA/ritonavir 800/100 mg once daily (greater than or equal to 5%) of at least moderate intensity (greater than or equal to Grade 2) were diarrhea, headache, abdominal pain and rash. 2.3% of subjects in the PREZISTA/ritonavir arm discontinued treatment due to ADRs.

ADRs to PREZISTA/ritonavir 800/100 mg once daily of at least moderate intensity (greater than or equal to Grade 2) in antiretroviral treatment-naïve HIV-1-infected adult subjects are presented in Table 7 and subsequent text below the table.

Table 7: Selected Clinical Adverse Drug Reactions to PREZISTA/ritonavir 800/100 mg Once Daily* of at Least Moderate Intensity (≥ Grade 2) Occurring in ≥ 2% of Antiretroviral Treatment-Naïve HIV-1-Infected Adult Subjects		
System Organ Class, Preferred Term, %	Randomized Study TMC114-C211	
	PREZISTA/ritonavir 800/100 mg once daily + TDF/FTC N=343	lopinavir/ritonavir 800/200 mg per day + TDF/FTC N=346
Gastrointestinal Disorders		
Abdominal pain	6%	6%
Diarrhea	9%	16%
Nausea	4%	4%
Vomiting	2%	4%
General Disorders and Administration Site Conditions		
Fatigue	<1%	3%
Metabolism and Nutrition Disorders		
Anorexia	2%	<1%
Nervous System Disorders		
Headache	7%	6%

Skin and Subcutaneous Tissue Disorders		
Rash	6%	7%

N=total number of subjects per treatment group

TDF=tenofovir disoproxil fumarate

FTC=emtricitabine

* Excluding laboratory abnormalities reported as ADRs

Less Common Adverse Reactions

Treatment-emergent ADRs of at least moderate intensity (greater than or equal to Grade 2) occurring in less than 2% of antiretroviral treatment-naïve subjects receiving PREZISTA/ritonavir 800/100 mg once daily are listed below by body system:

Gastrointestinal Disorders: acute pancreatitis, dyspepsia, flatulence

General Disorders and Administration Site Conditions: asthenia

Hepatobiliary Disorders: acute hepatitis (e.g., acute hepatitis, cytolytic hepatitis, hepatotoxicity)

Immune System Disorders: (drug) hypersensitivity, immune reconstitution syndrome

Metabolism and Nutrition Disorders: diabetes mellitus

Musculoskeletal and Connective Tissue Disorders: myalgia, osteonecrosis

Psychiatric Disorders: abnormal dreams

Skin and Subcutaneous Tissue Disorders: angioedema, pruritus, Stevens-Johnson Syndrome, urticaria

Laboratory abnormalities:

Selected Grade 2 to 4 laboratory abnormalities that represent a worsening from baseline observed in antiretroviral treatment-naïve adult subjects treated with PREZISTA/ritonavir 800/100 mg once daily are presented in Table 8.

Table 8: Grade 2 to 4 Laboratory Abnormalities Observed in Antiretroviral Treatment-Naïve HIV-1-Infected Adult Subjects*			
Laboratory Parameter Preferred Term, %	Limit	Randomized Study TMC114-C211	
		PREZISTA/ritonavir 800/100 mg once daily + TDF/FTC	lopinavir/ritonavir 800/200 mg per day + TDF/FTC
Biochemistry			
Alanine Aminotransferase			
Grade 2	> 2.5 to ≤ 5.0 X ULN	9%	9%
Grade 3	> 5.0 to ≤ 10.0 X ULN	3%	3%
Grade 4	> 10.0 X ULN	<1%	3%
Aspartate Aminotransferase			
Grade 2	> 2.5 to ≤ 5.0 X ULN	7%	10%
Grade 3	> 5.0 to ≤ 10.0 X ULN	4%	2%
Grade 4	> 10.0 X ULN	1%	3%
Alkaline Phosphatase			
Grade 2	> 2.5 to ≤ 5.0 X ULN	1%	1%
Grade 3	> 5.0 to ≤ 10.0 X ULN	0%	<1%

Grade 4	> 10.0 X ULN	0%	0%
Hyperbilirubinemia			
Grade 2	> 1.5 to ≤ 2.5 X ULN	<1%	5%
Grade 3	> 2.5 to ≤ 5.0 X ULN	<1%	<1%
Grade 4	> 5.0 X ULN	0%	0%
Triglycerides			
Grade 2	5.65-8.48 mmol/L 500-750 mg/dL	3%	10%
Grade 3	8.49-13.56 mmol/L 751-1200 mg/dL	2%	5%
Grade 4	> 13.56 mmol/L > 1200 mg/dL	1%	1%
Total Cholesterol			
Grade 2	6.20-7.77 mmol/L 240-300 mg/dL	23%	27%
Grade 3	> 7.77 mmol/L > 300 mg/dL	1%	5%
Low-Density Lipoprotein Cholesterol			
Grade 2	4.13-4.90 mmol/L 160-190 mg/dL	14%	12%
Grade 3	≥ 4.91 mmol/L ≥ 191 mg/dL	9%	6%
Elevated Glucose Levels			
Grade 2	6.95-13.88 mmol/L 126-250 mg/dL	11%	10%
Grade 3	13.89-27.75 mmol/L 251-500 mg/dL	1%	<1%
Grade 4	> 27.75 mmol/L > 500 mg/dL	0%	0%
Pancreatic Lipase			
Grade 2	> 1.5 to ≤ 3.0 X ULN	3%	2%
Grade 3	> 3.0 to ≤ 5.0 X ULN	<1%	1%
Grade 4	> 5.0 X ULN	0%	<1%
Pancreatic Amylase			
Grade 2	> 1.5 to ≤ 2.0 X ULN	5%	2%
Grade 3	> 2.0 to ≤ 5.0 X ULN	5%	4%
Grade 4	> 5.0 X ULN	0%	<1%

N=total number of subjects per treatment group

TDF=tenofovir disoproxil fumarate

FTC=emtricitabine

* Grade 4 data not applicable in Division of AIDS grading scale.

6.2 Clinical Trials Experience: Treatment-Experienced Adults

Study TMC114-C214

The safety assessment is based on all safety data from the Phase 3 trial TMC114-C214 comparing PREZISTA/ritonavir 600/100 mg twice daily versus lopinavir/ritonavir 400/100 mg twice daily in 595 antiretroviral treatment-experienced HIV-1-infected adult subjects. The total mean exposure for subjects in the PREZISTA/ritonavir 600/100 mg twice daily arm and in the lopinavir/ritonavir 400/100 mg twice daily arm was 80.7 and 76.4 weeks, respectively.

The majority of the ADRs reported during treatment with PREZISTA/ritonavir 600/100 mg twice daily were mild in severity. The most common clinical ADRs to PREZISTA/ritonavir 600/100 mg twice daily (greater than or equal to 5%) of at least moderate intensity (greater than or equal to Grade 2) were diarrhea, nausea, rash, abdominal pain and vomiting. 4.7% of subjects in the PREZISTA/ritonavir arm discontinued treatment due to ADRs.

ADRs to PREZISTA/ritonavir 600/100 mg twice daily of at least moderate intensity (greater than or equal to Grade 2) in antiretroviral treatment-experienced HIV-1-infected adult subjects are presented in Table 9 and subsequent text below the table.

Table 9: Selected Clinical Adverse Drug Reactions to PREZISTA/ritonavir 600/100 mg Twice Daily* of at Least Moderate Intensity (≥ Grade 2) Occurring in ≥2% of Antiretroviral Treatment-Experienced HIV-1-Infected Adult Subjects		
System Organ Class, Preferred Term, %	Randomized Study TMC114-C214	
	PREZISTA/ritonavir 600/100 mg twice daily + OBR N=298	lopinavir/ritonavir 400/100 mg twice daily + OBR N=297
Gastrointestinal Disorders		
Abdominal distension	2%	<1%
Abdominal pain	6%	3%
Diarrhea	14%	20%
Dyspepsia	2%	1%
Nausea	7%	6%
Vomiting	5%	3%
General Disorders and Administration Site Conditions		
Asthenia	3%	1%
Fatigue	2%	1%
Metabolism and Nutrition Disorders		
Anorexia	2%	2%
Diabetes mellitus	2%	<1%
Nervous System Disorders		
Headache	3%	3%
Skin and Subcutaneous Tissue Disorders		
Rash	7%	3%

N=total number of subjects per treatment group

OBR=optimized background regimen

* Excluding laboratory abnormalities reported as ADRs

Less Common Adverse Reactions

Treatment-emergent ADRs of at least moderate intensity (greater than or equal to Grade 2) occurring in less than 2% of antiretroviral treatment-experienced subjects receiving PREZISTA/ritonavir 600/100 mg twice daily are listed below by body system:

Gastrointestinal Disorders: acute pancreatitis, flatulence

Musculoskeletal and Connective Tissue Disorders: myalgia

Psychiatric Disorders: abnormal dreams

Skin and Subcutaneous Tissue Disorders: pruritus, urticaria

Laboratory abnormalities

Selected Grade 2 to 4 laboratory abnormalities that represent a worsening from baseline observed in antiretroviral treatment-experienced adult subjects treated with PREZISTA/ritonavir 600/100 mg twice daily are presented in Table 10.

Table 10: Grade 2 to 4 Laboratory Abnormalities Observed in Antiretroviral Treatment-Experienced HIV-1-Infected Adult Subjects*			
Laboratory Parameter Preferred Term, %	Limit	Randomized Study TMC114-C214	
		PREZISTA/ritonavir 600/100 mg twice daily + OBR	lopinavir/ritonavir 400/100 mg twice daily + OBR
Biochemistry			
Alanine Aminotransferase			
Grade 2	> 2.5 to ≤ 5.0 X ULN	7%	5%
Grade 3	> 5.0 to ≤ 10.0 X ULN	2%	2%
Grade 4	> 10.0 X ULN	1%	2%
Aspartate Aminotransferase			
Grade 2	> 2.5 to ≤ 5.0 X ULN	6%	6%
Grade 3	> 5.0 to ≤ 10.0 X ULN	2%	2%
Grade 4	> 10.0 X ULN	<1%	2%
Alkaline Phosphatase			
Grade 2	> 2.5 to ≤ 5.0 X ULN	<1%	0%
Grade 3	> 5.0 to ≤ 10.0 X ULN	<1%	<1%
Grade 4	> 10.0 X ULN	0%	0%
Hyperbilirubinemia			
Grade 2	> 1.5 to ≤ 2.5 X ULN	<1%	2%
Grade 3	> 2.5 to ≤ 5.0 X ULN	<1%	<1%
Grade 4	> 5.0 X ULN	<1%	0%
Triglycerides			
Grade 2	5.65-8.48 mmol/L 500-750 mg/dL	10%	11%
Grade 3	8.49-13.56 mmol/L 751-1200 mg/dL	7%	10%
Grade 4	> 13.56 mmol/L > 1200 mg/dL	3%	6%
Total Cholesterol			
Grade 2	6.20-7.77 mmol/L 240-300 mg/dL	25%	23%
Grade 3	> 7.77 mmol/L > 300 mg/dL	10%	14%
Low-Density Lipoprotein Cholesterol			
Grade 2	4.13-4.90 mmol/L 160-190 mg/dL	14%	14%
Grade 3	≥ 4.91 mmol/L ≥ 191 mg/dL	8%	9%
Elevated Glucose Levels			
Grade 2	6.95-13.88 mmol/L 126-250 mg/dL	10%	11%
Grade 3	13.89-27.75 mmol/L 251-500 mg/dL	1%	<1%
Grade 4	> 27.75 mmol/L	<1%	0%

	> 500 mg/dL		
Pancreatic Lipase			
Grade 2	> 1.5 to ≤ 3.0 X ULN	3%	4%
Grade 3	> 3.0 to ≤ 5.0 X ULN	2%	<1%
Grade 4	> 5.0 X ULN	<1%	0%
Pancreatic Amylase			
Grade 2	> 1.5 to ≤ 2.0 X ULN	6%	7%
Grade 3	> 2.0 to ≤ 5.0 X ULN	7%	3%
Grade 4	> 5.0 X ULN	0%	0%

N=total number of subjects per treatment group

OBR=optimized background regimen

* Grade 4 data not applicable in Division of AIDS grading scale

6.3 Serious ADRs

The following serious ADRs of at least moderate intensity (greater than or equal to Grade 2) occurred in the Phase 2b studies and Phase 3 studies with PREZISTA/ritonavir: abdominal pain, acute hepatitis, acute pancreatitis, anorexia, asthenia, diabetes mellitus, diarrhea, fatigue, headache, hepatic enzyme increased, hypercholesterolemia, hyperglycemia, hypertriglyceridemia, immune reconstitution syndrome, low density lipoprotein increased, nausea, pancreatic enzyme increased, rash, Stevens-Johnson Syndrome, and vomiting.

6.4 Patients co-infected with hepatitis B and/or hepatitis C virus

In subjects co-infected with hepatitis B or C virus receiving PREZISTA/ritonavir, the incidence of adverse events and clinical chemistry abnormalities was not higher than in subjects receiving PREZISTA/ritonavir who were not co-infected, except for increased hepatic enzymes [see *Warnings and Precautions (5.2)*]. The pharmacokinetic exposure in co-infected subjects was comparable to that in subjects without co-infection.

6.5 Clinical Trials Experience: Pediatric Patients

PREZISTA/ritonavir has been studied in combination with other antiretroviral agents in 3 Phase II trials. TMC114-C212, in which 80 antiretroviral treatment-experienced HIV-1-infected pediatric subjects 6 to less than 18 years of age and weighing at least 20 kg were included, TMC114-C228, in which 21 antiretroviral treatment-experienced HIV-1-infected pediatric subjects 3 to less than 6 years of age and weighing at least 10 kg were included, and TMC114-C230 in which 12 antiretroviral treatment-naïve HIV-1 infected pediatric patients aged from 12 to less than 18 years and weighing at least 40 kg were included. The TMC114-C212 and C228 trials evaluated PREZISTA/ritonavir twice daily dosing and the TMC114-C230 trial evaluated PREZISTA/ritonavir once daily dosing [see *Use in Specific Populations (8.4)* and *Clinical Studies (14.4)*].

Frequency, type, and severity of ADRs in pediatric subjects were comparable to those observed in adults.

Study TMC114-C212

Clinical ADRs to PREZISTA/ritonavir (all grades, greater than or equal to 3%), were vomiting (13%), diarrhea (11%), abdominal pain (10%), headache (9%), rash (5%), nausea (4%) and fatigue (3%).

Grade 3 or 4 laboratory abnormalities were ALT increased (Grade 3: 3%; Grade 4: 1%), AST increased (Grade 3: 1%), pancreatic amylase increased (Grade 3: 4%, Grade 4: 1%), pancreatic lipase increased (Grade 3: 1%), total cholesterol increased (Grade 3: 1%), and LDL increased (Grade 3: 3%).

Study TMC114-C228

Clinical ADRs to PREZISTA/ritonavir (all grades, greater than or equal to 5%), were diarrhea (24%), vomiting (19%), rash (19%), abdominal pain (5%) and anorexia (5%).

There were no Grade 3 or 4 laboratory abnormalities considered as ADRs in this study.

Study TMC114-C230

Clinical ADRs to PREZISTA/ritonavir (all grades, greater than or equal to 3%), were vomiting (33%), nausea (25%), diarrhea (16.7%), abdominal pain (8.3%), decreased appetite (8.3%), pruritus (8.3%), and rash (8.3%).

There were no Grade 3 or 4 laboratory abnormalities considered as ADRs in this study.

6.6 Postmarketing Experience

The following events have been identified during post approval use of PREZISTA. Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Redistribution of body fat has been reported.

Rarely, rhabdomyolysis (associated with co-administration with HMG-CoA reductase inhibitors and PREZISTA/ritonavir) has been reported.

In addition, toxic epidermal necrolysis, acute generalized exanthematous pustulosis and drug rash with eosinophilia and systemic symptoms have been reported rarely [see *Warnings and Precautions (5.3)*].

7 DRUG INTERACTIONS

See also *Contraindications (4)* and *Clinical Pharmacology (12.3)*.

7.1 Potential for PREZISTA/ritonavir to Affect Other Drugs

PREZISTA co-administered with ritonavir is an inhibitor of CYP3A, CYP2D6, and P-gp. Co-administration of PREZISTA and ritonavir with drugs that are primarily metabolized by CYP3A and CYP2D6, or are transported by P-gp may result in increased plasma concentrations of such drugs, which could increase or prolong their therapeutic effect and adverse events (see Table 11).

7.2 Potential for Other Drugs to Affect Darunavir

Darunavir and ritonavir are metabolized by CYP3A. *In vitro* data indicate that darunavir may be a P-gp substrate. Drugs that induce CYP3A activity would be expected to increase the clearance of darunavir and ritonavir, resulting in lowered plasma concentrations of darunavir and ritonavir. Co-administration of darunavir and ritonavir and other drugs that inhibit CYP3A, or P-gp may decrease the clearance of darunavir and ritonavir and may result in increased plasma concentrations of darunavir and ritonavir (see Table 11).

7.3 Established and Other Potentially Significant Drug Interactions

Table 11 provides dosing recommendations as a result of drug interactions with PREZISTA/ritonavir. These recommendations are based on either drug interaction studies or predicted interactions due to the expected magnitude of interaction and potential for serious adverse events or loss of efficacy.

Table 11: Established and Other Potentially Significant Drug Interactions: Alterations in Dose or Regimen May Be Recommended Based on Drug Interaction Studies or Predicted Interaction [See <i>Clinical Pharmacology</i> (12.3) for Magnitude of Interaction, Tables 14 and 15]		
Concomitant Drug Class: Drug Name	Effect on Concentration of Darunavir or Concomitant Drug	Clinical Comment
HIV-1-Antiviral Agents: Nucleoside Reverse Transcriptase Inhibitors (NRTIs)		
didanosine	↔ darunavir ↔ didanosine	Didanosine should be administered one hour before or two hours after PREZISTA/ritonavir (which are administered with food).
HIV-1-Antiviral Agents: HIV-Protease Inhibitors (PIs)		
indinavir (The reference regimen for indinavir was indinavir/ritonavir 800/100 mg twice daily.)	↑ darunavir ↑ indinavir	The appropriate dose of indinavir in combination with PREZISTA/ritonavir has not been established.
lopinavir/ritonavir	↓ darunavir ↔ lopinavir	Appropriate doses of the combination have not been established. Hence, it is not

		recommended to co-administer lopinavir/ritonavir and PREZISTA, with or without ritonavir.
saquinavir	↓ darunavir ↔ saquinavir	Appropriate doses of the combination have not been established. Hence, it is not recommended to co-administer saquinavir and PREZISTA, with or without ritonavir.
HIV-1-Antiviral Agents: CCR5 co-receptor antagonists		
maraviroc	↑ maraviroc	When used in combination with PREZISTA/ritonavir, the dose of maraviroc should be 150 mg twice daily.
Other Agents		
Antiarrhythmics: e.g. amiodarone, bepidil, disopyramide, flecainide, lidocaine (systemic), mexiletine, propafenone, quinidine, digoxin	↑ antiarrhythmics ↑ digoxin	Therapeutic concentration monitoring, if available, is recommended for antiarrhythmics when co-administered with PREZISTA/ritonavir. The lowest dose of digoxin should initially be prescribed. The serum digoxin concentrations should be monitored and used for titration of digoxin dose to obtain the desired clinical effect.
Antibacterial: clarithromycin	↔ darunavir ↑ clarithromycin	No dose adjustment of the combination is required for patients with normal renal function. For co-administration of clarithromycin and PREZISTA/ritonavir in patients with renal impairment, the following dose adjustments should be considered: <ul style="list-style-type: none"> • For subjects with CLcr of 30-60 mL/min, the dose of clarithromycin should be reduced by 50%. • For subjects with CLcr of <30 mL/min, the dose of clarithromycin should be reduced by 75%.
Anticoagulant: apixaban, dabigatran etexilate, rivaroxaban	↑ anticoagulant	Concomitant use of apixaban and PREZISTA/ritonavir is not recommended. The combination of PREZISTA/ritonavir and dabigatran etexilate is not recommended in specific renal impairment groups, depending on the indication. Please see the dabigatran US prescribing information for specific recommendations. Co-administration of PREZISTA/ritonavir and rivaroxaban is not recommended.

warfarin	↓ warfarin ↔ darunavir	Warfarin concentrations are decreased when co-administered with PREZISTA/ritonavir. It is recommended that the international normalized ratio (INR) be monitored when warfarin is combined with PREZISTA/ritonavir.
Anticonvulsant: carbamazepine	↔ darunavir ↑ carbamazepine	The dose of either darunavir/ritonavir or carbamazepine does not need to be adjusted when initiating co-administration with darunavir/ritonavir and carbamazepine. Clinical monitoring of carbamazepine concentrations and its dose titration is recommended to achieve the desired clinical response.
Anticonvulsant: phenobarbital, phenytoin	↔ darunavir ↓ phenytoin ↓ phenobarbital	Phenytoin and phenobarbital levels should be monitored when co-administering with PREZISTA/ritonavir.
Antidepressant: Selective Serotonin Reuptake Inhibitors (SSRIs): paroxetine, sertraline Tricyclic Antidepressants (TCAs): amitriptyline, desipramine imipramine, nortriptyline Other: trazodone	↓ paroxetine ↓ sertraline ↑ amitriptyline ↑ desipramine ↑ imipramine ↑ nortriptyline ↑trazodone	If either sertraline or paroxetine is initiated patients receiving PREZISTA/ritonavir, dose titrating the SSRI based on a clinical assessment of antidepressant response is recommended. Monitor for antidepressant response in patients on a stable dose of sertraline or paroxetine who start treatment with PREZISTA/ritonavir. Use a lower dose of the tricyclic antidepressants and trazodone due to potential increased adverse events such as nausea, dizziness, hypotension and syncope.
Antifungals: itraconazole, ketoconazole, posaconazole voriconazole	↑ darunavir ↑ itraconazole (not studied) ↑ ketoconazole ↔ posaconazole (not studied) ↓ voriconazole (not studied)	Monitor for increased PREZISTA/ritonavir adverse events with concomitant use of itraconazole, ketoconazole, or posaconazole. When co-administration is required, the daily dose of ketoconazole or itraconazole should not exceed 200 mg with monitoring for increased antifungal adverse events. Voriconazole is not recommended for patients receiving darunavir/ritonavir unless an assessment comparing predicted benefit to risk ratio justifies the use of voriconazole.
Anti-gout:	↑ colchicine	<u>Treatment of gout-flares –</u>

colchicine		<p><u>co-administration of colchicine in patients on PREZISTA/ritonavir:</u> 0.6 mg (1 tablet) × 1 dose, followed by 0.3 mg (half tablet) 1 hour later. Treatment course to be repeated no earlier than 3 days.</p> <p><u>Prophylaxis of gout-flares – co-administration of colchicine in patients on PREZISTA/ritonavir:</u> If the original regimen was 0.6 mg twice a day, the regimen should be adjusted to 0.3 mg once a day. If the original regimen was 0.6 mg once a day, the regimen should be adjusted to 0.3 mg once every other day.</p> <p><u>Treatment of familial Mediterranean fever – co-administration of colchicine in patients on PREZISTA/ritonavir:</u> maximum daily dose of 0.6 mg (may be given as 0.3 mg twice a day).</p> <p>Co-administration of PREZISTA/ritonavir with colchicine in patients with renal or hepatic impairment is CONTRAINDICATED.</p>
<p>Antimalarials: artemether/lumefantrine</p>	<p>↓ artemether ↓ dihydroartemisinin ↑ lumefantrine ↔ darunavir</p>	<p>The combination of PREZISTA/ritonavir and artemether/lumefantrine can be used without dose adjustments. However, the combination should be used with caution as increased lumefantrine exposure may increase the risk of QT prolongation.</p>
<p>Antimycobacterial: rifabutin</p> <p>The reference regimen for rifabutin was 300 mg once daily</p> <p>rifapentine</p>	<p>↑ darunavir ↑ rifabutin ↑ 25-O-desacetyl-rifabutin</p> <p>↓ darunavir</p>	<p>Dose reduction of rifabutin by at least 75% of the usual dose (300 mg once daily) is recommended (i.e., a maximum dose of 150 mg every other day). Increased monitoring for adverse events is warranted in patients receiving this combination and further dose reduction of rifabutin may be necessary.</p> <p>Co-administration of PREZISTA/ritonavir with rifapentine is not recommended.</p>
<p>Antineoplastics: dasatinib nilotinib vinblastine vincristine</p>	<p>↑ Antineoplastics</p>	<p>A decrease in the dosage or an adjustment of the dosing interval of dasatinib and nilotinib may be necessary for patients. Please refer to the dasatinib and nilotinib prescribing information for dosing instructions.</p> <p>For vincristine and vinblastine, consideration should be given to temporarily withholding the ritonavir-</p>

		containing antiretroviral regimen in patients who develop significant hematologic or gastrointestinal side effects when PREZISTA/ritonavir is administered concurrently with vincristine or vinblastine. If the antiretroviral regimen must be withheld for a prolonged period, consideration should be given to initiating a revised regimen that does not include a CYP3A or P-gp inhibitor.
Antipsychotics: quetiapine e.g. risperidone, thioridazine	↑ quetiapine ↑ antipsychotics	<u>Initiation of PREZISTA with ritonavir in patients taking quetiapine:</u> Consider alternative antiretroviral therapy to avoid increases in quetiapine exposures. If co-administration is necessary, reduce the quetiapine dose to 1/6 of the current dose and monitor for quetiapine-associated adverse reactions. Refer to the quetiapine prescribing information for recommendations on adverse reaction monitoring. <u>Initiation of quetiapine in patients taking PREZISTA with ritonavir:</u> Refer to the quetiapine prescribing information for initial dosing and titration of quetiapine. A dose decrease may be needed for these drugs when co-administered with PREZISTA/ritonavir.
β-Blockers: e.g. carvedilol, metoprolol, timolol	↑ beta-blockers	Clinical monitoring of patients is recommended. A dose decrease may be needed for these drugs when co-administered with PREZISTA/ritonavir and a lower dose of the beta blocker should be considered.
Calcium Channel Blockers: amlodipine, diltiazem, felodipine, nicardipine nifedipine, verapamil	↑ calcium channel blockers	Clinical monitoring of patients is recommended.
Corticosteroid (systemic): dexamethasone Corticosteroid (systemic): metabolized by CYP3A e.g. budesonide	↓ darunavir ↑ corticosteroid	Systemic dexamethasone induces CYP3A and can thereby decrease darunavir plasma concentrations. This may result in loss of therapeutic effect to PREZISTA. Concomitant use of corticosteroids metabolized by CYP3A may increase the

prednisolone		risk for development of systemic corticosteroid effects including Cushing's syndrome and adrenal suppression. Alternatives should be considered, particularly for long-term use.
Corticosteroid (inhaled/nasal): budesonide, fluticasone	↑ corticosteroid	Alternatives should be considered, particularly for long-term use.
Endothelin receptor antagonists: bosentan	↑ bosentan	<u>Co-administration of bosentan in patients on PREZISTA/ritonavir:</u> In patients who have been receiving PREZISTA/ritonavir for at least 10 days, start bosentan at 62.5 mg once daily or every other day based upon individual tolerability. <u>Co-administration of PREZISTA/ritonavir in patients on bosentan:</u> Discontinue use of bosentan at least 36 hours prior to initiation of PREZISTA/ritonavir. After at least 10 days following the initiation of PREZISTA/ritonavir, resume bosentan at 62.5 mg once daily or every other day based upon individual tolerability.
Hepatitis C Virus (HCV) Direct-Acting Agents: NS3-4A protease inhibitors: boceprevir telaprevir simeprevir	↓ darunavir ↓ boceprevir ↓ telaprevir ↑ simeprevir ↑ darunavir	Co-administration of boceprevir or telaprevir and PREZISTA/ritonavir is not recommended. Co-administration of darunavir/ritonavir and simeprevir is not recommended
HMG-CoA Reductase Inhibitors: atorvastatin, pravastatin, rosuvastatin	↑ HMG-CoA Reductase Inhibitors	Co-administration of PREZISTA/ritonavir with HMG-Co A reductase inhibitors may lead to adverse events such as myopathy. Titrate atorvastatin, pravastatin or rosuvastatin dose carefully and use the lowest necessary dose while monitoring for adverse events. Do not exceed atorvastatin 20 mg/day.
Immunosuppressants: e.g. cyclosporine, tacrolimus, sirolimus Immunosuppressants/neoplastic: everolimus	↑ immunosuppressants	Therapeutic concentration monitoring of the immunosuppressive agent is recommended when co-administered with PREZISTA/ritonavir. Co-administration of everolimus and PREZISTA/ritonavir is not recommended.
Inhaled beta agonist: salmeterol	↑ salmeterol	Co-administration of salmeterol and PREZISTA/ritonavir is not recommended. The combination may result in increased risk of cardiovascular adverse events

		associated with salmeterol, including QT prolongation, palpitations and sinus tachycardia.
Narcotic Analgesic/Treatment of Opioid Dependence: buprenorphine, buprenorphine/naloxone, methadone	↔ buprenorphine, naloxone ↑ norbuprenorphine (metabolite) ↓ methadone	No adjustment of methadone dosage is required when initiating co-administration of PREZISTA/ritonavir. However, clinical monitoring is recommended as the dose of methadone during maintenance therapy may need to be adjusted in some patients. No dose adjustment for buprenorphine or buprenorphine/naloxone is required with concurrent administration of PREZISTA/ritonavir. Clinical monitoring is recommended if PREZISTA/ritonavir and buprenorphine or buprenorphine/naloxone are coadministered.
Oral Contraceptives/estrogen: ethinyl estradiol, norethindrone	↓ ethinyl estradiol ↓ norethindrone	Alternative methods of nonhormonal contraception are recommended.
PDE-5 inhibitors: e.g. avanafil, sildenafil, tadalafil, vardenafil	↑ PDE-5 inhibitors (only the use of sildenafil at doses used for treatment of erectile dysfunction has been studied with PREZISTA/ritonavir)	Co-administration with PREZISTA/ritonavir may result in an increase in PDE-5 inhibitor-associated adverse events, including hypotension, syncope, visual disturbances and priapism. <u>Use of PDE-5 inhibitors for pulmonary arterial hypertension (PAH):</u> <ul style="list-style-type: none"> • Use of sildenafil is contraindicated when used for the treatment of pulmonary arterial hypertension (PAH) [see <i>Contraindications (4)</i>]. • The following dose adjustments are recommended for use of tadalafil with PREZISTA/ritonavir: <u>Co-administration of tadalafil in patients on PREZISTA/ritonavir:</u> In patients receiving PREZISTA/ritonavir for at least one week, start tadalafil at 20 mg once daily. Increase to 40 mg once daily based upon individual tolerability. <u>Co-administration of PREZISTA/ritonavir in patients on tadalafil:</u> Avoid use of tadalafil during the initiation of PREZISTA/ritonavir. Stop tadalafil at least 24 hours prior to starting PREZISTA/ritonavir. After at least one week following the initiation of PREZISTA/ritonavir, resume tadalafil at 20 mg once daily. Increase to 40 mg once daily based upon individual tolerability.

		<p><u>Use of PDE-5 inhibitors for erectile dysfunction:</u> Sildenafil at a single dose not exceeding 25 mg in 48 hours, vardenafil at a single dose not exceeding 2.5 mg dose in 72 hours, or tadalafil at a single dose not exceeding 10 mg dose in 72 hours can be used with increased monitoring for PDE-5 inhibitor-associated adverse events.</p> <p>Co-administration of PREZISTA/ritonavir and avanafil is not recommended.</p>
<p>Sedatives/Hypnotics: Metabolized by CYP3A e.g. buspirone, diazepam, estazolam, parenterally administered midazolam, zoldipem</p>	<p>↑ sedatives/hypnotics</p>	<p>Titration is recommended when co-administering PREZISTA/ritonavir with sedatives/hypnotics metabolized by CYP3A and a lower dose of the sedatives/hypnotics should be considered with monitoring for adverse events.</p> <p>Co-administration of parenteral midazolam should be done in a setting which ensures close clinical monitoring and appropriate medical management in case of respiratory depression and/or prolonged sedation. Dosage reduction for midazolam should be considered, especially if more than a single dose of midazolam is administered. Co-administration of oral midazolam or triazolam with PREZISTA/ritonavir is CONTRAINDICATED.</p>

In addition to the drugs included in Table 11, the interaction between PREZISTA/ritonavir and the following drugs were evaluated in clinical studies and no dose adjustments are needed for either drug [see *Clinical Pharmacology (12.3)*]: atazanavir, efavirenz, etravirine, nevirapine, omeprazole, ranitidine, and rilpivirine. Using cross-trial comparisons to historical pharmacokinetic data, dolutegravir did not appear to affect the pharmacokinetics of darunavir. Darunavir/ritonavir had no clinically significant effect on the pharmacokinetics of dolutegravir. Pitavastatin had no clinically significant effect on the pharmacokinetics of darunavir/ritonavir.

Acid modifying medications are not predicted to alter darunavir exposure.

Other nucleoside reverse transcriptase inhibitors (NRTIs):

Based on the different elimination pathways of the other NRTIs (abacavir, emtricitabine, lamivudine, stavudine, tenofovir disoproxil fumarate, zidovudine) that are primarily renally excreted, no drug interactions are expected for these drugs and PREZISTA/ritonavir.

Other PIs:

The co-administration of PREZISTA/ritonavir and PIs other than lopinavir/ritonavir, saquinavir, atazanavir, and indinavir has not been studied. Therefore, such co-administration is not recommended.

Integrase strand transfer inhibitors:

No dose adjustments are needed for either raltegravir or darunavir based on pharmacokinetic data from literature references. Please refer to the elvitegravir/cobicistat/emtricitabine/tenofovir disoproxil fumarate U.S. prescribing information for information regarding concomitant use with other antiretroviral medications.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C: PREZISTA should be used during pregnancy only if the potential benefit justifies the potential risk.

No adequate and well-controlled studies have been conducted in pregnant women. Reproduction studies conducted with darunavir showed no embryotoxicity or teratogenicity in mice and rats in the presence or absence of ritonavir as well as in rabbits with darunavir alone. In these studies, darunavir exposures (based on AUC) were higher in rats (3-fold), whereas in mice and rabbits, exposures were lower (less than 1-fold) compared to those obtained in humans at the recommended clinical dose of darunavir boosted with ritonavir.

In the rat pre- and postnatal development study, a reduction in pup body weight gain was observed with darunavir alone or in combination with ritonavir during lactation. This was due to exposure of pups to drug substances via the milk. Sexual development, fertility and mating performance of offspring were not affected by maternal treatment with darunavir alone or in combination with ritonavir. The maximal plasma exposures achieved in rats were approximately 50% of those obtained in humans at the recommended clinical dose boosted with ritonavir.

In the juvenile toxicity study where rats were directly dosed with darunavir, deaths occurred from post-natal day 5 through 11 at plasma exposure levels ranging from 0.1 to 1.0 of the human exposure levels. In a 4-week rat toxicology study, when dosing was initiated on post-natal day 23 (the human equivalent of 2 to 3 years of age), no deaths were observed with a plasma exposure (in combination with ritonavir) of 0.1 of the human plasma exposure levels.

Antiretroviral Pregnancy Registry: *To monitor maternal-fetal outcomes of pregnant women exposed to PREZISTA, an Antiretroviral Pregnancy Registry has been established. Physicians are encouraged to register patients by calling 1-800-258-4263.*

8.3 Nursing Mothers

The Centers for Disease Control and Prevention recommend that HIV-infected mothers in the United States not breastfeed their infants to avoid risking postnatal transmission of HIV. Although it is not known whether darunavir is secreted in human milk, darunavir is secreted into the milk of lactating rats. Because of both the potential for HIV transmission and the potential for serious adverse reactions in nursing infants, **mothers should be instructed not to breastfeed if they are receiving PREZISTA.**

8.4 Pediatric Use

Do not administer PREZISTA/ritonavir in pediatric patients below 3 years of age because of toxicity and mortality observed in juvenile rats dosed with darunavir (from 20 mg/kg to 1000 mg/kg) up to days 23 to 26 of age [see *Warnings and Precautions (5.11)*, *Use in Specific Populations (8.1)*, *Clinical Pharmacology (12.3)* and *Nonclinical Toxicology (13.2)*].

The safety, pharmacokinetic profile, and virologic and immunologic responses of PREZISTA/ritonavir were evaluated in treatment-experienced HIV-1-infected pediatric subjects 3 to less than 18 years of age and weighing at least 10 kg. These subjects were evaluated in clinical trials TMC114-C212 (80 subjects, 6 to less than 18 years of age) and TMC114-228 (21 subjects, 3 to less than 6 years of age) [see *Adverse Reactions (6.5)*, *Clinical Pharmacology (12.3)* and *Clinical Studies (14.4)*]. Frequency, type, and severity of adverse drug reactions in pediatric subjects were comparable to those observed in adults [see *Adverse Reactions (6.1, 6.2)*]. Please see *Dosage and Administration (2.2)* for twice-daily dosing recommendations for pediatric subjects 3 to less than 18 years of age and weighing at least 10 kg.

In clinical trial TMC114-C230, the safety, pharmacokinetic profile and virologic and immunologic responses of PREZISTA/ritonavir administered once daily were evaluated in treatment-naïve HIV-1 infected pediatric subjects 12 to less than 18 years of age (12 subjects) [see *Adverse Reactions (6.5)*, *Clinical Pharmacology (12.3)* and *Clinical Studies (14.4)*]. Frequency, type, and severity of adverse drug reactions in pediatric subjects were comparable to those observed in adults [see *Adverse Reactions (6.1, 6.2)*]. Once daily dosing recommendations for pediatric patients 3 to less than 12 years of age were derived using population pharmacokinetic modeling and simulation. Although a PREZISTA/ritonavir once daily dosing pediatric trial was not conducted in children less than 12 years of age, there is sufficient clinical safety data to support the predicted PREZISTA exposures for the dosing recommendations in this age group [see *Clinical Pharmacology (12.3)*]. Please see *Dosage and Administration (2.2)* for once-daily dosing recommendations for pediatric subjects 3 to less than 18 years of age and weighing at least 10 kg.

8.5 Geriatric Use

Clinical studies of PREZISTA did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients. In general, caution should be exercised in the administration and monitoring of PREZISTA in elderly patients, reflecting the greater frequency of decreased hepatic function, and of concomitant disease or other drug therapy [see *Clinical Pharmacology (12.3)*].

8.6 Hepatic Impairment

No dose adjustment of PREZISTA/ritonavir is necessary for patients with either mild or moderate hepatic impairment. No pharmacokinetic or safety data are available regarding the use of PREZISTA/ritonavir in subjects with severe hepatic impairment. Therefore, PREZISTA/ritonavir is not recommended for use in patients with severe hepatic impairment [see *Dosage and Administration (2.3)* and *Clinical Pharmacology (12.3)*].

8.7 Renal Impairment

Population pharmacokinetic analysis showed that the pharmacokinetics of darunavir were not significantly affected in HIV-infected subjects with moderate renal impairment (CrCL between 30-60 mL/min, n=20). No pharmacokinetic data are available in HIV-1-infected patients with severe renal impairment or end stage renal disease; however, because the renal clearance of darunavir is limited, a decrease in total body clearance is not expected in patients with renal impairment. As darunavir and ritonavir are highly bound to plasma proteins, it is unlikely that they will be significantly removed by hemodialysis or peritoneal dialysis [see *Clinical Pharmacology (12.3)*].

10 OVERDOSAGE


Human experience of acute overdose with PREZISTA/ritonavir is limited. Single doses up to 3200 mg of the oral solution of darunavir alone and up to 1600 mg of the tablet formulation of darunavir in combination with ritonavir have been administered to healthy volunteers without untoward symptomatic effects.

No specific antidote is available for overdose with PREZISTA. Treatment of overdose with PREZISTA consists of general supportive measures including monitoring of vital signs and observation of the clinical status of the patient. If indicated, elimination of unabsorbed active substance is to be achieved by emesis. Administration of activated charcoal may also be used to aid in removal of unabsorbed active substance. Since PREZISTA is highly protein bound, dialysis is unlikely to be beneficial in significant removal of the active substance.

11 DESCRIPTION

PREZISTA (darunavir) is an inhibitor of the human immunodeficiency virus (HIV-1) protease.

PREZISTA (darunavir), in the form of darunavir ethanolate, has the following chemical name: [(1S,2R)-3-[[[(4-aminophenyl)sulfonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-carbamic acid (3R,3aS,6aR)-hexahydrofuro[2,3-b]furan-3-yl ester monoethanolate. Its molecular formula is $C_{27}H_{37}N_3O_7S \cdot C_2H_5OH$ and its molecular weight is 593.73. Darunavir ethanolate has the following structural formula:



Darunavir ethanolate is a white to off-white powder with a solubility of approximately 0.15 mg/mL in water at 20°C.

PREZISTA 100 mg/mL oral suspension is available as a white to off-white opaque suspension for oral administration.

PREZISTA 75 mg tablets are available as white, caplet-shaped, film-coated tablets for oral administration. PREZISTA 150 mg tablets are available as white, oval-shaped, film-coated tablets for oral administration. PREZISTA 600 mg tablets are available as orange, oval-shaped, film-coated tablets for oral administration.

PREZISTA 800 mg tablets are available as dark red, oval-shaped, film-coated tablets for oral administration.

Each mL of the oral suspension contains darunavir ethanolate equivalent to 100 mg darunavir. In addition, each mL contains the inactive ingredients hydroxypropyl cellulose, microcrystalline cellulose, sodium carboxymethylcellulose, methylparaben sodium, citric acid monohydrate, sucralose, masking flavor, strawberry cream flavor, hydrochloric acid (for pH adjustment) and purified water.

Each 75 mg tablet contains darunavir ethanolate equivalent to 75 mg of darunavir. Each 150 mg tablet contains darunavir ethanolate equivalent to 150 mg of darunavir. Each 600 mg tablet contains darunavir ethanolate equivalent to 600 mg of darunavir. Each 800 mg tablet contains darunavir ethanolate equivalent to 800 mg of darunavir. During storage, partial conversion from ethanolate to hydrate may occur; however, this does not affect product quality or performance. Each tablet also contains the inactive ingredients colloidal silicon dioxide, crospovidone, magnesium stearate, and microcrystalline cellulose. The 800 mg tablet also contains hypromellose. The 75 and 150 mg tablet film coating, OPADRY® White, contains polyethylene glycol 3350, polyvinyl alcohol-

partially hydrolyzed, talc, and titanium dioxide. The 600 mg tablet film coating, OPADRY® Orange, contains FD&C Yellow No. 6, polyethylene glycol 3350, polyvinyl alcohol-partially hydrolyzed, talc, and titanium dioxide. The 800 mg tablet film coating, OPADRY® Dark Red, contains iron oxide red, polyethylene glycol 3350, polyvinyl alcohol-partially hydrolyzed, talc, and titanium dioxide.

All dosages for PREZISTA are expressed in terms of the free form of darunavir.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Darunavir is an HIV-1 antiviral drug [see *Microbiology (12.4)*].

12.2 Pharmacodynamics

In an open-label, randomized, placebo- and active-controlled, four-way crossover trial, 40 healthy subjects were administered supratherapeutic doses of darunavir/ritonavir 1600/100 mg once daily and 800/100 mg twice daily for seven days.

At the mean maximum darunavir concentration of 6599 ng/mL observed in this study, the mean increase in QTcF was 2.2 ms with a 90% two-sided confidence interval (CI) of -2.0 to 6.3 ms. When evaluating the 2-sided 90% CI on the time-matched mean changes in QTcF versus placebo control, the upper bounds of both darunavir/ritonavir groups never exceeded the 10 ms boundary. In the setting of this trial, darunavir/ritonavir did not appear to prolong the QTc interval.

12.3 Pharmacokinetics

Pharmacokinetics in Adults

General

Darunavir is primarily metabolized by CYP3A. Ritonavir inhibits CYP3A, thereby increasing the plasma concentrations of darunavir. When a single dose of PREZISTA 600 mg was given orally in combination with 100 mg ritonavir twice daily, there was an approximate 14-fold increase in the systemic exposure of darunavir. Therefore, PREZISTA should only be used in combination with 100 mg of ritonavir to achieve sufficient exposures of darunavir.

The pharmacokinetics of darunavir, co-administered with low dose ritonavir (100 mg), has been evaluated in healthy adult volunteers and in HIV-1-infected subjects. Table 12 displays the population pharmacokinetic estimates of darunavir after oral administration of PREZISTA/ritonavir 600/100 mg twice daily [based on sparse sampling in 285 patients in study TMC114-C214, 278 patients in Study TMC114-C229 and 119 patients (integrated data) from Studies TMC114-C202 and TMC114-C213] and PREZISTA/ritonavir 800/100 mg once daily [based on sparse sampling in 335 patients in

Study TMC114-C211 and 280 patients in Study TMC114-C229] to HIV-1-infected patients.

Table 12: Population Pharmacokinetic Estimates of Darunavir at PREZISTA/ritonavir 800/100 mg Once Daily (Study TMC114-C211, 48-Week Analysis and Study TMC114-C229, 48-Week Analysis) and PREZISTA/ritonavir 600/100 mg Twice Daily (Study TMC114-C214, 48-Week Analysis, Study TMC114-C229, 48-Week Analysis and Integrated Data from Studies TMC114-C213 and TMC114-C202, Primary 24-Week Analysis)					
Parameter	Study TMC114-C211 PREZISTA / ritonavir 800/100 mg once daily N=335	Study TMC114-C229 PREZISTA / ritonavir 800/100 mg once daily N=280	Study TMC114-C214 PREZISTA / ritonavir 600/100 mg twice daily N=285	Study TMC114-C229 PREZISTA / ritonavir 600/100 mg twice daily N=278	Studies TMC114-C213 and TMC114- C202 (integrated data) PREZISTA / ritonavir 600/100 mg twice daily N=119
AUC_{24h} (ng·h/mL)*					
Mean ± Standard Deviation	93026 ± 27050	93334 ± 28626	116796 ± 33594	114302 ± 32681	124698 ± 32286
Median (Range)	87854 (45000-219240)	87788 (45456-236920)	111632 (64874-355360)	109401 (48934-323820)	123336 (67714-212980)
C_{0h} (ng/mL)					
Mean ± Standard Deviation	2282 ± 1168	2160 ± 1201	3490 ± 1401	3386 ± 1372	3578 ± 1151
Median (Range)	2041 (368-7242)	1896 (184-7881)	3307 (1517-13198)	3197 (250-11865)	3539 (1255-7368)

N=number of subjects with data

* AUC_{24h} is calculated as AUC_{12h}*2

Absorption and Bioavailability

Darunavir, co-administered with 100 mg ritonavir twice daily, was absorbed following oral administration with a T_{max} of approximately 2.5-4 hours. The absolute oral bioavailability of a single 600 mg dose of darunavir alone and after co-administration with 100 mg ritonavir twice daily was 37% and 82%, respectively. *In vivo* data suggest that darunavir/ritonavir is an inhibitor of the p-glycoprotein (p-gp) transporters.

Effects of Food on Oral Absorption

When PREZISTA tablets were administered with food, the C_{max} and AUC of darunavir, co-administered with ritonavir, is approximately 40% higher relative to the fasting state. Therefore, PREZISTA tablets, co-administered with ritonavir, should always be taken with food. Within the range of meals studied, darunavir exposure is similar. The total caloric content of the various meals evaluated ranged from 240 Kcal (12 gms fat) to 928 Kcal (56 gms fat).

Distribution

Darunavir is approximately 95% bound to plasma proteins. Darunavir binds primarily to plasma alpha 1-acid glycoprotein (AAG).

Metabolism

In vitro experiments with human liver microsomes (HLMs) indicate that darunavir primarily undergoes oxidative metabolism. Darunavir is extensively metabolized by CYP enzymes, primarily by CYP3A. A mass balance study in healthy volunteers showed that after a single dose administration of 400 mg ¹⁴C-darunavir, co-administered with 100 mg ritonavir, the majority of the radioactivity in the plasma was due to darunavir. At least 3 oxidative metabolites of darunavir have been identified in humans; all showed activity that was at least 90% less than the activity of darunavir against wild-type HIV-1.

Elimination

A mass balance study in healthy volunteers showed that after single dose administration of 400 mg ¹⁴C-darunavir, co-administered with 100 mg ritonavir, approximately 79.5% and 13.9% of the administered dose of ¹⁴C-darunavir was recovered in the feces and urine, respectively. Unchanged darunavir accounted for approximately 41.2% and 7.7% of the administered dose in feces and urine, respectively. The terminal elimination half-life of darunavir was approximately 15 hours when co-administered with ritonavir. After intravenous administration, the clearance of darunavir, administered alone and co-administered with 100 mg twice daily ritonavir, was 32.8 L/h and 5.9 L/h, respectively.

Special Populations

Hepatic Impairment

Darunavir is primarily metabolized by the liver. The steady-state pharmacokinetic parameters of darunavir were similar after multiple dose co-administration of PREZISTA/ritonavir 600/100 mg twice daily to subjects with normal hepatic function (n=16), mild hepatic impairment (Child-Pugh Class A, n=8), and moderate hepatic impairment (Child-Pugh Class B, n=8). The effect of severe hepatic impairment on the pharmacokinetics of darunavir has not been evaluated [see *Dosage and Administration* (2.3) and *Use in Specific Populations* (8.6)].

Hepatitis B or Hepatitis C Virus Co-infection

The 48-week analysis of the data from Studies TMC114-C211 and TMC114-C214 in HIV-1-infected subjects indicated that hepatitis B and/or hepatitis C virus co-infection status had no apparent effect on the exposure of darunavir.

Renal Impairment

Results from a mass balance study with ¹⁴C-darunavir/ritonavir showed that approximately 7.7% of the administered dose of darunavir is excreted in the urine as unchanged drug. As darunavir and ritonavir are highly bound to plasma proteins, it is unlikely that they will be significantly removed by hemodialysis or peritoneal dialysis. Population pharmacokinetic analysis showed that the pharmacokinetics of darunavir were not significantly affected in HIV-1-infected subjects with moderate renal impairment

(CrCL between 30-60 mL/min, n=20). There are no pharmacokinetic data available in HIV-1-infected patients with severe renal impairment or end stage renal disease [see *Use in Specific Populations* (8.7)].

Gender

Population pharmacokinetic analysis showed higher mean darunavir exposure in HIV-1-infected females compared to males. This difference is not clinically relevant.

Race

Population pharmacokinetic analysis of darunavir in HIV-1-infected subjects indicated that race had no apparent effect on the exposure to darunavir.

Geriatric Patients

Population pharmacokinetic analysis in HIV-1-infected subjects showed that darunavir pharmacokinetics are not considerably different in the age range (18 to 75 years) evaluated in HIV-1-infected subjects (n=12, age greater than or equal to 65) [see *Use in Specific Populations* (8.5)].

Pediatric Patients

PREZISTA/ritonavir administered twice daily:

The pharmacokinetics of darunavir in combination with ritonavir in 93 antiretroviral treatment-experienced HIV-1-infected pediatric subjects 3 to less than 18 years of age and weighing at least 10 kg showed that the administered weight-based dosages resulted in similar darunavir exposure when compared to the darunavir exposure achieved in treatment-experienced adults receiving PREZISTA/ritonavir 600/100 mg twice daily [see *Dosage and Administration* (2.2)].

PREZISTA/ritonavir administered once daily:

The pharmacokinetics of darunavir in combination with ritonavir in 12 antiretroviral treatment-naïve HIV-1-infected pediatric subjects 12 to less than 18 years of age and weighing at least 40 kg receiving PREZISTA/ritonavir 800/100 mg once daily resulted in similar darunavir exposures when compared to the darunavir exposure achieved in treatment-naïve adults receiving PREZISTA/ritonavir 800/100 mg once daily. [see *Dosage and Administration* (2.2)].

Based on population pharmacokinetic modeling and simulation, the proposed PREZISTA/ritonavir once daily dosing regimens for pediatric patients 3 to less than 12 years of age is predicted to result in similar darunavir exposures when compared to the darunavir exposures achieved in treatment-naïve adults receiving PREZISTA/ritonavir 800/100 mg once daily [see *Dosage and Administration* (2.2)].

The population pharmacokinetic parameters in pediatric subjects with PREZISTA/ritonavir administered once or twice daily are summarized in the table below.

Table 13: Population Pharmacokinetic Estimates of Darunavir Exposure (Study TMC114-C230, Study TMC114-C212 and Study TMC114-C228) Following Administration of Doses in Tables 2 and 3				
Parameter	Study TMC114-C230 PREZISTA/ ritonavir once daily^β N=12	Study TMC114-C212 PREZISTA/ritonavir twice daily N=74	Study TMC114-C228 PREZISTA/ritonavir twice daily*	
			10 to less than 15 kg[‡] N=10	15 to less than 20 kg[§] N=13
AUC_{24h} (ng·h/mL) †				
Mean ± Standard Deviation	84390 ± 23587	126377 ± 34356	137896 ± 51420	157760 ± 54080
Median (Range)	86741 (35527–123325)	127340 (67054–230720)	124044 (89688–261090)	132698 (112310–294840)
C_{0h} (ng/mL)				
Mean ± Standard Deviation	2141 ± 865	3948 ± 1363	4510 ± 2031	4848 ± 2143
Median (Range)	2234 (542–3776)	3888 (1836–7821)	4126 (2456–9361)	3927 (3046–10292)

N=number of subjects with data.

* Subjects may have contributed pharmacokinetic data to both the 10 kg to less than 15 kg weight group and the 15 kg to less than 20 kg weight group.

† AUC_{24h} is calculated as AUC_{12h}*2

‡ Calculated from individual pharmacokinetic parameters estimated for Week 2 and Week 4, based on the Week 48 analysis that evaluated a darunavir dose of 20 mg/kg twice daily with ritonavir 3 mg/kg twice daily.

§ The 15 kg to less than 20 kg weight group received 380 mg (3.8 mL) PREZISTA oral suspension twice daily with 48 mg (0.6 mL) ritonavir oral solution twice daily in TMC114-C228. Calculated from individual pharmacokinetic parameters estimated for Week 2 post-dose adjustment visit; Week 24 and Week 48 based on the – Week 48 analysis that evaluated a darunavir dose of 380 mg twice daily.

β Summary statistics for population pharmacokinetic parameter estimates for DRV after administration of DRV/rtv at 800/100 mg q.d. in treatment-naïve HIV-1 infected subjects from 12 to <18 years of age – Week-48 Analyses

Drug Interactions

[See also *Contraindications (4)*, *Warnings and Precautions (5.5)*, and *Drug Interactions (7)*.]

Darunavir co-administered with ritonavir is an inhibitor of CYP3A, CYP2D6, and P-gp. Co-administration of darunavir and ritonavir with drugs primarily metabolized by CYP3A and CYP2D6, or are transported by P-gp, may result in increased plasma concentrations of such drugs, which could increase or prolong their therapeutic effect and adverse events.

Darunavir and ritonavir are metabolized by CYP3A. *In vitro* data indicate that darunavir may be a P-gp substrate. Drugs that induce CYP3A activity would be expected to increase the clearance of darunavir and ritonavir, resulting in lowered plasma concentrations of darunavir and ritonavir. Co-administration of darunavir and ritonavir

and other drugs that inhibit CYP3A or P-gp may decrease the clearance of darunavir and ritonavir and may result in increased plasma concentrations of darunavir and ritonavir.

Drug interaction studies were performed with darunavir and other drugs likely to be co-administered and some drugs commonly used as probes for pharmacokinetic interactions. The effects of co-administration of darunavir on the AUC, C_{max} , and C_{min} values are summarized in Table 14 (effect of other drugs on darunavir) and Table 15 (effect of darunavir on other drugs). For information regarding clinical recommendations, see *Drug Interactions (7)*.

Several interaction studies have been performed with a dose other than the recommended dose of the co-administered drug or darunavir; however, the results are applicable to the recommended dose of the co-administered drug and/or darunavir.

Table 14: Drug Interactions: Pharmacokinetic Parameters for <u>Darunavir</u> in the Presence of Co-administered Drugs							
Co-Administered Drug	Dose/Schedule		N	PK	LS Mean Ratio (90% CI) of <u>Darunavir</u> Pharmacokinetic Parameters With/Without Co-administered Drug No Effect =1.00		
	Co-Administered Drug	Darunavir/ritonavir			C_{max}	AUC	C_{min}
Co-Administration With Other HIV Protease Inhibitors							
Atazanavir	300 mg q.d.*	400/100 mg b.i.d.†	13	↔	1.02 (0.96-1.09)	1.03 (0.94-1.12)	1.01 (0.88-1.16)
Indinavir	800 mg b.i.d.	400/100 mg b.i.d.	9	↑	1.11 (0.98-1.26)	1.24 (1.09-1.42)	1.44 (1.13-1.82)
Lopinavir/ Ritonavir	400/100 mg b.i.d. 533/133.3 mg b.i.d.	1200/100 mg b.i.d.‡	14	↓	0.79 (0.67-0.92)	0.62 (0.53-0.73)	0.49 (0.39-0.63)
		1200 mg b.i.d.‡	15	↓	0.79 (0.64-0.97)	0.59 (0.50-0.70)	0.45 (0.38-0.52)
Saquinavir hard gel capsule	1000 mg b.i.d.	400/100 mg b.i.d.	14	↓	0.83 (0.75-0.92)	0.74 (0.63-0.86)	0.58 (0.47-0.72)
Co-Administration With Other HIV Antiretrovirals							
Didanosine	400 mg q.d.	600/100 mg b.i.d.	17	↔	0.93 (0.86-1.00)	1.01 (0.95-1.07)	1.07 (0.95-1.21)
Efavirenz	600 mg q.d.	300/100 mg b.i.d.	12	↓	0.85 (0.72-1.00)	0.87 (0.75-1.01)	0.69 (0.54-0.87)
Etravirine	200 mg b.i.d.	600/100 mg b.i.d.	15	↔	1.11 (1.01-1.22)	1.15 (1.05-1.26)	1.02 (0.90-1.17)
Nevirapine	200 mg b.i.d.	400/100 mg b.i.d.	8	↑	1.40 [§] (1.14-1.73)	1.24 [§] (0.97-1.57)	1.02 [§] (0.79-1.32)
Rilpivirine	150 mg q.d.	800/100 mg q.d.	15	↔	0.90 (0.81-1.00)	0.89 (0.81-0.99)	0.89 (0.68-1.16)
Tenofovir Disoproxil Fumarate	300 mg q.d.	300/100 mg b.i.d.	12	↑	1.16 (0.94-1.42)	1.21 (0.95-1.54)	1.24 (0.90-1.69)

Co-Administration With HCV NS3-4A Protease Inhibitors							
Boceprevir [^]	800 mg three times daily	600/100 mg b.i.d.	11	↓	0.64 (0.58-0.71)	0.56 (0.51-0.61)	0.41 (0.38-0.45)
Telaprevir	750 mg every 8 hours	600/100 mg b.i.d.	11	↓	0.60 (0.56-0.64)	0.60 (0.57-0.63)	0.58 (0.52-0.64)
	1125 mg every 12 hours	600/100 mg b.i.d.	15	↓	0.53 (0.47-0.59)	0.49 (0.43-0.55)	0.42 (0.35-0.51)
Simeprevir	50 mg q.d. [€]	800 mg q.d.	25 ⁺	↑	1.04 (0.99-1.10)	1.18 (1.11-1.25)	1.31 (1.13-1.52)
Co-Administration With Other Drugs							
Artemether/ Lumefantrine	80/480 mg (6 doses at 0, 8, 24, 36, 48, and 60 hours)	600/100 mg b.i.d.	14	↔	1.00 (0.93-1.07)	0.96 (0.90-1.03)	0.87 (0.77-0.98)
Carbamazepine	200 mg b.i.d.	600/100 mg b.i.d.	16	↔	1.04 (0.93-1.16)	0.99 (0.90-1.08)	0.85 (0.73-1.00)
Clarithromycin	500 mg b.i.d.	400/100 mg b.i.d.	17	↔	0.83 (0.72-0.96)	0.87 (0.75-1.01)	1.01 (0.81-1.26)
Ketoconazole	200 mg b.i.d.	400/100 mg b.i.d.	14	↑	1.21 (1.04-1.40)	1.42 (1.23-1.65)	1.73 (1.39-2.14)
Omeprazole	20 mg q.d.	400/100 mg b.i.d.	16	↔	1.02 (0.95-1.09)	1.04 (0.96-1.13)	1.08 (0.93-1.25)
Paroxetine	20 mg q.d.	400/100 mg b.i.d.	16	↔	0.97 (0.92-1.02)	1.02 (0.95-1.10)	1.07 (0.96-1.19)
Pitavastatin	4 mg q.d.	800/100 mg q.d.	27	↔	1.06 (1.00-1.12)	1.03 (0.95-1.12)	NA
Ranitidine	150 mg b.i.d.	400/100 mg b.i.d.	16	↔	0.96 (0.89-1.05)	0.95 (0.90-1.01)	0.94 (0.90-0.99)
Rifabutin	150 mg q.o.d. [¶]	600/100 mg b.i.d.	11	↑	1.42 (1.21-1.67)	1.57 (1.28-1.93)	1.75 (1.28-2.37)
Sertraline	50 mg q.d.	400/100 mg b.i.d.	13	↔	1.01 (0.89-1.14)	0.98 (0.84-1.14)	0.94 (0.76-1.16)

N = number of subjects with data

* q.d. = once daily

† b.i.d. = twice daily

‡ The pharmacokinetic parameters of darunavir in this study were compared with the pharmacokinetic parameters following administration of darunavir/ritonavir 600/100 mg b.i.d.

§ Ratio based on between-study comparison.

¶ q.o.d. = every other day

^ AUC is AUC_(0-last); N=10 for C_{min} in the reference arm

|| N=14 for C_{max}

€ The dose of simeprevir in this interaction study was 50 mg when co-administered in combination with darunavir/ritonavir compared to 150 mg once daily in the simeprevir alone treatment group

+ Maximum number of subjects

Table 15: Drug Interactions: Pharmacokinetic Parameters for <u>Co-administered Drugs</u> in the Presence of Darunavir/Ritonavir							
Co-Administered Drug	Dose/Schedule		N	PK	LS Mean Ratio (90% CI) of <u>Co-Administered Drug</u> Pharmacokinetic Parameters With/Without Darunavir No effect =1.00		
	Co-Administered Drug	Darunavir/ritonavir			C_{max}	AUC	C_{min}
Co-Administration With Other HIV Protease Inhibitors							
Atazanavir	300 mg q.d.* /100 mg ritonavir q.d. when administered alone 300 mg q.d. when administered with darunavir/ritonavir	400/100 mg b.i.d. †	13	↔	0.89 (0.78-1.01)	1.08 (0.94-1.24)	1.52 (0.99-2.34)
Indinavir	800 mg b.i.d. /100 mg ritonavir b.i.d. when administered alone 800 mg b.i.d. when administered with darunavir/ritonavir	400/100 mg b.i.d.	9	↑	1.08 (0.95-1.22)	1.23 (1.06-1.42)	2.25 (1.63-3.10)
Lopinavir/ Ritonavir	400/100 mg b.i.d. ‡ 533/133.3 mg b.i.d. ‡	1200/100 mg b.i.d.	14	↔	0.98 (0.78-1.22)	1.09 (0.86-1.37)	1.23 (0.90-1.69)
		1200 mg b.i.d.	15	↔	1.11 (0.96-1.30)	1.09 (0.96-1.24)	1.13 (0.90-1.42)
Saquinavir hard gel capsule	1000 mg b.i.d. /100 mg ritonavir b.i.d. when administered alone 1000 mg b.i.d. when administered with darunavir/ritonavir	400/100 mg b.i.d.	12	↔	0.94 (0.78-1.13)	0.94 (0.76-1.17)	0.82 (0.52-1.30)
Co-Administration With Other HIV Antiretrovirals							
Didanosine	400 mg q.d.	600/100 mg b.i.d.	17	↔	0.84 (0.59-1.20)	0.91 (0.75-1.10)	-
Dolutegravir	30 mg q.d.	600/100 mg b.i.d.	15	↓	0.89 (0.83-	0.78 (0.72-	0.62 ^Ω (0.56-

					0.97)	0.85)	0.69)
Dolutegravir	50 mg q.d.	600/100 mg b.i.d. with 200 mg b.i.d. etravirine	9	↓	0.88 (0.78-1.00)	0.75 (0.69-0.81)	0.63 ^Ω (0.52-0.76)
Efavirenz	600 mg q.d.	300/100 mg b.i.d.	12	↑	1.15 (0.97-1.35)	1.21 (1.08-1.36)	1.17 (1.01-1.36)
Etravirine	100 mg b.i.d.	600/100 mg b.i.d.	14	↓	0.68 (0.57-0.82)	0.63 (0.54-0.73)	0.51 (0.44-0.61)
Nevirapine	200 mg b.i.d.	400/100 mg b.i.d.	8	↑	1.18 (1.02-1.37)	1.27 (1.12-1.44)	1.47 (1.20-1.82)
Rilpivirine	150 mg q.d.	800/100 mg q.d.	14	↑	1.79 (1.56-2.06)	2.30 (1.98-2.67)	2.78 (2.39-3.24)
Tenofovir Disoproxil Fumarate	300 mg q.d.	300/100 mg b.i.d.	12	↑	1.24 (1.08-1.42)	1.22 (1.10-1.35)	1.37 (1.19-1.57)
Maraviroc	150 mg b.i.d.	600/100 mg b.i.d.	12	↑	2.29 (1.46-3.59)	4.05 (2.94-5.59)	8.00 (6.35-10.1)
Maraviroc	150 mg b.i.d.	600/100 mg b.i.d. with 200 mg b.i.d. etravirine	10	↑	1.77 (1.20-2.60)	3.10 (2.57-3.74)	5.27 (4.51-6.15)
Co-Administration With HCV NS3-4A Protease Inhibitors							
Boceprevir	800 mg three times daily	600/100 mg b.i.d.	12 [^]	↓	0.75 (0.67-0.85)	0.68 (0.65-0.72)	0.65 (0.56-0.76)
Telaprevir	750 mg every 8 hours	600/100 mg b.i.d.	11	↓	0.64 (0.61-0.67)	0.65 (0.61-0.69)	0.68 (0.63-0.74)
Simeprevir	50 mg q.d. [€]	800/100 mg q.d.	25 ⁺	↑	1.79 (1.55-2.06)	2.59 (2.15-3.11)	4.58 (3.54-5.92)
Co-Administration With Other Drugs							
Atorvastatin	40 mg q.d. when administered alone 10 mg q.d. when administered with darunavir/ritonavir	300/100 mg b.i.d.	15	↑	0.56 (0.48-0.67)	0.85 (0.76-0.97)	1.81 (1.37-2.40)
Artemether Dihydroartemisinin	80 mg single dose	600/100 mg b.i.d.	15	↓	0.85 (0.68-1.05)	0.91 (0.78-1.06)	-
			15	↑	1.06 (0.82-1.39)	1.12 (0.96-1.30)	-
Artemether	Artemether/	600/100 mg	15	↓	0.82	0.84	0.97

Dihydroartemisinin	lumefantrine 80/480 mg (6 doses at 0, 8, 24, 36, 48, and 60 hours)	b.i.d.			(0.61- 1.11)	(0.69- 1.02)	(0.90- 1.05)
Lumefantrine			15	↓	0.82 (0.66- 1.01)	0.82 (0.74- 0.91)	1.00 (0.82- 1.22)
			15	↑	1.65 (1.49- 1.83)	2.75 (2.46- 3.08)	2.26 (1.92- 2.67)
Buprenorphine/ Naloxone	8/2 mg to 16/4 mg q.d.	600/100 mg b.i.d.	17	↔	0.92 [§] (0.79- 1.08)	0.89 [§] (0.78- 1.02)	0.98 [§] (0.82- 1.16)
Norbuprenorphine			17	↑	1.36 (1.06- 1.74)	1.46 (1.15- 1.85)	1.71 (1.29- 2.27)
Carbamazepine	200 mg b.i.d.	600/100 mg b.i.d.	16	↑	1.43 (1.34- 1.53)	1.45 (1.35- 1.57)	1.54 (1.41- 1.68)
Carbamazepine epoxide			16	↓	0.46 (0.43- 0.49)	0.46 (0.44- 0.49)	0.48 (0.45- 0.51)
Clarithromycin	500 mg b.i.d.	400/100 mg b.i.d.	17	↑	1.26 (1.03- 1.54)	1.57 (1.35- 1.84)	2.74 (2.30- 3.26)
Dextromethorphan	30 mg	600/100 mg b.i.d.	12	↑	2.27 (1.59- 3.26)	2.70 (1.80- 4.05)	-
Dextrophan				↓	0.87 (0.77- 0.98)	0.96 (0.90- 1.03)	-
Digoxin	0.4 mg	600/100 mg b.i.d.	8	↑	1.15 (0.89- 1.48)	1.36 (0.81- 2.27)	-
Ethinyl estradiol (EE)	Ortho-Novum 1/35 (35 µg EE / 1 mg NE)	600/100 mg b.i.d.	11	↓	0.68 (0.61- 0.74)	0.56 (0.50- 0.63)	0.38 (0.27- 0.54)
Norethindrone (NE)			11	↓	0.90 (0.83- 0.97)	0.86 (0.75- 0.98)	0.70 (0.51- 0.97)
Ketoconazole	200 mg b.i.d.	400/100 mg b.i.d.	15	↑	2.11 (1.81- 2.44)	3.12 (2.65- 3.68)	9.68 (6.44- 14.55)
R-Methadone	55-150 mg q.d.	600/100 mg b.i.d.	16	↓	0.76 (0.71- 0.81)	0.84 (0.78- 0.91)	0.85 (0.77- 0.94)
Omeprazole	40 mg single dose	600/100 mg b.i.d.	12	↓	0.66 (0.48- 0.90)	0.58 (0.50- 0.66)	-

5-hydroxy omeprazole				↓	0.93 (0.71-1.21)	0.84 (0.77-0.92)	-
Paroxetine	20 mg q.d.	400/100 mg b.i.d.	16	↓	0.64 (0.59-0.71)	0.61 (0.56-0.66)	0.63 (0.55-0.73)
Pitavastatin	4 mg q.d.	800/100 mg q.d.	27	↓	0.96 (0.84-1.09)	0.74 (0.69-0.80)	NA
Pravastatin	40 mg single dose	600/100 mg b.i.d.	14	↑	1.63 (0.95-2.82)	1.81 (1.23-2.66)	-
Rifabutin	150 mg q.o.d. [†] when administered with PREZISTA/ ritonavir	600/100 mg b.i.d. [#]	11	↑	0.72 (0.55-0.93)	0.93 (0.80-1.09)	1.64 (1.48-1.81)
25-O-desacetyl-rifabutin	300 mg q.d. when administered alone		11	↑	4.77 (4.04-5.63)	9.81 (8.09-11.9)	27.1 (22.2-33.2)
Sertraline	50 mg q.d.	400/100 mg b.i.d.	13	↓	0.56 (0.49-0.63)	0.51 (0.46-0.58)	0.51 (0.45-0.57)
Sildenafil	100 mg (single dose) administered alone 25 mg (single dose) when administered with darunavir/ ritonavir	400/100 mg b.i.d.	16	↑	0.62 (0.55-0.70)	0.97 (0.86-1.09)	-
S-warfarin	10 mg single dose	600/100 mg b.i.d.	12	↓	0.92 (0.86-0.97)	0.79 (0.73-0.85)	-
7-OH-S-warfarin			12	↑	1.42 (1.24-1.63)	1.23 (0.97-1.57)	-

N = number of subjects with data; - = no information available

* q.d. = once daily

† b.i.d. = twice daily

‡ The pharmacokinetic parameters of lopinavir in this study were compared with the pharmacokinetic parameters following administration of lopinavir/ritonavir 400/100 mg b.i.d.

§ ratio is for buprenorphine; mean C_{max} and AUC₂₄ for naloxone were comparable when buprenorphine/naloxone was administered with or without PREZISTA/ritonavir

¶ q.o.d. = every other day

In comparison to rifabutin 300 mg q.d.

^ N=11 for the test arm

|| N=14 for C_{max}

Ω Noted as C_r or C₂₄ in the dolutegravir U.S. prescribing information

+ Maximum number of subjects

A cocktail study was conducted in 12 healthy volunteers to evaluate the effect of steady state pharmacokinetics of darunavir/ritonavir on the activity of CYP2D6 (using dextromethorphan as probe substrate), CYP2C9 (using warfarin as probe substrate), and CYP2C19 (using omeprazole as probe substrate). The pharmacokinetic results are shown in Table 15.

€ The dose of simeprevir in this interaction study was 50 mg when co-administered in combination with darunavir/ritonavir compared to 150 mg once daily in the simeprevir alone treatment group.

12.4 Microbiology

Mechanism of Action

Darunavir is an inhibitor of the HIV-1 protease. It selectively inhibits the cleavage of HIV-1 encoded Gag-Pol polyproteins in infected cells, thereby preventing the formation of mature virus particles.

Antiviral Activity

Darunavir exhibits activity against laboratory strains and clinical isolates of HIV-1 and laboratory strains of HIV-2 in acutely infected T-cell lines, human peripheral blood mononuclear cells and human monocytes/macrophages with median EC₅₀ values ranging from 1.2 to 8.5 nM (0.7 to 5.0 ng/mL). Darunavir demonstrates antiviral activity in cell culture against a broad panel of HIV-1 group M (A, B, C, D, E, F, G), and group O primary isolates with EC₅₀ values ranging from less than 0.1 to 4.3 nM. The EC₅₀ value of darunavir increases by a median factor of 5.4 in the presence of human serum. Darunavir did not show antagonism when studied in combination with the PIs amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir, saquinavir, or tipranavir, the N(t)RTIs abacavir, didanosine, emtricitabine, lamivudine, stavudine, tenofovir, zalcitabine, or zidovudine, the NNRTIs delavirdine, rilpivirine, efavirenz, etravirine, or nevirapine, and the fusion inhibitor enfuvirtide.

Resistance

Cell Culture: HIV-1 isolates with a decreased susceptibility to darunavir have been selected in cell culture and obtained from subjects treated with darunavir/ritonavir. Darunavir-resistant virus derived in cell culture from wild-type HIV-1 had 21- to 88-fold decreased susceptibility to darunavir and developed 2 to 4 of the following amino acid substitutions S37D, R41E/T, K55Q, H69Q, K70E, T74S, V77I, or I85V in the protease. Selection in cell culture of darunavir resistant HIV-1 from nine HIV-1 strains harboring multiple PI resistance-associated mutations resulted in the overall emergence of 22 mutations in the protease gene, coding for amino acid substitutions L10F, V11I, I13V, I15V, G16E, L23I, V32I, L33F, S37N, M46I, I47V, I50V, F53L, L63P, A71V, G73S, L76V, V82I, I84V, T91A/S, and Q92R, of which L10F, V32I, L33F, S37N, M46I, I47V, I50V, L63P, A71V, and I84V were the most prevalent. These darunavir-resistant viruses had at least eight protease substitutions and exhibited 50- to 641-fold decreases in darunavir susceptibility with final EC₅₀ values ranging from 125 nM to 3461 nM.

Clinical studies of PREZISTA/ritonavir in treatment-experienced subjects: In a pooled analysis of the 600/100 mg PREZISTA/ritonavir twice daily arms of Studies TMC114-C213, TMC114-C202, TMC114-C215, and the control arms of etravirine studies TMC125-C206 and TMC125-C216, the amino acid substitutions V32I and I54L or M developed most frequently on PREZISTA/ritonavir in 41% and 25%, respectively, of the treatment-experienced subjects who experienced virologic failure, either by rebound or by never being suppressed (less than 50 copies/mL). Other substitutions that developed frequently in PREZISTA/ritonavir virologic failure isolates occurred at amino acid positions V11I, I15V, L33F, I47V, I50V, and L89V. These amino acid substitutions were associated with decreased susceptibility to darunavir; 90% of the virologic failure isolates had a greater than 7-fold decrease in susceptibility to darunavir at failure. The median darunavir phenotype (fold change from reference) of the virologic failure isolates was 4.3-fold at baseline and 85-fold at failure. Amino acid substitutions were also observed in the protease cleavage sites in the Gag polyprotein of some PREZISTA/ritonavir virologic failure isolates. In Study TMC114-C212 of treatment-experienced pediatric subjects, the amino acid substitutions V32I, I54L and L89M developed most frequently in virologic failures on PREZISTA/ritonavir.

In the 96-week as-treated analysis of the Phase 3 Study TMC114-C214, the percent of virologic failures (never suppressed, rebounders and discontinued before achieving suppression) was 21% (62/298) in the group of subjects receiving PREZISTA/ritonavir 600/100 mg twice daily compared to 32% (96/297) of subjects receiving lopinavir/ritonavir 400/100 mg twice daily. Examination of subjects who failed on PREZISTA/ritonavir 600/100 mg twice daily and had post-baseline genotypes and phenotypes showed that 7 subjects (7/43; 16%) developed PI substitutions on darunavir/ritonavir treatment resulting in decreased susceptibility to darunavir. Six of the 7 had baseline PI resistance-associated substitutions and baseline darunavir phenotypes greater than 7. The most common emerging PI substitutions in these virologic failures were V32I, L33F, M46I or L, I47V, I54L, T74P and L76V. These amino acid substitutions were associated with 59- to 839-fold decreased susceptibility to darunavir at failure. Examination of individual subjects who failed in the comparator arm on lopinavir/ritonavir and had post-baseline genotypes and phenotypes showed that 31 subjects (31/75; 41%) developed substitutions on lopinavir treatment resulting in decreased susceptibility to lopinavir (greater than 10-fold) and the most common substitutions emerging on treatment were L10I or F, M46I or L, I47V or A, I54V and L76V. Of the 31 lopinavir/ritonavir virologic failure subjects, 14 had reduced susceptibility (greater than 10-fold) to lopinavir at baseline.

In the 48-week analysis of the Phase 3 Study TMC114-C229, the number of virologic failures (including those who discontinued before suppression after Week 4) was 26% (75/294) in the group of subjects receiving PREZISTA/ritonavir 800/100 mg once daily compared to 19% (56/296) of subjects receiving PREZISTA/ritonavir 600/100 mg twice daily. Examination of isolates from subjects who failed on PREZISTA/ritonavir 800/100

mg once daily and had post-baseline genotypes showed that 8 subjects (8/60; 13%) had isolates that developed IAS-USA defined PI resistance-associated substitutions compared to 5 subjects (5/39; 13%) on PREZISTA/ritonavir 600/100 mg twice daily. Isolates from 2 subjects developed PI resistance associated substitutions associated with decreased susceptibility to darunavir; 1 subject isolate in the PREZISTA/ritonavir 800/100 mg once daily arm, developed substitutions V32I, M46I, L76V and I84V associated with a 24-fold decreased susceptibility to darunavir, and 1 subject isolate in the PREZISTA/ritonavir 600/100 mg twice daily arm developed substitutions L33F and I50V associated with a 40-fold decreased susceptibility to darunavir. In the PREZISTA/ritonavir 800/100 mg once daily and PREZISTA/ritonavir 600/100 mg twice daily groups, isolates from 7 (7/60, 12%) and 4 (4/42, 10%) virologic failures, respectively, developed decreased susceptibility to an NRTI included in the treatment regimen.

Clinical studies of PREZISTA/ritonavir in treatment-naïve subjects: In the 192-week as-treated analysis censoring those who discontinued before Week 4 of the Phase 3 Study TMC114-C211, the percentage of virologic failures (never suppressed, rebounders and discontinued before achieving suppression) was 22% (64/288) in the group of subjects receiving PREZISTA/ritonavir 800/100 mg once daily compared to 29% (76/263) of subjects receiving lopinavir/ritonavir 800/200 mg per day. In the PREZISTA/ritonavir arm, emergent PI resistance-associated substitutions were identified in 11 of the virologic failures with post-baseline genotypic data (n=43). However, none of the darunavir virologic failures had a decrease in darunavir susceptibility (greater than 7-fold change) at failure. In the comparator lopinavir/ritonavir arm, emergent PI resistance-associated substitutions were identified in 17 of the virologic failures with post-baseline genotypic data (n=53), but none of the lopinavir/ritonavir virologic failures had decreased susceptibility to lopinavir (greater than 10-fold change) at failure. The reverse transcriptase M184V substitution and/or resistance to emtricitabine, which was included in the fixed background regimen, was identified in 4 virologic failures from the PREZISTA/ritonavir arm and 7 virologic failures in the lopinavir/ritonavir arm.

Cross-resistance

Cross-resistance among PIs has been observed. Darunavir has a less than 10-fold decreased susceptibility in cell culture against 90% of 3309 clinical isolates resistant to amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir, saquinavir and/or tipranavir showing that viruses resistant to these PIs remain susceptible to darunavir.

Darunavir-resistant viruses were not susceptible to amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir or saquinavir in cell culture. However, six of nine darunavir-resistant viruses selected in cell culture from PI-resistant viruses showed a fold change in EC50 values less than 3 for tipranavir, indicative of limited cross-resistance between darunavir and tipranavir. In Studies TMC114-C213, TMC114-C202, and TMC114-C215, 34% (64/187) of subjects in the darunavir/ritonavir arm whose baseline isolates had decreased susceptibility to tipranavir (tipranavir fold change greater than 3)

achieved less than 50 copies/mL serum HIV-1 RNA levels at Week 96. Of the viruses isolated from subjects experiencing virologic failure on PREZISTA/ritonavir 600/100 mg twice daily (greater than 7 fold change), 41% were still susceptible to tipranavir and 10% were susceptible to saquinavir while less than 2% were susceptible to the other protease inhibitors (amprenavir, atazanavir, indinavir, lopinavir or nelfinavir).

In Study TMC114-C214, the 7 darunavir/ritonavir virologic failures with reduced susceptibility to darunavir at failure were also resistant to the approved PIs (fos)amprenavir, atazanavir, lopinavir, indinavir, and nelfinavir at failure. Six of these 7 were resistant to saquinavir and 5 were resistant to tipranavir. Four of these virologic failures were already PI-resistant at baseline.

Cross-resistance between darunavir and nucleoside/nucleotide reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, fusion inhibitors, CCR5 co-receptor antagonists, or integrase inhibitors is unlikely because the viral targets are different.

Baseline Genotype/Phenotype and Virologic Outcome Analyses

Genotypic and/or phenotypic analysis of baseline virus may aid in determining darunavir susceptibility before initiation of PREZISTA/ritonavir 600/100 mg twice daily therapy. The effect of baseline genotype and phenotype on virologic response at 96 weeks was analyzed in as-treated analyses using pooled data from the Phase 2b studies (Studies TMC114-C213, TMC114-C202, and TMC114-C215) (n=439). The findings were confirmed with additional genotypic and phenotypic data from the control arms of etravirine Studies TMC125-C206 and TMC125-C216 at Week 24 (n=591).

Diminished virologic responses were observed in subjects with 5 or more baseline IAS-defined primary protease inhibitor resistance-associated substitutions (D30N, V32I, L33F, M46I/L, I47A/V, G48V, I50L/V, I54L/M, L76V, V82A/F/L/S/T, I84V, N88S, L90M) (see Table 16).

Table 16: Response to PREZISTA/ritonavir 600/100 mg Twice Daily by Baseline Number of IAS-Defined Primary PI Resistance-Associated Substitutions: As-treated Analysis of Studies TMC114-C213, TMC114-C202, and TMC114-C215			
	Studies TMC114-C213, TMC114-C202, TMC114-C215 <50 copies/mL at Week 96 N=439		
# IAS-Defined Primary PI Substitutions	Overall	De Novo ENF	Re-Used/No ENF
All	44% (192/439)	54% (61/112)	40% (131/327)
0 – 4	50% (162/322)	58% (49/85)	48% (113/237)
5	22% (16/74)	47% (9/19)	13% (7/55)
≥ 6	9% (3/32)	17% (1/6)	8% (2/26)

IAS Primary PI Substitutions (2008): D30N, V32I, L33F, M46I/L, I47A/V, G48V, I50L/V, I54L/M, L76V, V82A/F/L/S/T, I84V, N88S, L90M

The presence at baseline of two or more of the substitutions V11I, V32I, L33F, I47V, I50V, I54L or M, T74P, L76V, I84V or L89V was associated with a decreased virologic response to PREZISTA/ritonavir. In subjects not taking enfuvirtide *de novo*, the proportion of subjects achieving viral load less than 50 plasma HIV-1 RNA copies/mL at 96 weeks was 59%, 29%, and 12% when the baseline genotype had 0-1, 2 and greater than or equal to 3 of these substitutions, respectively.

Baseline darunavir phenotype (shift in susceptibility relative to reference) was shown to be a predictive factor of virologic outcome. Response rates assessed by baseline darunavir phenotype are shown in Table 17. These baseline phenotype groups are based on the select patient populations in the Studies TMC114-C213, TMC114-C202, and TMC114-C215, and are not meant to represent definitive clinical susceptibility breakpoints for PREZISTA/ritonavir. The data are provided to give clinicians information on the likelihood of virologic success based on pre-treatment susceptibility to darunavir.

Table 17: Response (HIV-1 RNA <50 copies/mL at Week 96) to PREZISTA/ritonavir 600/100 mg Twice Daily by Baseline Darunavir Phenotype and by Use of Enfuvirtide (ENF): As-treated Analysis of Studies TMC114-C213, TMC114-C202, and TMC114-C215			
Proportion of Subjects with <50 copies/mL at Week 96			
N=417			
Baseline DRV Phenotype	All	De Novo ENF	Re-Used/No ENF
Overall	175/417 (42%)	61/112 (54%)	131/327 (40%)
0 – 7	148/270 (55%)	44/65 (68%)	104/205 (51%)
> 7 – 20	16/53 (30%)	7/17 (41%)	9/36 (25%)
> 20	11/94 (12%)	6/23 (26%)	5/71 (7%)

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis and Mutagenesis

Darunavir was evaluated for carcinogenic potential by oral gavage administration to mice and rats up to 104 weeks. Daily doses of 150, 450 and 1000 mg/kg were administered to mice and doses of 50, 150 and 500 mg/kg was administered to rats. A dose-related increase in the incidence of hepatocellular adenomas and carcinomas were observed in males and females of both species as well as an increase in thyroid follicular cell adenomas in male rats. The observed hepatocellular findings in rodents are considered to be of limited relevance to humans. Repeated administration of darunavir to rats caused hepatic microsomal enzyme induction and increased thyroid hormone elimination, which predispose rats, but not humans, to thyroid neoplasms. At the highest tested doses, the systemic exposures to darunavir (based on AUC) were between 0.4- and 0.7-fold (mice) and 0.7- and 1-fold (rats), relative to those observed in humans at the recommended therapeutic doses (600/100 mg twice daily or 800/100 mg once daily).

Darunavir was not mutagenic or genotoxic in a battery of *in vitro* and *in vivo* assays including bacterial reserve mutation (Ames), chromosomal aberration in human lymphocytes and *in vivo* micronucleus test in mice.

Impairment of Fertility

No effects on fertility or early embryonic development were observed with darunavir in rats and darunavir has shown no teratogenic potential in mice or rats (in the presence or absence of ritonavir), and rabbits.

13.2 Animal Toxicology and/or Pharmacology

In juvenile rats single doses of darunavir (20 mg/kg to 160 mg/kg at ages 5-11 days) or multiple doses of darunavir (40 mg/kg to 1000 mg/kg at age 12 days) caused mortality. The mortalities were associated with convulsions in some of the animals. Within this age range exposures in plasma, liver and brain were dose and age dependent and were considerably greater than those observed in adult rats. These findings were attributed to the ontogeny of the CYP450 liver enzymes involved in the metabolism of darunavir and the immaturity of the blood-brain barrier. No treatment-related mortalities were noted in juvenile rats after a single dose of darunavir at 1000 mg/kg on day 26 of age or after repeat dosing at 500 mg/kg from day 23 to 50 of age. The exposures and toxicity profile in the older animals (day 23 or day 26) were comparable to those observed in adult rats. Due to uncertainties regarding the rate of development of the human blood-brain barrier and liver enzymes, do not administer PREZISTA/ritonavir in pediatric patients below 3 years of age.

14 CLINICAL STUDIES

14.1 Description of Adult Clinical Studies

The evidence of efficacy of PREZISTA/ritonavir is based on the analyses of 192-week data from a randomized, controlled open-label Phase 3 trial in treatment-naïve (TMC114-C211) HIV-1-infected adult subjects and 96-week data from a randomized, controlled, open-label Phase 3 trial in antiretroviral treatment-experienced (TMC114-C214) HIV-1-infected adult subjects. In addition, 96-week data are included from 2 randomized, controlled Phase 2b trials, TMC114-C213 and TMC114-C202, in antiretroviral treatment-experienced HIV-1-infected adult subjects.

14.2 Treatment-Naïve Adult Subjects

Study TMC114-C211

Study TMC114-C211 is a randomized, controlled, open-label Phase 3 trial comparing PREZISTA/ritonavir 800/100 mg once daily versus lopinavir/ritonavir 800/200 mg per day (given as a twice daily or as a once daily regimen) in antiretroviral treatment-naïve HIV-1-infected adult subjects. Both arms used a fixed background regimen consisting of tenofovir disoproxil fumarate 300 mg once daily (TDF) and emtricitabine 200 mg once daily (FTC).

HIV-1-infected subjects who were eligible for this trial had plasma HIV-1 RNA greater than or equal to 5000 copies/mL. Randomization was stratified by screening plasma viral load (HIV-1 RNA less than 100,000 copies/mL or greater than or equal to 100,000 copies/mL) and screening CD4+ cell count (less than 200 cells/mm³ or greater than or equal to 200 cells/mm³). Virologic response was defined as a confirmed plasma HIV-1 RNA viral load less than 50 copies/mL. Analyses included 689 subjects in Study TMC114-C211 who had completed 192 weeks of treatment or discontinued earlier.

Demographics and baseline characteristics were balanced between the PREZISTA/ritonavir arm and the lopinavir/ritonavir arm (see Table 18). Table 18 compares the demographic and baseline characteristics between subjects in the PREZISTA/ritonavir 800/100 mg once daily arm and subjects in the lopinavir/ritonavir 800/200 mg per day arm in Study TMC114-C211.

Table 18: Demographic and Baseline Characteristics of Subjects in Study TMC114-C211 □		
	Randomized Study TMC114-C211	
	PREZISTA/ritonavir 800/100 mg once daily + TDF/FTC N=343	lopinavir/ritonavir 800/200 mg per day + TDF/FTC N=346
Demographic Characteristics		
Median Age (years) (range, years)	34 (18-70)	33 (19-68)
Sex		
Male	70%	70%
Female	30%	30%
Race		
White	40%	45%
Black	23%	21%
Hispanic	23%	22%
Asian	13%	11%
Baseline Characteristics		
Mean Baseline Plasma HIV-1 RNA (log ₁₀ copies/mL)	4.86	4.84
Median Baseline CD4+ Cell Count (cells/mm ³) (range, cells/mm ³)	228 (4-750)	218 (2-714)
Percentage of Patients with Baseline Viral Load ≥ 100,000 copies/mL	34%	35%
Percentage of Patients with Baseline CD4+ Cell Count <200 cells/mm ³	41%	43%

Week 192 outcomes for subjects on PREZISTA/ritonavir 800/100 mg once daily from Study TMC114-C211 are shown in Table 19.

Table 19: Virologic Outcome of Randomized Treatment of Study TMC114-C211 at 192 Weeks □		
	PREZISTA/ritonavir 800/100 mg once daily + TDF/FTC N=343	lopinavir/ritonavir 800/200 mg per day + TDF/FTC N=346
Virologic success HIV-1 RNA <50 copies/mL	70%*	61%

Virologic failure [†]	12%	15%
No virologic data at Week 192 window [‡]		
<u>Reasons</u>		
Discontinued study due to adverse event or death [§]	5%	13%
Discontinued study for other reasons [¶]	13%	12%
Missing data during window [‡] but on study	<1%	0%

N = total number of subjects with data

* 95% CI: 1.9; 16.1

[†] Includes patients who discontinued prior to Week 192 for lack or loss of efficacy and patients who are ≥ 50 copies in the 192-week window and patients who had a change in their background regimen that was not permitted by the protocol

[‡] Window 186-198 Weeks

[§] Includes patients who discontinued due to adverse event or death at any time point from Day 1 through the time window if this resulted in no virologic data on treatment during the specified window

[¶] Other includes: withdrew consent, loss to follow-up, etc., if the viral load at the time of discontinuation was <50 copies/mL

In Study TMC114-C211 at 192 weeks of treatment, the median increase from baseline in CD4+ cell counts was 258 cells/mm³ in the PREZISTA/ritonavir 800/100 mg once daily arm and 263 cells/mm³ in the lopinavir/ritonavir 800/200 mg per day arm. Of the PREZISTA/ritonavir subjects with a confirmed virologic response of <50 copies/mL at Week 48, 81% remained undetectable at Week 192 versus 68% with lopinavir/ritonavir. In the 192 week analysis, statistical superiority of the PREZISTA/ritonavir regimen over the lopinavir/ritonavir regimen was demonstrated for both ITT and OP populations.

14.3 Treatment-Experienced Adult Subjects

Study TMC114-C229

Study TMC114-C229 is a randomized, open-label trial comparing PREZISTA/ritonavir 800/100 mg once daily to PREZISTA/ritonavir 600/100 mg twice daily in treatment-experienced HIV-1-infected patients with screening genotype resistance test showing no darunavir resistance associated substitutions (i.e. V11I, V32I, L33F, I47V, I50V, I54L, I54M, T74P, L76V, I84V, L89V) and a screening viral load of greater than 1,000 HIV-1 RNA copies/mL. Both arms used an optimized background regimen consisting of greater than or equal to 2 NRTIs selected by the investigator.

HIV-1-infected subjects who were eligible for this trial were on a highly active antiretroviral therapy regimen (HAART) for at least 12 weeks. Virologic response was defined as a confirmed plasma HIV-1 RNA viral load less than 50 copies/mL. Analyses included 590 subjects who had completed 48 weeks of treatment or discontinued earlier.

Table 20 compares the demographic and baseline characteristics between subjects in the PREZISTA/ritonavir 800/100 mg once daily arm and subjects in the PREZISTA/ritonavir 600/100 mg twice daily arm in Study TMC114-C229. No imbalances between the 2 arms were noted.

Table 20: Demographic and Baseline Characteristics of Subjects in Study TMC114-C229		
	Randomized Study TMC114-C229	
	PREZISTA/ritonavir 800/100 mg once daily + OBR N=294	PREZISTA/ritonavir 600/100 mg twice daily + OBR N=296
Demographic Characteristics		
Median Age (years) (range, years)	40 (18-70)	40 (18-77)
Sex		
Male	61%	67%
Female	39%	33%
Race		
White	35%	37%
Black	28%	24%
Hispanic	16%	20%
Asian	16%	14%
Baseline Characteristics		
Mean Baseline Plasma HIV-1 RNA (log ₁₀ copies/mL)	4.19	4.13
Median Baseline CD4+ Cell Count (cells/mm ³) (range, cells/mm ³)	219 (24-1306)	236 (44-864)
Percentage of Patients with Baseline Viral Load ≥ 100,000 copies/mL	13%	11%
Percentage of Patients with Baseline CD4+ Cell Count <200 cells/mm ³	43%	39%
Median Darunavir Fold Change (range)*	0.50 (0.1-1.8)	0.50 (0.1-1.9)
Median Number of Resistance-Associated†:		
PI mutations	3	4
NNRTI mutations	2	1
NRTI mutations	1	1
Percentage of Subjects Susceptible to All Available PIs at Baseline	88%	86%
Percentage of Subjects with Number of Baseline Primary Protease Inhibitor Mutations‡:		
0	84%	84%
1	8%	9%
2	5%	4%
≥ 3	3%	2%
Median Number of ARVs Previously Used‡:		
NRTIs	3	3
NNRTIs	1	1
PIs (excluding low-dose ritonavir)	1	1

* Based on phenotype (Antivirogram®)

† Johnson VA, Brun-Vézinet F, Clotet B, et al. Update of the drug resistance mutations in HIV-1: December 2008. Top HIV Med 2008; 16(5): 138-145

‡ Only counting ARVs, excluding low-dose ritonavir

Week 48 outcomes for subjects on PREZISTA/ritonavir 800/100 mg once daily from Study TMC114-C229 are shown in Table 21.

Table 21: Virologic Outcome of Randomized Treatment of Study TMC114-C229 at 48 Weeks		
	Randomized Study TMC114-C229	
	PREZISTA/ritonavir 800/100 mg once daily + OBR N=294	PREZISTA/ritonavir 600/100 mg twice daily + OBR N=296
Virologic success HIV-1 RNA <50 copies/mL	69%	69%
Virologic failure*	26%	23%
No virologic data at Week 48 window†		
Reasons		
Discontinued study due to adverse event or death‡	3%	4%
Discontinued study for other reasons§	2%	3%
Missing data during window† but on study	0%	<1%

N = total number of subjects with data

* Includes patients who discontinued prior to Week 48 for lack or loss of efficacy, patients who are ≥ 50 copies in the 48-week window, patients who had a change in their background regimen that was not permitted in the protocol (provided the switch occurred before the earliest onset of an AE leading to permanent stop of study medication) and patients who discontinued for reasons other than AEs/death and lack or loss of efficacy (provided their last available viral load was detectable (HIV RNA ≥ 50 copies/mL).

† Window 42-54 Weeks

‡ Patients who discontinued due to adverse event or death at any time point from Day 1 through the time window if this resulted in no virologic data on treatment during the specified window.

§ Other includes: withdrew consent, loss to follow-up, etc., if the viral load at the time of discontinuation was < 50 copies/mL.

The mean increase from baseline in CD4+ cell counts was comparable for both treatment arms (108 cells/mm³ and 112 cells/mm³ in the PREZISTA/ritonavir 800/100 mg once daily arm and the PREZISTA/ritonavir 600/100 mg twice daily arm, respectively).

Study TMC114-C214

Study TMC114-C214 is a randomized, controlled, open-label Phase 3 trial comparing PREZISTA/ritonavir 600/100 mg twice daily versus lopinavir/ritonavir 400/100 mg twice daily in antiretroviral treatment-experienced, lopinavir/ritonavir-naïve HIV-1-infected adult subjects. Both arms used an optimized background regimen (OBR) consisting of at least 2 antiretrovirals (NRTIs with or without NNRTIs).

HIV-1-infected subjects who were eligible for this trial had plasma HIV-1 RNA greater than 1000 copies/mL and were on a highly active antiretroviral therapy regimen (HAART) for at least 12 weeks. Virologic response was defined as a confirmed plasma HIV-1 RNA viral load less than 400 copies/mL. Analyses included 595 subjects in Study TMC114-C214 who had completed 96 weeks of treatment or discontinued earlier.

Demographics and baseline characteristics were balanced between the PREZISTA/ritonavir arm and the lopinavir/ritonavir arm (see Table 22). Table 22 compares the demographic and baseline characteristics between subjects in the PREZISTA/ritonavir 600/100 mg twice daily arm and subjects in the lopinavir/ritonavir 400/100 mg twice daily arm in Study TMC114-C214.

Table 22: Demographic and Baseline Characteristics of Subjects in Study TMC114-C214		
	Randomized Study TMC114-C214	
	PREZISTA/ritonavir 600/100 mg twice daily + OBR N=298	lopinavir/ritonavir 400/100 mg twice daily + OBR N=297
Demographic Characteristics		
Median Age (years) (range, years)	40 (18-68)	41 (22-76)
Sex		
Male	77%	81%
Female	23%	19%
Race		
White	54%	57%
Black	18%	17%
Hispanic	15%	15%
Asian	9%	9%
Baseline Characteristics		
Mean Baseline Plasma HIV-1 RNA (log ₁₀ copies/mL)	4.33	4.28
Median Baseline CD4+ Cell Count (cells/mm ³) (range, cells/mm ³)	235 (3-831)	230 (2-1096)
Percentage of Patients with Baseline Viral Load ≥ 100,000 copies/mL	19%	17%
Percentage of Patients with Baseline CD4+ Cell Count <200 cells/mm ³	40%	40%
Median Darunavir Fold Change (range)	0.60 (0.10-37.40)	0.60 (0.1-43.8)
Median Lopinavir Fold Change (range)	0.70 (0.40-74.40)	0.80 (0.30-74.50)
Median Number of Resistance-Associated*:		
PI mutations	4	4
NNRTI mutations	1	1
NRTI mutations	2	2
Percentage of Subjects with Number of Baseline Primary Protease Inhibitor Mutations*:		
≤ 1	78%	80%
2	8%	9%
≥ 3	13%	11%
Median Number of ARVs Previously Used†:		
NRTIs	4	4
NNRTIs	1	1
PIs (excluding low-dose ritonavir)	1	1
Percentage of Subjects Resistant‡ to All Available§ PIs at Baseline, excluding Darunavir	2%	3%

* Johnson VA, Brun-Vezinet F, Clotet B, et al. Update of the drug resistance mutations in HIV-1: Fall 2006. Top HIV Med 2006; 14(3): 125-130

† Only counting ARVs, excluding low-dose ritonavir

‡ Based on phenotype (Antivirogram®)

§ Commercially available PIs at the time of study enrollment

Week 96 outcomes for subjects on PREZISTA/ritonavir 600/100 mg twice daily from Study TMC114-C214 are shown in Table 23.

	PREZISTA /ritonavir 600/100 mg twice daily + OBR N=298	lopinavir/ritonavir 400/100 mg twice daily + OBR N=297
Virologic success HIV-1 RNA <50 copies/mL	58%	52%
Virologic failure*	26%	33%
No virologic data at Week 96 window†		
Reasons		
Discontinued study due to adverse event or death‡	7%	8%
Discontinued study for other reasons§	8%	7%
Missing data during window† but on study	1%	<1%

N = total number of subjects with data

* Includes patients who discontinued prior to Week 96 for lack or loss of efficacy and patients who are ≥ 50 copies in the 96-week window and patients who had a change in their OBR that was not permitted by the protocol.

† Window 90-102 Weeks

‡ Includes patients who discontinued due to adverse event or death at any time point from Day 1 through the time window if this resulted in no virologic data on treatment during the specified window.

§ Other includes: withdrew consent, loss to follow-up, etc., if the viral load at the time of discontinuation was <50 copies/mL.

In Study TMC114-C214 at 96 weeks of treatment, the median increase from baseline in CD4+ cell counts was 81 cells/mm³ in the PREZISTA/ritonavir 600/100 mg twice daily arm and 93 cells/mm³ in the lopinavir/ritonavir 400/100 mg twice daily arm.

Studies TMC114-C213 and TMC114-C202

Studies TMC114-C213 and TMC114-C202 are randomized, controlled, Phase 2b trials in adult subjects with a high level of PI resistance consisting of 2 parts: an initial partially-blinded, dose-finding part and a second long-term part in which all subjects randomized to PREZISTA/ritonavir received the recommended dose of 600/100 mg twice daily.

HIV-1-infected subjects who were eligible for these trials had plasma HIV-1 RNA greater than 1000 copies/mL, had prior treatment with PI(s), NNRTI(s) and NRTI(s), had at least one primary PI mutation (D30N, M46I/L, G48V, I50L/V, V82A/F/S/T, I84V, L90M) at screening, and were on a stable PI-containing regimen at screening for at least 8 weeks. Randomization was stratified by the number of PI mutations, screening viral load, and the use of enfuvirtide.

The virologic response rate was evaluated in subjects receiving PREZISTA/ritonavir plus an OBR versus a control group receiving an investigator-selected PI(s) regimen plus an OBR. Prior to randomization, PI(s) and OBR were selected by the investigator based on genotypic resistance testing and prior ARV history. The OBR consisted of at least 2 NRTIs with or without enfuvirtide. Selected PI(s) in the control arm included: lopinavir in 36%, (fos)amprenavir in 34%, saquinavir in 35% and atazanavir in 17%; 98% of control subjects received a ritonavir boosted PI regimen out of which 23% of control subjects used dual-boosted PIs. Approximately 47% of all subjects used enfuvirtide, and 35% of the use was in subjects who were ENF-naïve. Virologic response was defined as a decrease in plasma HIV-1 RNA viral load of at least 1 log₁₀ versus baseline.

In the pooled analysis for TMC114-C213 and TMC114-C202, demographics and baseline characteristics were balanced between the PREZISTA/ritonavir arm and the comparator PI arm (see Table 24). Table 24 compares the demographic and baseline characteristics between subjects in the PREZISTA/ritonavir 600/100 mg twice daily arm and subjects in the comparator PI arm in the pooled analysis of Studies TMC114-C213 and TMC114-C202.

Table 24: Demographic and Baseline Characteristics of Subjects in the Studies TMC114-C213 and TMC114-C202 (Pooled Analysis)		
	Randomized Studies TMC114-C213 and TMC114-C202	
	PREZISTA/ritonavir 600/100 mg twice daily + OBR N=131	Comparator PI(s) + OBR N=124
Demographic Characteristics		
Median Age (years) (range, years)	43 (27-73)	44 (25-65)
Sex		
Male	89%	88%
Female	11%	12%
Race		
White	81%	73%
Black	10%	15%
Hispanic	7%	8%
Baseline Characteristics		
Mean Baseline Plasma HIV-1 RNA (log ₁₀ copies/mL)	4.61	4.49
Median Baseline CD4+ Cell Count (cells/mm ³) (range, cells/mm ³)	153 (3-776)	163 (3-1274)
Percentage of Patients with Baseline Viral Load > 100,000 copies/mL	24%	29%
Percentage of Patients with Baseline CD4+ Cell Count <200 cells/mm ³	67%	58%
Median Darunavir Fold Change	4.3	3.3
Median Number of Resistance-Associated*:		
PI mutations	12	12
NNRTI mutations	1	1
NRTI mutations	5	5
Percentage of Subjects with Number of Baseline Primary Protease Inhibitor Mutations*:		
≤ 1	8%	9%
2	22%	21%
≥ 3	70%	70%
Median Number of ARVs Previously Used†:		
NRTIs	6	6
NNRTIs	1	1
PIs (excluding low-dose ritonavir)	5	5
Percentage of Subjects Resistant† to All Available‡ PIs at Baseline, excluding Tipranavir and Darunavir	63%	61%
Percentage of Subjects with Prior Use of Enfuvirtide	20%	17%

* Johnson VA, Brun-Vezinet F, Clotet B, et al. Update of the drug resistance mutations in HIV-1: Fall 2006. Top HIV Med 2006; 14(3): 125-130

† Based on phenotype (Antivirogram®)

‡ Commercially available PIs at the time of study enrollment

Week 96 outcomes for subjects on the recommended dose PREZISTA/ritonavir 600/100 mg twice daily from the pooled Studies TMC114-C213 and TMC114-C202 are shown in Table 25.

	Randomized Studies TMC114-C213 and TMC114-C202	
	PREZISTA/ritonavir 600/100 mg twice daily + OBR N=131	Comparator PI(s) + OBR N=124
Virologic Responders confirmed at least 1 log ₁₀ HIV-1 RNA below baseline through Week 96 (<50 copies/mL at Week 96)	57% (39%)	10% (9%)
Virologic failures	29%	80%
Lack of initial response*	8%	53%
Rebounder†	17%	19%
Never Suppressed‡	4%	8%
Death or discontinuation due to adverse events	9%	3%
Discontinuation due to other reasons	5%	7%

* Subjects who did not achieve at least a confirmed 0.5 log₁₀ HIV-1 RNA drop from baseline at Week 12

† Subjects with an initial response (confirmed 1 log₁₀ drop in viral load), but without a confirmed 1 log₁₀ drop in viral load at Week 96

‡ Subjects who never reached a confirmed 1 log₁₀ drop in viral load before Week 96

In the pooled Studies TMC114-C213 and TMC114-C202 through 48 weeks of treatment, the proportion of subjects with HIV-1 RNA less than 400 copies/mL in the arm receiving PREZISTA/ritonavir 600/100 mg twice daily compared to the comparator PI arm was 55.0% and 14.5%, respectively. In addition, the mean changes in plasma HIV-1 RNA from baseline were -1.69 log₁₀ copies/mL in the arm receiving PREZISTA/ritonavir 600/100 mg twice daily and -0.37 log₁₀ copies/mL for the comparator PI arm. The mean increase from baseline in CD4+ cell counts was higher in the arm receiving PREZISTA/ritonavir 600/100 mg twice daily (103 cells/mm³) than in the comparator PI arm (17 cells/mm³).

14.4 Pediatric Patients

The pharmacokinetic profile, safety and antiviral activity of PREZISTA/ritonavir were evaluated in 3 randomized, open-label, multicenter studies.

Study TMC114-C212

Treatment-experienced pediatric subjects between the ages of 6 and less than 18 years and weighing at least 20 kg were stratified according to their weight (greater than or equal to 20 kg to less than 30 kg, greater than or equal to 30 kg to less than 40 kg, greater than or equal to 40 kg) and received PREZISTA tablets with either ritonavir capsules or oral solution plus background therapy consisting of at least two non-protease inhibitor

antiretroviral drugs. Eighty patients were randomized and received at least one dose of PREZISTA/ritonavir. Pediatric subjects who were at risk of discontinuing therapy due to intolerance of ritonavir oral solution (e.g., taste aversion) were allowed to switch to the capsule formulation. Of the 44 pediatric subjects taking ritonavir oral solution, 23 subjects switched to the 100 mg capsule formulation and exceeded the weight-based ritonavir dose without changes in observed safety.

The 80 randomized pediatric subjects had a median age of 14 (range 6 to less than 18 years), and were 71% male, 54% Caucasian, 30% Black, 9% Hispanic and 8% other. The mean baseline plasma HIV-1 RNA was 4.64 log₁₀ copies/mL, and the median baseline CD4+ cell count was 330 cells/mm³ (range: 6 to 1505 cells/mm³). Overall, 38% of pediatric subjects had baseline plasma HIV-1 RNA ≥100,000 copies/mL. Most pediatric subjects (79%) had previous use of at least one NNRTI and 96% of pediatric subjects had previously used at least one PI.

Seventy-seven pediatric subjects (96%) completed the 24-week period. Of the patients who discontinued, one patient discontinued treatment due to an adverse event. An additional 2 patients discontinued for other reasons, one patient due to compliance and another patient due to relocation.

The proportion of pediatric subjects with HIV-1 RNA less than 400 copies/mL and less than 50 copies/mL was 64% and 50%, respectively. The mean increase in CD4+ cell count from baseline was 117 cells/mm³.

Study TMC114-C228

Treatment-experienced pediatric subjects between the ages of 3 and less than 6 years and weighing greater than or equal to 10 kg to less than 20 kg received PREZISTA oral suspension with ritonavir oral solution plus background therapy consisting of at least two active non-protease inhibitor antiretroviral drugs. Twenty-one subjects received at least one dose of PREZISTA/ritonavir.

The 21 subjects had a median age of 4.4 years (range 3 to less than 6 years), and were 48% male, 57% Black, 29% Caucasian and 14% other. The mean baseline plasma HIV-1 was 4.34 log₁₀ copies/mL, the median baseline CD4+ cell count was 927 × 10⁶ cells/L (range: 209 to 2,429 × 10⁶ cells/L) and the median baseline CD4+ percentage was 27.7% (range: 15.6% to 51.1%). Overall, 24% of subjects had a baseline plasma HIV-1 RNA greater than or equal to 100,000 copies/mL. All subjects had used greater than or equal to 2 NRTIs, 62% of subjects had used greater than or equal to 1 NNRTI and 76% had previously used at least one HIV PI.

Twenty subjects (95%) completed the 48 week period. One subject prematurely discontinued treatment due to vomiting assessed as related to ritonavir.

The proportion of subjects with HIV-1 RNA less than 50 copies/mL at Week 48 was 71%. The mean increase in CD4+ percentage from baseline was 4%. The mean change in CD4+ cell count from baseline was 187×10^6 cells/L.

Study TMC114-C230

Treatment-naïve pediatric subjects between the ages of 12 and less than 18 years and weighing at least 40 kg received the adult recommended dose of PREZISTA/ritonavir 800/100 mg once daily plus background therapy consisting of at least two non-protease inhibitor antiretroviral drugs.

The 12 randomized pediatric subjects had a median age of 14.4 years (range 12.6 to 17.3 years), and were 33.3% male, 58.3% Caucasian and 41.7% Black. The mean baseline plasma HIV-1 RNA was 4.72 log₁₀ copies/mL, and the median baseline CD4+ cell count was 282 cells/mm³ (range: 204 to 515 cells/mm³). Overall, 41.7% of pediatric subjects had baseline plasma HIV-1 RNA $\geq 100,000$ copies/mL.

All subjects completed the 48 week treatment period.

The proportion of subjects with HIV-1 RNA less than 50 copies/mL and less than 400 copies/mL was 83.3% and 91.7%, respectively. The mean increase in CD4+ cell count from baseline was 221×10^6 cells/L.

16 HOW SUPPLIED/STORAGE AND HANDLING

PREZISTA (darunavir) 100 mg/mL oral suspension is a white to off-white opaque liquid supplied in amber-colored multiple-dose bottles containing darunavir ethanolate equivalent to 100 mg of darunavir per mL packaged with a 6 mL oral dosing syringe with 0.2 mL gradations.

PREZISTA (darunavir) 75 mg tablets are supplied as white, caplet-shaped, film-coated tablets containing darunavir ethanolate equivalent to 75 mg of darunavir per tablet. Each tablet is debossed with “75” on one side and “TMC” on the other side.

PREZISTA (darunavir) 150 mg tablets are supplied as white, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 150 mg of darunavir per tablet. Each tablet is debossed with “150” on one side and “TMC” on the other side.

PREZISTA (darunavir) 600 mg tablets are supplied as orange, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 600 mg of darunavir per tablet. Each tablet is debossed with “600MG” on one side and “TMC” on the other side.

PREZISTA (darunavir) 800 mg tablets are supplied as dark red, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 800 mg of darunavir per tablet. Each tablet is debossed with “800” on one side and “T” on the other side.

PREZISTA is packaged in bottles in the following configuration:
100 mg/mL oral suspension – 200 mL bottles (NDC 59676-565-01)
75 mg tablets—bottles of 480 (NDC 59676-563-01)
150 mg tablets—bottles of 240 (NDC 59676-564-01)
600 mg tablets—bottles of 60 (NDC 59676-562-01)
800 mg tablets—bottles of 30 (NDC 59676-566-30)

Storage:

PREZISTA Oral Suspension

- Store at 25°C (77°F); with excursions permitted to 15°-30°C (59°-86°F).
- Do not refrigerate or freeze. Avoid exposure to excessive heat.
- Store in the original container.
- Shake well before each usage.

PREZISTA Tablets

- Store at 25°C (77°F); with excursions permitted to 15°-30°C (59°-86°F).

17 PATIENT COUNSELING INFORMATION

[See *FDA-approved patient labeling (Patient Information and Instruction for Use)*]

A statement to patients and healthcare providers is included on the product's bottle label:
ALERT: Find out about medicines that should NOT be taken with PREZISTA. A Patient Package Insert for PREZISTA is available for patient information.

Information About Therapy with PREZISTA

PREZISTA is not a cure for HIV-1 infection and patients may continue to experience illnesses associated with HIV-1 infection, including opportunistic infections. Patients should remain under the care of a physician when using PREZISTA.

Patients should be advised to avoid doing things that can spread HIV-1 infection to others.

- **Do not share needles or other injection equipment.**
- **Do not share personal items that can have blood or body fluids on them, like toothbrushes and razor blades.**
- **Do not have any kind of sex without protection.** Always practice safe sex by using a latex or polyurethane condom to lower the chance of sexual contact with semen, vaginal secretions, or blood.
- **Do not breastfeed.** We do not know if PREZISTA can be passed to the baby through breast milk and whether it could harm the baby. Also, mothers with HIV-1 should not breastfeed because HIV-1 can be passed to the baby in the breast milk.

Instructions for Use

Patients should be advised to take PREZISTA and ritonavir (NORVIR[®]) with food every day as prescribed. Patients should be instructed to swallow whole tablets with a drink such as water or milk. PREZISTA must always be used with ritonavir (NORVIR[®]) in combination with other antiretroviral drugs. Patients should not alter the dose of either PREZISTA or ritonavir (NORVIR[®]), discontinue ritonavir (NORVIR[®]), or discontinue therapy with PREZISTA without consulting their physician.

Patients Taking PREZISTA Once Daily

If a patient misses a dose of PREZISTA or ritonavir (NORVIR[®]) by more than 12 hours, the patient should be told to wait and then take the next dose of PREZISTA and ritonavir (NORVIR[®]) at the regularly scheduled time. If the patient misses a dose of PREZISTA or ritonavir (NORVIR[®]) by less than 12 hours, the patient should be told to take PREZISTA and ritonavir (NORVIR[®]) immediately, and then take the next dose of PREZISTA and ritonavir (NORVIR[®]) at the regularly scheduled time. If a dose of PREZISTA or ritonavir (NORVIR[®]) is skipped, the patient should not double the next dose. Inform the patient that he or she should not take more or less than the prescribed dose of PREZISTA or ritonavir (NORVIR[®]).

Patients Taking PREZISTA Twice Daily

If a patient misses a dose of PREZISTA or ritonavir (NORVIR[®]) by more than 6 hours, the patient should be told to wait and then take the next dose of PREZISTA and ritonavir (NORVIR[®]) at the regularly scheduled time. If the patient misses a dose of PREZISTA or ritonavir (NORVIR[®]) by less than 6 hours, the patient should be told to take PREZISTA and ritonavir (NORVIR[®]) immediately, and then take the next dose of PREZISTA and ritonavir (NORVIR[®]) at the regularly scheduled time. If a dose of PREZISTA or ritonavir (NORVIR[®]) is skipped, the patient should not double the next dose. Inform the patient that he or she should not take more or less than the prescribed dose of PREZISTA or ritonavir (NORVIR[®]).

Hepatotoxicity

Patients should be informed that Drug-induced hepatitis (e.g., acute hepatitis, cytolytic hepatitis) has been reported with PREZISTA co-administered with 100 mg of ritonavir. Monitor liver function before and during therapy, especially in patients with underlying chronic hepatitis, cirrhosis, or in patients who have pre-treatment elevations of transaminases. Post-marketing cases of liver injury, including some fatalities, have been reported. Patients should be advised about the signs and symptoms of liver problems. These may include jaundice of the skin or eyes, dark (tea colored) urine, pale colored stools, nausea, vomiting, loss of appetite, or pain, aching or sensitivity in the right upper quadrant of the abdomen.

Severe Skin Reactions

Patients should be informed that skin reactions ranging from mild to severe, including Stevens-Johnson Syndrome, drug rash with eosinophilia and systemic symptoms, and toxic epidermal necrolysis, have been reported with PREZISTA co-administered with 100 mg of ritonavir. Patients should be advised to discontinue PREZISTA/ritonavir immediately if signs or symptoms of severe skin reactions develop. These can include but are not limited to severe rash or rash accompanied with fever, general malaise, fatigue, muscle or joint aches, blisters, oral lesions, conjunctivitis, hepatitis and/or eosinophilia.

Drug Interactions

PREZISTA/ritonavir may interact with many drugs; therefore, patients should be advised to report to their healthcare provider the use of any other prescription or nonprescription medication or herbal products, including St. John's wort.

Patients receiving estrogen-based contraceptives should be instructed to use alternate contraceptive measures during therapy with PREZISTA/ritonavir because hormonal levels may decrease.

Fat Redistribution

Patients should be informed that redistribution or accumulation of body fat may occur in patients receiving antiretroviral therapy, including PREZISTA/ritonavir, and that the cause and long-term health effects of these conditions are not known at this time.

Product of Ireland

Manufactured by:

PREZISTA oral suspension
Janssen Pharmaceutica, N.V.
Beerse, Belgium

PREZISTA tablets
Janssen Ortho LLC,
Gurabo,
PR 00778

Manufactured for:

Janssen Therapeutics,
Division of Janssen Products, LP,
Titusville NJ 08560

NORVIR[®] is a registered trademark of its respective owner.

PREZISTA[®] is a registered trademark of Janssen Pharmaceuticals.

© Janssen Pharmaceuticals, Inc. 2006

PATIENT INFORMATION

PREZISTA (pre-ZIS-ta)

(darunavir)
Oral Suspension

PREZISTA (pre-ZIS-ta)

(darunavir)
Tablets

Read this Patient Information before you start taking PREZISTA and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

Also read the Patient Information leaflet for NORVIR[®] (ritonavir).

What is the most important information I should know about PREZISTA?

- **PREZISTA can interact with other medicines and cause serious side effects.** It is important to know the medicines that should not be taken with PREZISTA. See the section “**Who should not take PREZISTA?**”
- **PREZISTA may cause liver problems.** Some people taking PREZISTA in combination with NORVIR[®] (ritonavir) have developed liver problems which may be life-threatening. Your healthcare provider should do blood tests before and during your combination treatment with PREZISTA. If you have chronic hepatitis B or C infection, your healthcare provider should check your blood tests more often because you have an increased chance of developing liver problems.
- Tell your healthcare provider if you have any of the below signs and symptoms of liver problems.
 - Dark (tea colored) urine
 - yellowing of your skin or whites of your eyes
 - pale colored stools (bowel movements)
 - nausea

- vomiting
- pain or tenderness on your right side below your ribs
- loss of appetite

PREZISTA may cause severe or life-threatening skin reactions or rash. Sometimes these skin reactions and skin rashes can become severe and require treatment in a hospital. You should call your healthcare provider immediately if you develop a rash. However, **stop** taking PREZISTA and ritonavir combination treatment and call your healthcare provider immediately if you develop any skin changes with symptoms below:

- fever
- tiredness
- muscle or joint pain
- blisters or skin lesions
- mouth sores or ulcers
- red or inflamed eyes, like “pink eye” (conjunctivitis)

Rash occurred more often in people taking PREZISTA and raltegravir together than with either drug separately, but was generally mild.

See “**What are the possible side effects of PREZISTA?**” for more information about side effects.

What is PREZISTA?

PREZISTA is a prescription anti-HIV medicine used with ritonavir and other anti-HIV medicines to treat adults and children 3 years of age and older with human immunodeficiency virus (HIV-1) infection. PREZISTA is a type of anti-HIV medicine called a protease inhibitor. HIV is the virus that causes AIDS (Acquired Immune Deficiency Syndrome).

When used with other HIV medicines, PREZISTA may help to reduce the amount of HIV in your blood (called “viral load”). PREZISTA may also help to increase the number of white blood cells called CD4 (T) cell which help fight off other infections. Reducing the amount of HIV and increasing the CD4 (T) cell count may improve your immune system. This may reduce your risk of death or infections that can happen when your immune system is weak (opportunistic infections).

Children under 3 years of age should not take PREZISTA.

PREZISTA does not cure HIV infection or AIDS and you may continue to experience illnesses associated with HIV-1 infection, including opportunistic infections. You should remain under the care of a doctor when using PREZISTA.

Avoid doing things that can spread HIV-1 infection.

- **Do not share needles or other injection equipment.**
- **Do not share personal items that can have blood or body fluids on them, like toothbrushes and razor blades.**
- **Do not have any kind of sex without protection.** Always practice safe sex by using a latex or polyurethane condom to lower the chance of sexual contact with semen, vaginal secretions, or blood.

Ask your healthcare provider if you have any questions on how to prevent passing HIV to other people.

Who should not take PREZISTA?

Do not take PREZISTA with any of the following medicines:

- alfuzosin (Uroxatral[®])
- ergot-containing medicines: dihydroergotamine (D.H.E. 45[®], Embolex[®], Migranal[®]), ergotamine (Cafergot[®], Ergomar[®])
methylergonovine
- cisapride
- colchicine, if you have liver or kidney problems
- dronedarone (Multaq[®])
- pimozone (Orap[®])
- oral midazolam (Versed[®]), triazolam (Halcion[®])
- ranolazine (Ranexa[®])
- the herbal supplement St. John's Wort (*Hypericum perforatum*)
- the cholesterol lowering medicines lovastatin (Mevacor[®], Altoprev[®], Advicor[®]) or simvastatin (Zocor[®], Simcor[®], Vytorin[®])
- rifampin (Rifadin[®], Rifater[®], Rifamate[®], Rimactane[®])
- sildenafil (Revatio[®]) only when used for the treatment of pulmonary arterial hypertension.

Serious problems can happen if you or your child take any of these medicines with PREZISTA.

What should I tell my doctor before I take PREZISTA?

PREZISTA may not be right for you. Before taking PREZISTA, tell your healthcare provider if you:

- have liver problems, including hepatitis B or hepatitis C
- are allergic to sulfa medicines
- have high blood sugar (diabetes)
- have hemophilia
- are pregnant or planning to become pregnant. It is not known if PREZISTA will harm your unborn baby.

Pregnancy Registry: You and your healthcare provider will need to decide if taking PREZISTA is right for you. If you take PREZISTA while you are pregnant, talk to your healthcare provider about how you can be included in the Antiretroviral Pregnancy Registry. The purpose of the registry is follow the health of you and your baby.

- are breastfeeding or plan to breastfeed. **Do not breastfeed.** We do not know if PREZISTA can be passed to your baby in your breast milk and whether it could harm your baby. Also, mothers with HIV-1 should not breastfeed because HIV-1 can be passed to the baby in the breast milk.

Tell your healthcare provider about all the medicines you take including prescription and nonprescription medicines, vitamins, and herbal supplements. Using PREZISTA and certain other medicines may affect each other causing serious side effects. PREZISTA may affect the way other medicines work and other medicines may affect how PREZISTA works.

Especially tell your healthcare provider if you take any of the medicines listed below. The generic name is provided, followed by examples of possible brand names for the drug product:

- medicine to treat HIV
- estrogen-based contraceptives (birth control). PREZISTA might reduce the effectiveness of estrogen-based contraceptives. You must take additional precautions for birth control such as a condom.

- medicines to prevent organ transplant rejection such as cyclosporine (Gengraf[®], Sandimmune[®], Neoral[®]), tacrolimus (Prograf[®]), sirolimus (Rapamune[®])
- any medicines that contain steroids
- amiodarone (Pacerone[®], Cardarone[®])
- amitriptyline
- amlodipine
- apixaban (Eliquis[®])
- artemether/lumefantrine (Coartem[®])
- atorvastatin (Lipitor[®])
- avanafil (Stendra[®])
- bepridil (Bepadin[®], Vascor[®])
- boceprevir (Victrelis[™])
- bosentan (Tracleer[®])
- budesonide (Entocort[®], Pulmicort Flexhaler[®], Pulmicort respules[®], Rhinocort[®], Symbicort[®], Uceris[®])
- buprenorphine (Butrans[®], Buprenex[®], Subutex[®])
- buprenorphine/naloxone (Suboxone[®], Zubsolv[®])
- buspirone
- carbamazepine (Carbatrol[®], Equetro[®], Tegretol[®], Eptol[®])
- carvedilol (Coreg[®])
- clarithromycin (Prevpac[®], Biaxin[®])
- colchicine (Colcrys[®], Col-Probenecid[®])
- dabigatran etexilate (Pradaxa[®])
- dasatinib (Sprycel[®])
- desipramine (Norpramin[®])
- dexamethasone (Ozurdex[®])
- diazepam (Valium[®])
- digoxin (Lanoxin[®])
- diltiazem
- disopyramide (Norpace[®])
- estazolam

- everolimus (Zortress[®], Afinitor[®])
- felodipine (Plendil[®])
- flecainide (Tambocor[®])
- fluticasone (Advair Diskus[®], Veramyst[®], Flovent[®], Flonase[®])
- imipramine
- itraconazole (Sporanox[®], Onmel[®])
- ketoconazole (Nizoral[®])
- lidocaine (Xylocaine Viscous[®])
- methadone (Methadose[®])
- metoprolol (Lopressor[®], Toprol-XL[®])
- mexiletine (Mexitil[®])
- nifedipine (Procardia[®], Adalat CC[®], Afeditab CR[®])
- nilotinib (Tasigna[®])
- nortriptyline
- paroxetine (Paxil[®], Pexeva[®])
- phenobarbital
- phenytoin (Dilantin[®], Phenytek[®])
- posaconazole
- pravastatin (Pravachol[®])
- propafenone (Rythmol[®])
- quetiapine (Seroquel)
- quinidine (Nuedexta[®])
- rifabutin (Mycobutin[®])
- rifapentine (Priftin[®])
- risperidone (Risperdal[®])
- rivaroxaban (Xaralto[®])
- rosuvastatin (Crestor[®])
- salmeterol (Advair[®], Serevent[®])
- sertraline (Zoloft[®])
- sildenafil (Viagra[®], Revatio[®])

- simeprevir (OLYSIO™)
- tadalafil (Cialis® , Adcirca®)
- telaprevir (Incivek™)
- thioridazine (Mellaril®)
- timolol (Cosopt® , Betimol® , Timoptic® , Isatolol® , Combigan®)
- trazodone (Oleptro® , Desyrel®)
- warfarin (Coumadin® , Jantoven®)
- vardenafil (Levitra® , Staxyn®)
- verapamil
- vinblastine
- vincristine (Marquibo®)
- voriconazole (VFend®)
- zolpidem (Ambien®)

This is **not** a complete list of medicines that you should tell your healthcare provider that you are taking. Ask your healthcare provider or pharmacist if you are not sure if your medicine is one that is listed above. Know the medicines you take. Keep a list of them to show your doctor or pharmacist when you get a new medicine. Do not start any new medicines while you are taking PREZISTA without first talking with your healthcare provider.

How should I take PREZISTA?

- Take PREZISTA every day exactly as prescribed by your healthcare provider.
- You must take ritonavir (NORVIR®) at the same time as PREZISTA.
- Do not change your dose of PREZISTA or stop treatment without talking to your healthcare provider first.
- Take PREZISTA and ritonavir (NORVIR®) with food.
- Swallow PREZISTA tablets whole with a drink. If you have difficulty swallowing PREZISTA tablets, PREZISTA oral suspension is also available. Your health care provider will help decide whether PREZISTA tablets or oral suspension is right for you.
- If your child is taking PREZISTA, your child's healthcare provider will decide the right dose based on your child's weight. Your

child's healthcare provider will tell you how much PREZISTA (tablets or oral suspension) and how much ritonavir (NORVIR[®]) (capsules, tablets or solution) your child should take. Your child should take PREZISTA with ritonavir with food. If your child does not tolerate ritonavir oral solution, ask your child's healthcare provider for advice.

- PREZISTA oral suspension should be given with the supplied oral dosing syringe. Shake the suspension well before each use. See the Instructions for Use that come with PREZISTA oral suspension for information about the right way to prepare and take a dose.
- If your or your child's prescribed dose of PREZISTA oral suspension is more than 6 mL, you will need to divide the dose. Follow the instructions given to you by your healthcare provider or pharmacist about how to divide the dose. Ask your healthcare provider or pharmacist if you are not sure.
- If you take too much PREZISTA, call your healthcare provider or go to the nearest hospital emergency room right away.

What should I do if I miss a dose?

People who take PREZISTA one time a day:

- If you miss a dose of PREZISTA by less than 12 hours, take your missed dose of PREZISTA right away. Then take your next dose of PREZISTA at your regularly scheduled time.
- If you miss a dose of PREZISTA by more than 12 hours, wait and then take the next dose of PREZISTA at your regularly scheduled time.

People who take PREZISTA two times a day

- If you miss a dose of PREZISTA by less than 6 hours, take your missed dose of PREZISTA right away. Then take your next dose of PREZISTA at your regularly scheduled time.
- If you miss a dose of PREZISTA by more than 6 hours, wait and then take the next dose of PREZISTA at your regularly scheduled time.

If a dose of PREZISTA is skipped, do not double the next dose. Do not take more or less than your prescribed dose of PREZISTA at any one time.

What are the possible side effects of PREZISTA?

PREZISTA can cause side effects including:

- See **“What is the most important information I should know about PREZISTA?”**
- **Diabetes and high blood sugar (hyperglycemia).** Some people who take protease inhibitors including PREZISTA can get high blood sugar, develop diabetes, or your diabetes can get worse. Tell your healthcare provider if you notice an increase in thirst or urinate often while taking PREZISTA.
- **Changes in body fat.** These changes can happen in people who take antiretroviral therapy. The changes may include an increased amount of fat in the upper back and neck (“buffalo hump”), breast, and around the back, chest, and stomach area. Loss of fat from the legs, arms, and face may also happen. The exact cause and long-term health effects of these conditions are not known.
- **Changes in your immune system (Immune Reconstitution Syndrome)** can happen when you start taking HIV medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Call your healthcare provider right away if you start having new symptoms after starting your HIV medicine.
- **Increased bleeding for hemophiliacs.** Some people with hemophilia have increased bleeding with protease inhibitors including PREZISTA.

The most common side effects of PREZISTA include:

- diarrhea
- nausea
- rash
- headache
- abdominal pain
- vomiting

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of PREZISTA. For more information, ask your health care provider.

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

How should I store PREZISTA?

- Store PREZISTA oral suspension and tablets at room temperature [77°F (25°C)].
- Do not refrigerate or freeze PREZISTA oral suspension.
- Keep PREZISTA away from high heat.
- PREZISTA oral suspension should be stored in the original container.

Keep PREZISTA and all medicines out of the reach of children.

General information about PREZISTA

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use PREZISTA for a condition for which it was not prescribed. Do not give PREZISTA to other people even if they have the same condition you have. It may harm them.

This leaflet summarizes the most important information about PREZISTA. If you would like more information, talk to your healthcare provider. You can ask your healthcare provider or pharmacist for information about PREZISTA that is written for health professionals.

For more information, call 1-800-526-7736.

What are the ingredients in PREZISTA?

Active ingredient: darunavir

Inactive ingredients:

PREZISTA Oral Suspension: hydroxypropyl cellulose, microcrystalline cellulose, sodium carboxymethylcellulose, methylparaben sodium, citric acid monohydrate, sucralose, masking flavor, strawberry cream flavor, hydrochloric acid (for pH adjustment), purified water.

PREZISTA 75 mg and 150 mg Tablets: colloidal silicon dioxide, crospovidone, magnesium stearate, microcrystalline cellulose. The film

coating contains: OPADRY[®] White (polyethylene glycol 3350, polyvinyl alcohol-partially hydrolyzed, talc, titanium dioxide).

PREZISTA 600 mg Tablets: colloidal silicon dioxide, crospovidone, magnesium stearate, microcrystalline cellulose. The film coating contains: OPADRY[®] Orange (FD&C Yellow No. 6, polyethylene glycol 3350, polyvinyl alcohol-partially hydrolyzed, talc, titanium dioxide).

PREZISTA 800 mg Tablets: colloidal silicon dioxide, crospovidone, magnesium stearate, microcrystalline cellulose, hypromellose. The film coating contains: OPADRY[®] Dark Red (iron oxide red, polyethylene glycol 3350, polyvinyl alcohol-partially hydrolyzed, talc, titanium dioxide).

This Patient Information has been approved by the U.S Food and Drug Administration.

Product of Ireland

Manufactured by:

PREZISTA Oral Suspension
Janssen Pharmaceutica, N.V.
Beerse, Belgium

PREZISTA Tablets
Janssen Ortho LLC, Gurabo, PR 00778

Manufactured for:

Janssen Therapeutics, Division of Janssen Products, LP,
Titusville NJ 08560

Revised: May 2015

NORVIR[®] is a registered trademark of its respective owner.

PREZISTA[®] is a registered trademark of Janssen Pharmaceuticals

© Janssen Pharmaceuticals, Inc. 2006