



 <p>Actelion Pharmaceuticals Ltd. Gewerbstrasse 16 CH-4123 Allschwil Switzerland E-mail: top-text_artwork@actelion.com</p>	<b>Product name</b> <b>Non-medicated Valchlor Demonstration Gel US</b> <b>Front tube label</b>		<b>Printing colors</b> <div> <div>Process black</div> <div>Pantone 199</div> <div></div> </div>
	<b>Mock up code</b> <b>VAL-FTL-non-US-514</b>	<b>Proof no. / date</b> <b>4 / 22-Apr 2015</b>	
	<b>Dimension</b> <b>1.5" x 0.625"</b>	<b>Technical drawing no</b>	<b>Technical colors</b>



 <b>ACTELION</b>  Actelion Pharmaceuticals Ltd. Gewerbestrasse 16 CH-4123 Allschwil Switzerland E-mail: top-text_artwork@actelion.com	Product name <b>Non-medicated Valchlor Demonstration Gel US</b> <b>Back tube label</b>		Printing colors <b>Process black</b> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="width: 40px; height: 20px; background-color: black;"></div> <div style="width: 60px; height: 20px; border: 1px solid black;"></div> <div style="width: 60px; height: 20px; border: 1px solid black;"></div> </div>
	Mock up code <b>VAL-BTL-non-US-515</b>	Proof no. / date <b>2 / 22-Apr 2015</b>	
	Dimension <b>1.5" x 0.625"</b>	Technical drawing no	Technical colors <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="width: 60px; height: 20px; border: 1px solid black;"></div> <div style="width: 60px; height: 20px; border: 1px solid black;"></div> </div>



 <p>Actelion Pharmaceuticals Ltd. Gewerbstrasse 16 CH-4123 Allschwil Switzerland E-mail: top-text_artwork@actelion.com</p>	<b>Product name</b> <b>Non-medicated Valchlor Demonstration Gel US</b> <b>Outer Carton</b>		<b>Printing colors</b> <div> <div>Process black</div> <div>Pantone 199</div> </div>	<div> <div></div> <div></div> </div>
	<b>Mock up code</b> <b>VAL-PC-non-US-571</b>	<b>Proof no. / date</b> <b>5 / 28-Apr 2015</b>		
	<b>Dimension</b> <b>28.57 x 28.57 x 82.55 mm</b>	<b>Technical drawing no</b> <b>150459 (Nosco)</b>	<b>Technical colors</b> <div>die cut</div>	

## Non-medicated VALCHLOR<sup>®</sup> Demonstration Gel

### Instructions For Use<sup>1</sup>

Non-medicated VALCHLOR<sup>®</sup> Demonstration Gel **DOES NOT** contain mechlorethamine (the medicine in VALCHLOR).


The purpose of VALCHLOR Demonstration Gel is to show healthcare providers and patients how to properly apply VALCHLOR. This will also allow the user to experience the consistency of VALCHLOR and how it may feel on the skin.

This information does not take the place of talking to your healthcare provider about your condition or your treatment.


#### How should VALCHLOR Demonstration Gel be used?

- Use VALCHLOR Demonstration Gel on clean and dry, healthy skin
- If you take a bath or shower, be sure to use the gel after your skin is dry (at least 30 minutes after bathing or showering)

#### How to use VALCHLOR Demonstration Gel:




STEP 1: Open the 5g tube, use cap to puncture seal



STEP 2: Squeeze a small pea-sized amount on your fingertip

---

<sup>1</sup> Reference VALCHLOR US Prescribing Information.



STEP 3: Gently spread a thin layer of the gel to dry healthy skin



STEP 4: Allow the gel to dry before covering with clothing (5 to 10 minutes after applying)



STEP 5: Throw the tube away in the household trash



STEP 6: Wash your hands with soap and water

**Some Important Do's and Don'ts about VALCHLOR Demonstration Gel:**

- When applying VALCHLOR Demonstration Gel, avoid the eyes, mouth, and nose.
- VALCHLOR Demonstration Gel should only be used on healthy skin since it is non-medicated.
- The 5g tube of VALCHLOR Demonstration Gel is meant for single use only.
- VALCHLOR Demonstration Gel contains alcohol. Since alcohol-based gels are flammable, avoid fire, flame, and smoking until the gel has dried.
- VALCHLOR Demonstration Gel can be stored at room temperature.
- Keep out of reach of children.

This demonstration gel contains the following nonactive ingredients: diethylene glycol monoethyl ether, propylene glycol, isopropyl alcohol, glycerin, lactic acid, hydroxypropylcellulose, sodium chloride, menthol, edetate disodium, and butylated hydroxytoluene.

Please see VALCHLOR full Prescribing Information including Medication Guide for information for active drug product.

To report **SUSPECTED ADVERSE REACTIONS** to the VALCHLOR Demonstration Gel, contact Actelion Pharmaceuticals US, Inc., at 1-855-4-VALCHLOR (1-855-482-5245) or FDA at 1-800-FDA-1088 or visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

ACT20150424