



art bleeds off 1/16" (0.0625") out from die (dash line does not print)


This line indicates die and does not print

no copy 1/16" (0.0625") in from die (dash line does not print)

Gloss and Matt Varnish - Full size file on Varnish layer



MICRO PRINT (0.8 pt dedicated plate)




3/16" HINGE AREA UNVARNISHED



This line indicates die and does not print

no copy 1/16" (0.0625") in from die (dash line does not print)

BASE



This line indicates die and does not print

no copy 1/16" (0.0625") in from die (dash line does not print)

3/16" HINGE AREA UNVARNISHED

RELEASED BACK

PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
Date 21-NOV-2014 PASS # 1 Component # PAA056448 Drawing # (b) (4) -50013882v04	(b) (4)
Identification	
	(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	1 pt, .5 pt



PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
Date 24-NOV-2014 PASS # 2 Component # PAA056698 Drawing # (b) (4) 50021037_v02	(b) (4)
Identification	
	(b) (4)

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<.39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

UNCOATED
NO COPY
NO PRINT

128 Code = 056449

128 CODE
AREA

128 Code = 056449


128 CODE
AREA

128 Code = 056449


128 CODE
AREA

MICRO PRINT
LOCATION

Gloss and Matte Varnish -
Full size file on Varnish layer



Emboss -
Full size file on Emboss layer*
* There are two levels of Embossing.



KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts
Active ingredient (in each capsule) Purpose
Esomeprazole 20 mg Acid reducer
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent chest pain
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking
■ warfarin, clopidogrel or cilostazol (blood-thinning medicines)
■ prescription antifungal or anti-yeast medicines
■ digoxin (heart medicine)
■ diazepam (anxiety medicine)
■ tacrolimus or mycophenolate mofetil (immune system medicines)
■ prescription antiretrovirals (medicines for HIV infection)
■ methotrexate (arthritis medicine)

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days

Drug Facts (continued)
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ may take 1 to 4 days for full effect
14-Day Course of Treatment
■ swallow 1 capsule with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 capsule a day
■ swallow whole. Do not crush or chew capsules.
■ do not use for more than 14 days unless directed by your doctor
Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F)

Inactive ingredients
corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments?
call toll-free 1-866-226-1600

Tips for Managing Heartburn
• Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
• Eat slowly and do not eat big meals.
• Do not eat late at night or just before bedtime.
• Do not lie flat or bend over soon after eating.
• Raise the head of your bed.
• Wear loose-fitting clothing around your stomach.
• If you are overweight, lose weight.
• If you smoke, quit smoking.



Marketed by:
Pfizer, Madison, NJ 07940 USA
© 2014 Pfizer Inc.
Made in France
For most recent product information, visit
www.Nexium24HR.com
Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

Nexium[®]
24HR | Clinically Proven to
Treat Frequent Heartburn

Treats Frequent Heartburn



28
CAPSULES

Two 14-day courses of treatment

May take 1 to 4 days for full effect

2 BOTTLES
INSIDE



Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

PAA056449



128 CODE
AREA

128 Code = 056449

128 CODE AREA - SUBSET C 10 MIL DENSITY
WITH 4 OR 6 DIGITS (CODE TO EXTEND
TO EDGE OF FLAP - UNVARNISHED)
PRINTED SIDE OF CARTON
128 Code = 056449

128 CODE
AREA

MICRO PRINT
LOCATION

128 Code = 056449

LOT & EXP AREA.
DARK COLOR FOR
LASER PRINTING.

3/16" DATA MATRIX CODE
WITH 1/8" QUIET ZONE
ON ALL SIDES
DATA MATRIX CODE = 056449

PRODUCTION

(NOT COLOR ACCURATE)

Job Information	
Date	21-NOV-2014 PASS # 1
Component #	PAA056449
Drawing #	(b)(4) 50013884v05

Identification	
Pfizer	Consumer Healthcare (b)(4)

Color Legend	
(b)(4)	(b)(4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Treats Frequent Heartburn

Nexium[®]

(esomeprazole magnesium)
Delayed-Release Capsules
22.3 mg/Acid Reducer


24HR

May take 1 to 4 days for full effect

2 Pack

28
CAPSULES

Two 14-day courses
of treatment



D000050549



Marketed by: Pfizer, Madison, NJ 07940 USA
© 2014 Pfizer Inc.
Made in France
For most recent product information, visit www.Nexium24HR.com
Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.


REMOVE VARNISH
WHITE HOLDOUT
LOT/EXP

WHITE HOLDOUT
QUIET AREA
DATA MATRIX CODE
050549

Drug Facts
Active ingredient (in each capsule) Esomeprazole 20 mg Acid reducer
Purpose Esomeprazole 20 mg Acid reducer
Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
Warnings ■ **Do not use if you have:** ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulder; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ nausea or vomiting ■ unexplained weight loss ■ stomach pain
Ask a doctor before use if you are taking ■ prescription antifungal or anti-yeast medicines ■ dioxin (heart medicine) ■ diazepam (anxiety medicine) ■ tacrolimus or mycophenolate mofetil (immune system medicines) ■ prescription antiretrovirals (medicines for HIV infection) ■ methotrexate (arthritis medicine)
Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days
Other information ■ read the directions and warnings before use ■ keep the card. It contains important information. ■ store at 20-25°C (68-77°F)
Inactive ingredients com starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate
Questions or comments? call toll-free 1-866-226-1600

KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Gloss and Matte Varnish -
Full size file on Varnish layers



Tips for Managing Heartburn
• Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
• Eat slowly and do not eat big meals.
• Do not eat late at night or just before bedtime.
• Do not lie flat or bend over soon after eating.
• Raise the head of your bed.
• Wear loose-fitting clothing around your stomach.
• If you are overweight, lose weight.
• If you smoke, quit smoking.



Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
Date 02-DEC-2014 PASS # 1 Component # D000050549 Drawing # 119216AP10	(b) (4)
Identification	
Consumer Healthcare	(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	10 pt
• DRUG FACTS CONTINUED	9 pt
• HEADINGS	9 pt
• SUBHEADINGS/BODY TEXT	7 pt
• LEADING	7.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	6 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

UNCOATED
NO COPY
NO PRINT

128 CODES = 056450

128 CODE
AREA


128 CODE
AREA

128 CODE
AREA

MICRO PRINT
LOCATION

Gloss and Matte Varnish -
Full size file on Varnish layer

Emboss -
Full size file on Emboss layer*
* There are two levels of Embossing.



KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts
Active ingredient (in each capsule) Purpose
Esomeprazole 20 mg Acid reducer
(Each delayed-release capsule corresponds to 22.3 mg
esomeprazole magnesium trihydrate)

Uses
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take
1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody
or black stools
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath; sweating;
pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent chest pain
These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more
serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain
Ask a doctor or pharmacist before use if you are taking
■ warfarin, clopidogrel or cimetidine (blood-thinning medicines)
■ prescription antifungal or anti-yeast medicines
■ digoxin (heart medicine)
■ diazepam (anxiety medicine)
■ tacrolimus or mycophenolate mofetil (immune system
medicines)
■ prescription antiretrovirals (medicines for HIV infection)
■ methotrexate (arthritis medicine)
Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days

Drug Facts (continued)
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help
or contact a Poison Control Center right away.

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day
for 14 days
■ may take 1 to 4 days for full effect
14-Day Course of Treatment
■ swallow 1 capsule with a glass of water before eating in the
morning
■ take every day for 14 days
■ do not take more than 1 capsule a day
■ swallow whole. Do not crush or chew capsules.
■ do not use for more than 14 days unless directed by your doctor
Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every
4 months unless directed by a doctor
■ children under 18 years of age: ask a doctor before use. Heartburn
in children may sometimes be caused by a serious condition.

Other Information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F)

Inactive ingredients
corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40,
ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose,
hypromellose, magnesium stearate, methacrylic acid copolymer,
pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide,
triethyl citrate

Questions or comments?
call toll-free 1-866-226-1600

**Tips for
Managing Heartburn**
• Avoid foods or drinks that are
more likely to cause heartburn, such
as rich, spicy, fatty and fried foods,
chocolate, caffeine, alcohol and even
some acidic fruits and vegetables.
• Eat slowly and do not eat big meals.
• Do not eat late at night or just before
bedtime.
• Do not lie flat or bend over soon
after eating.
• Raise the head of your bed.
• Wear loose-fitting clothing around
your stomach.
• If you are overweight, lose weight.
• If you smoke, quit smoking.



Marketed by:
Pfizer, Madison, NJ 07940 USA
© 2014 Pfizer Inc.
Made in France
For most recent product information, visit
www.Nexium24HR.com
Nexium and the colors purple and gold
as applied to the capsule are registered
trademarks of AstraZeneca AB and are
used under license.

Nexium[®]
24HR | Clinically Proven to
**Treat Frequent
Heartburn**

Treats Frequent Heartburn



**42
CAPSULES**
Three 14-day courses of treatment

May take 1 to 4 days for full effect

**3 BOTTLES
INSIDE**



Do Not Use if seal under
bottle cap imprinted with
"SEALED FOR YOUR PROTECTION"
or yellow band around the center of
each capsule is broken or missing.



PAA056450

128 CODE
AREA

128 CODE
AREA

128 CODE
AREA

MICRO PRINT
LOCATION

LOT & EXP AREA.
DARK COLOR FOR
LASER PRINTING.

3/16" DATA MATRIX CODE
WITH 1/8" QUIET ZONE
ON ALL SIDES
DATA MATRIX CODE = 056450

128 CODE AREA - SUBSET C 10 MIL DENSITY
WITH 4 OR 6 DIGITS (CODE TO EXTEND
TO EDGE OF FLAP - UNVARNISHED)
PRINTED SIDE OF CARTON

PRODUCTION (NOT COLOR ACCURATE)

Job Information	Color Legend
Date 21-NOV-2014 PASS # 1 Component # PAA056450 Drawing # (b) (4) 50013884v05	(b) (4)
Identification	
Consumer Healthcare (b) (4)	

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

