

Proposed Blister Label for Cetirizine Hydrochloride Orally Disintegrating Tablets, 10 mg (6 tablets)



Proposed Carton Label for Cetirizine Hydrochloride Orally Disintegrating Tablets, 10 mg - 12 tablets (2X6)

Cetirizine Hydrochloride
Orally Disintegrating Tablets
10 mg
 Antihistamine
Dissolve Tabs



Melts in Your Mouth
ORANGE FLAVOR

NDC 49884 229 87

Cetirizine Hydrochloride
Orally Disintegrating Tablets
10 mg
 Antihistamine

Dissolve Tabs

24 HOUR RELIEF OF:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

ALLERGY
 INDOOR + OUTDOOR
 ALLERGIES

12
 ORALLY
 DISINTEGRATING
 TABLETS
10 mg each

Cetirizine Hydrochloride
 Orally Disintegrating Tablets
 10 mg
 Antihistamine

LOT: _____
 EXP: _____

Distributed by:
 Par Pharmaceutical Companies, Inc.
 Spring Valley, NY 10977 U.S.A.

Manufactured by:
 Par Formulations Private Limited,
 1/58, Pudupakkam,
 Kelambakkam 603 103.
 Made in India
 10/2/15



N
 2 4 9 8 8 4 1 2 2 9 8 7 0

Dissolve Tabs
 Antihistamine
10 mg
 Cetirizine Hydrochloride
 Orally Disintegrating Tablets

Drug Facts		Drug Facts (continued)							
<p>Active ingredient (in each tablet) Purpose Cetirizine Hydrochloride 10 mg.....Antihistamine</p> <p>Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: • runny nose • sneezing • itchy, watery eyes • itching of the nose or throat</p> <p>Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.</p> <p>Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.</p> <p>Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.</p> <p>When using this product • drowsiness may occur • avoid alcoholic drinks • alcohol, sedatives, and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery</p> <p>Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.</p> <p>If pregnant or breast-feeding: • if breast-feeding: not recommended • if pregnant: ask a health professional before use.</p>	<p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions Tablet melts in mouth. Can be taken with or without water.</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border-right: 1px solid black; padding: 2px;">adults and children 6 years and over</td> <td style="padding: 2px;">one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 2px;">adults 65 years and over</td> <td style="padding: 2px;">ask a doctor</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 2px;">children under 6 years of age</td> <td style="padding: 2px;">ask a doctor</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 2px;">consumers with liver or kidney disease</td> <td style="padding: 2px;">ask a doctor</td> </tr> </table> <p>Other information • store between 20° to 25°C (68° to 77°F). Avoid high humidity. • do not use if carton or blister unit is opened or broken • see side panel for lot number and expiration date</p> <p>Inactive ingredients butyl hydroxyanisole, butyl hydroxytoluene, colloidal silicon dioxide, corn starch, croscopolone, edetate disodium, hydroxy propyl beta cyclodextrin, magnesium stearate, microcrystalline cellulose, maize maltodextrin, peppermint and orange flavor, and sucralose.</p> <p>Questions? call 1-800-828-9393</p>	adults and children 6 years and over	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.	adults 65 years and over	ask a doctor	children under 6 years of age	ask a doctor	consumers with liver or kidney disease	ask a doctor
adults and children 6 years and over	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.								
adults 65 years and over	ask a doctor								
children under 6 years of age	ask a doctor								
consumers with liver or kidney disease	ask a doctor								

Proposed Carton Label for Cetirizine Hydrochloride Orally Disintegrating Tablets, 10 mg - 24 tablets (4X6)

Cetirizine Hydrochloride
Orally Disintegrating Tablets
10 mg
 Antihistamine
Dissolve Tabs



Melts in Your Mouth
ORANGE FLAVOR

190

NDC 49884 229 35

Original Prescription Strength

Cetirizine Hydrochloride
Orally Disintegrating Tablets
10 mg
 Antihistamine

Dissolve Tabs



24 HOUR RELIEF OF:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

ALLERGY

INDOOR + OUTDOOR
 ALLERGIES

24
 ORALLY
 DISINTEGRATING
 TABLETS
10 mg each

Distributed by:
 Par Pharmaceutical Companies, Inc.
 Spring Valley, NY 10977 U.S.A.

Manufactured by:
 Par Formulations Private Limited,
 1/58, Pudupakkam,
 Kelambakkam 603 103.
 Made in India
 10/2/15



N
 2 4 9 8 8 4 1 2 2 9 3 5 1

Dissolve Tabs

Antihistamine
10 mg

Cetirizine Hydrochloride
Orally Disintegrating Tablets

Cetirizine Hydrochloride
Orally Disintegrating Tablets
 10 mg
 Antihistamine

Un varnish Area

LOT :
 EXP :

Drug Facts	Drug Facts (continued)								
<p>Active ingredient (in each tablet) Purpose Cetirizine Hydrochloride 10 mg.....Antihistamine</p> <p>Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:</p> <ul style="list-style-type: none"> • runny nose • sneezing • itchy, watery eyes • itching of the nose or throat <p>Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.</p> <p>Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.</p> <p>Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.</p> <p>When using this product</p> <ul style="list-style-type: none"> • drowsiness may occur • avoid alcoholic drinks • alcohol, sedatives, and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery <p>Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.</p> <p>If pregnant or breast-feeding:</p> <ul style="list-style-type: none"> • if breast-feeding: not recommended • if pregnant: ask a health professional before use. 	<p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions Tablet melts in mouth. Can be taken with or without water.</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%;">adults and children 6 years and over</td> <td style="width: 50%;">one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.</td> </tr> <tr> <td>adults 65 years and over</td> <td>ask a doctor</td> </tr> <tr> <td>children under 6 years of age</td> <td>ask a doctor</td> </tr> <tr> <td>consumers with liver or kidney disease</td> <td>ask a doctor</td> </tr> </table> <p>Other information</p> <ul style="list-style-type: none"> • store between 20° to 25°C (68° to 77°F). Avoid high humidity. • do not use if carton or blister unit is opened or broken • see side panel for lot number and expiration date <p>Inactive ingredients butyl hydroxyanisole, butyl hydroxytoluene, colloidal silicon dioxide, corn starch, crospovidone, edetate disodium, hydroxy propyl beta cyclodextrin, magnesium stearate, microcrystalline cellulose, maize maltodextrin, peppermint and orange flavor, and sucralose.</p> <p>Questions? call 1-800-828-9393</p>	adults and children 6 years and over	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.	adults 65 years and over	ask a doctor	children under 6 years of age	ask a doctor	consumers with liver or kidney disease	ask a doctor
adults and children 6 years and over	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.								
adults 65 years and over	ask a doctor								
children under 6 years of age	ask a doctor								
consumers with liver or kidney disease	ask a doctor								