

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- If needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients

acesulfame potassium, anhydrous citric acid, D&C red no. 33, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?

Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA
 ©J&JCI 2015
www.motrin.com 00000000

New Information: Read Entire Label

NDC 50580-603-04

For Ages 2 to 11 Years

Children's Motrin®

contains **Ibuprofen Oral Suspension (NSAID)**
100 mg per 5 mL

Pain Reliever / Fever Reducer

Lasts up to 8 hours

Alcohol Free

Grape Flavor
4 fl oz (120 mL)



Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL)

Ibuprofen 100 mg (NSAID)*.....Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery



Children's Motrin®

contains
Ibuprofen Oral Suspension (NSAID)

Bubble Gum Flavor

New Information: Read Entire Label

NDC 50580-604-04

For Ages 2 to 11 Years

Children's Motrin®

contains **Ibuprofen Oral Suspension (NSAID)**
100 mg per 5 mL

Pain Reliever / Fever Reducer

Lasts up to
8 hours

Alcohol Free

Bubble Gum Flavor

4 fl oz (120 mL)



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- If needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients

acesulfame potassium, anhydrous citric acid, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?

Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2015
www.motrin.com 00000000

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Ibuprofen 100 mg (NSAID)*	Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses

- temporarily:
- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
 - reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- shock
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

Active ingredient (in each 5 mL): Ibuprofen 100 mg (NSAID)[†] Purpose: Pain reliever/fever reducer

Warnings: Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, wheezing, rash, or skin redness. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is higher if your child has had stomach ulcers or bleeding problems, takes a blood thinner (an anticoagulant) or steroid drug, or has had ulcers combining prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) or takes more than 10 days. **Severe:** Stomach bleeding, heart failure, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days to relieve pain in children under 3 years of age unless directed by doctor. Do not use until the child has recovered from all other pain relievers. Ask a doctor before use if the child has a history of stomach problems, serious side effects from taking pain relievers, or if the child has been drinking fluids, has lost a lot of fluid due to vomiting or diarrhea, has high blood pressure, heart disease, liver or kidney disease, or is taking a diuretic. Ask a doctor or pharmacist before use if the child is under 6 months of age or for any serious condition. When using this product, do not take with food or milk if stomach upset occurs. If a doctor tells you to stop use, stop use if you experience any of the following signs of stomach bleeding: vomit with blood, has bloody or black stools, has stomach pain that does not get better, or if the child does not get any relief within 24 hours of treatment. If the child has a fever, do not use for more than 3 days unless a doctor says to use it.

Directions: Shake well before using. Use only the right dose on the label. If possible, use weight to dose, or use age. Use only enclosed dosing cup. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain child resistance.

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

Other information: Each 5 mL contains sodium 2 mg. Store between 20-25°C (68-77°F). Do not use if the bottle wrap is imprinted "SAFETY SEAL" is broken or missing. Inactive ingredients: acacia, croscarmellose, hydroxypropyl methylcellulose, polyethylene glycol, polyethylene glycol, purified water, sodium benzoate, sucrose, xanthan gum. Questions or comments? Call 1-877-895-5665 (Toll-Free) or 215-273-4933 (collect).
 Distributed by: JOHNSON & JOHNSON CONSUMER INC.
 Johnson Consumer Healthcare Division
 Fort Washington, PA 19034 USA ©J&JCI 2015 www.motrin.com 00000000

(b) (4)



Children's Motrin®

contains
Ibuprofen Oral Suspension (NSAID)

Original Berry Flavor

NDC 50580-601-20

For Ages 2 to 11 Years

Children's Motrin®

contains
Ibuprofen
Oral Suspension (NSAID)
100 mg per 5 mL

Pain Reliever / Fever Reducer

Lasts up to
8 hours

Alcohol Free

Original
Berry Flavor



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- If needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients

acesulfame potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?

Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2015
www.motrin.com 00000000

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Ibuprofen 100 mg (NSAID)*	Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- shock
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

Active ingredient (in each 5 mL): Ibuprofen 100 mg (NSAID)[†]

Purpose: Pain reliever/fever reducer

†Nonsteroidal anti-inflammatory drug

Uses: Temporarily relieves minor aches and pains (due to the common cold, flu, sore throat, headache and toothache) and reduces fever.

Warnings: Allergy alert: Ibuprofen may cause an allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, wheezing, rash, or skin redness. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is higher if your child has had stomach ulcers or bleeding problems, or takes a blood thinner (an anticoagulant) or steroid drug, or takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) or takes more than one longer than directed. **Severe:** Sore throat, fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to child under 3 years of age unless directed by doctor. **Do not use until the child has recovered from all other illnesses.** Ask a doctor before use if stomach bleeding warning applies to your child or if child has a history of stomach problems, such as heartburn or indigestion, or if child has serious side effects from taking pain relievers or other drugs.

NDC 50589-601-20

Children's Motrin

contains **Ibuprofen Oral Suspension (NSAID)**

100 mg per 5 mL

Pain Reliever/Fever Reducer

Original Berry Flavor

4 fl oz (120 mL)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Directions: This product does not contain directions or complete warnings for a full use. Do not give more than directed. Shake well before using. 5 mL = milliliter. Find right dose on label below. If possible, use weight to dose; otherwise use age. Use only enclosed dosing cup. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain child resistance.

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information: Each 5 mL contains sodium 2 mg. Store between 20-25°C (68-77°F). Do not use if bottle wrap imprinted "SAFETY SEAL" is broken or missing. Inactive ingredients: saccharin potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, iron oxide black, polyethylene glycol 400, propylparaben, purified water, sodium benzoate, sucrose, vanillin gum. Questions or comments? Call 1-877-895-5663 (toll-free) or 215-273-8755 (toll-free).

Distributed by: JOHNSON & JOHNSON CONSUMER INC.

McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA ©J&JCI 2015 www.motrin.com 00000000

(b) (4)



Children's Motrin®

DYE - FREE

contains
Ibuprofen Oral Suspension (NSAID)
Non-Staining
Original Berry Flavor

NDC 50580-184-50

For Ages 2 to 11 Years

Children's Motrin®

contains **Ibuprofen Oral Suspension (NSAID)**
100 mg per 5 mL
Pain Reliever / Fever Reducer

DYE - FREE

Lasts up to 8 hours

Alcohol Free

Non-Staining
Original Berry Flavor
4 fl oz (120 mL)



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2015
www.motrin.com 00000000

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Ibuprofen 100 mg (NSAID)*	Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.


Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- If needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients

acesulfame potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?

Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA
 ©J&JCI 2015
www.motrin.com 00000000

New Information: Read Entire Label

NDC 50580-601-21

FOR HOSPITAL/GOVERNMENT USE ONLY

For Ages 2 to 11 Years

Children's Motrin®
 contains **Ibuprofen**
Oral Suspension (NSAID)
 100 mg per 5 mL

Pain Reliever / Fever Reducer

Original Berry Flavor

Lasts up to 8 hours

Alcohol Free

Not for Retail Sale

4 fl oz (120 mL)



Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Ibuprofen 100 mg (NSAID)*	Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- shock
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

FOR HOSPITAL/GOVERNMENT USE ONLY
Not for Retail Sale NDC 50590-001-21

Children's Motrin®
contains Ibuprofen
Oral Suspension (NSAID)
100 mg per 5 mL
Pain Reliever/Fever Reducer 4 fl oz (120 mL)
Original Berry Flavor

Active ingredient (in each 5 mL): Ibuprofen 100 mg (NSAID)
Purpose: Pain reliever/fever reducer
Indications: Ibuprofen is used to relieve minor aches and pains due to the common cold, flu, sore throat, headache and toothache.

Warnings: Allergy alert: Ibuprofen may cause an allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, wheezing, rash, or skin redness. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is higher if your child has had stomach ulcers or bleeding problems, takes a blood thinner (an anticoagulant) or steroid drug, or takes drugs containing aspirin or non-steroidal NSAIDs (aspirin, ibuprofen, naproxen, or others) for a long time. **Severe:** Stomach bleeding, fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days to relieve fever in children under 3 years of age unless directed by doctor. Do not use with other pain relievers or aspirin unless directed by doctor. Ask a doctor before use if your child has a history of stomach problems or serious side effects from taking pain relievers or aspirin.

Directions: Shake well before using. Use only the enclosed dosing cup. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain child resistance.

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

Other information: Each 5 mL contains sodium 2 mg. Store between 20-25°C (68-77°F). Do not use if the bottle wrap imprinted "SAFETY SEAL" is broken or missing. **Inactive ingredients:** saccharin potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, iron oxide black, polyethylene glycol 400, propylparaben, purified water, sodium benzoate, sucrose, vanillin gum. Questions or comments? Call 1-877-895-5663 (toll-free) or 215-273-8755 (local).
Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McKeesport Consumer Healthcare Division
Fort Washington, PA 19034 USA ©J&JCI 2015 www.motrin.com 00000000

(b) (4)

Trim Safe Area Bleed Score

w 3.875"

Convenient 2-PACK!

NDC 50580-601-25

For Ages 2 to 11 Years

**Children's
Motrin[®]**

contains ibuprofen
Oral Suspension (NSAID)
100 mg per 5 mL

Pain Reliever / Fever Reducer

**Lasts up to
8 hours**

Alcohol Free

*Original
Berry Flavor*

NDC 50580-601-25

For Ages 2 to 11 Years

**Children's
Motrin[®]**

contains ibuprofen
Oral Suspension (NSAID)
100 mg per 5 mL

Pain Reliever / Fever Reducer

**Lasts up to
8 hours**

Alcohol Free

*Original
Berry Flavor*

h 5.65625"

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2015 www.motrin.com
McN-XX-0000



3 0045-0192-08 0

Score 0.65625"
from bottom

(b) (4)



Drug Facts (continued)

Ask a doctor before use if:

- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is:

- under a doctor's care for any serious condition
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
 - child does not get any relief within last day (24 hours) of treatment
 - fever or pain gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away, (1-800-232-1222)

Directions

- this product does not contain directions or complete warnings for adult use—do not give more than directed
- shake well before using
- mL = milliliter
- find right dose in chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain child resistance.

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other Information

- each 5 mL contains sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL" is broken or missing

Inactive Ingredients ascorbate potassium, anhydrous citric acid, D-xylose, yellow no. 10, FD&C red no. 40, flavors, glycerin, polyacrylate-80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)



New Information: Read Entire Label For Ages 2 to 11 Years

Motrin
contains **Ibuprofen Oral Suspension**
100 mg per 5 mL

Original Berry Flavor
1 fl oz (30 mL)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active Ingredient (in each 5 mL)	Purpose
Ibuprofen 100 mg (NSAID)	Pain reliever/fever reducer
*Non-steroidal anti-inflammatory drug	

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- stomach pain
- skin redness
- rash
- blistering

 If a allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing aspirin or other NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed
- has a liver or kidney problem
- has a history of stomach or intestinal bleeding
- has a history of stomach or intestinal pain
- has a history of stomach or intestinal pain
- has a history of stomach or intestinal pain

 Severe or persistent stomach or intestinal pain may be accompanied by high fever, headache, nausea, and vomiting may be seen. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use:

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

USA
000000
000

LOT:

EXP:



This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
06/16/2016