



Allegra-D® Allergy and Congestion 12 Hour 05.02.16

ALG-D_12hr_10ct_Blister_2015



Allegra-D Allergy & Congestion 24 Hour Blister Carton

05.02.16

ALG-D_24hr_15ct_CTN_2015_Cold_30pct

Drug Facts (continued)

Active ingredients (in each tablet)

Fexofenadine HCl 180 mg
Antihistamine

Pseudoephedrine HCl 240 mg
Nasal decongestant

Purpose

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

temporarily relieves sinus congestion and pressure

temporarily restores free breathing through the nose

Drug Facts (continued)

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

temporarily restores free breathing through the nose

NDC XXXX-XXXX-XXXX

Allegra-D[®]

ALLERGY & CONGESTION

24 HR

fexofenadine HCl 180 mg/antihistamine
pseudoephedrine HCl 240 mg/nasal decongestant

INDOOR / OUTDOOR RELIEF OF

- ✓ Nasal and Sinus Congestion Due to Colds or Allergies
- ✓ Sneezing; Runny Nose; Itchy, Watery Eyes and Itchy Nose or Throat Due to Allergies

NON-DROWSY

Extended Release

15 TABLETS

Drug Facts (continued)

Questions or comments?

call toll-free 1-800-633-1610 or www.allegra.com

The makers of Allegra[®] do not make store brand products. The trade dress of this Allegra[®] package is subject to trademark protection. Dist. By: Chatham, Inc., a Sanofi Company
Chattanooga, TN 37409-0219 ©2015 Origin France
XXXXXXXXXX

UPC

FPO

Drug Facts (continued)

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing

Ask a doctor before use if you have

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever.
- you get nervous, dizzy, or sleepless

Drug Facts (continued)

Directions

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

If pregnant or breast-feeding, ask a health professional before use.

adults and children 12 years and over

take 1 tablet with a glass of water every 24 hours on an empty stomach; do not take more than 1 tablet in 24 hours

children under 12 years of age

do not use

adults 65 years of age and older

ask a doctor

consumers with kidney disease

ask a doctor

Other information

each tablet contains sodium 33 mg

- safely sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° and 25°C (68° and 77°F)

Inactive ingredients acetone, black iron oxide, cellulose acetate, colloidal silicon dioxide, copovidone, croscarmellose sodium, FD&C blue #1 aluminum lake, glycerol triacetate, hypromellose, isopropyl alcohol, magnesium stearate, methyl alcohol, methylene chloride, microcrystalline cellulose, polyethylene glycol, propylene glycol, povidone, sodium chloride, talc, titanium dioxide, water

Drug Facts (continued)

Directions

do not divide, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years and over

take 1 tablet with a glass of water every 24 hours on an empty stomach; do not take more than 1 tablet in 24 hours

children under 12 years of age

do not use

adults 65 years of age and older

ask a doctor

consumers with kidney disease

ask a doctor

Other information

each tablet contains sodium 33 mg

- safely sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° and 25°C (68° and 77°F)

Inactive ingredients acetone, black iron oxide, cellulose acetate, colloidal silicon dioxide, copovidone, croscarmellose sodium, FD&C blue #1 aluminum lake, glycerol triacetate, hypromellose, isopropyl alcohol, magnesium stearate, methyl alcohol, methylene chloride, microcrystalline cellulose, polyethylene glycol, propylene glycol, povidone, sodium chloride, talc, titanium dioxide, water

Allegra-D[®]

ALLERGY & CONGESTION

24 HR

LOT

EXP

Allegra-D® Allergy and Congestion 24 Hour

05.02.16

ALG-D_24hr_5ct_Blister_2015



This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
06/02/2016