



Rx Only

**WARNING: RESPIRATORY DEPRESSION IN CHILDREN; DEATH RELATED TO ULTRA RAPID METABOLISM OF CODEINE TO MORPHINE and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS**

**Respiratory Depression in Children**

The combination of promethazine hydrochloride and codeine phosphate is contraindicated in pediatric patients less than 6 years of age. Concomitant administration of promethazine products with other respiratory depressants has an association with respiratory depression, and sometimes death, in pediatric patients.

Postmarketing cases of respiratory depression, including fatalities, have been reported with use of promethazine hydrochloride in pediatric patients less than 2 years of age. A wide range of weight-based doses of promethazine hydrochloride have resulted in respiratory depression in these patients.

**Death Related to Ultra-Rapid Metabolism of Codeine to Morphine**

Respiratory depression and death have occurred in children who received codeine following tonsillectomy and/or adenoidectomy and had evidence of being ultra-rapid metabolizers of codeine due to a CYP2D6 polymorphism.

**Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death (see WARNINGS, PRECAUTIONS - Drug Interactions). Avoid use of opioid cough medications in patients taking benzodiazepines, other CNS depressants, or alcohol.

**DESCRIPTION**


Each 5 mL (one teaspoonful), for oral administration contains: Promethazine hydrochloride 6.25 mg; codeine phosphate 10 mg in a flavored syrup base with a pH between 4.8 and 5.4. Alcohol 7%.

Inactive ingredients: Artificial and natural flavors, citric acid, D&C Red 33, FD&C Blue 1, FD&C Yellow 6, glycerin, saccharin sodium, sodium benzoate, sodium citrate, sodium propionate, water, and other ingredients.

Codeine is one of the naturally occurring phenanthrene alkaloids of opium derived from the opium poppy, it is classified pharmacologically as a narcotic analgesic. Codeine phosphate may be chemically designated as 7,8-Didehydro-4, 5 $\alpha$ -epoxy-3-methoxy-17-methylmorphinan-6 $\alpha$ -ol phosphate (1:1) (salt) hemihydrate.


The phosphate salt of codeine occurs as white, needle-shaped crystals or white crystalline powder. Codeine phosphate is freely soluble in water and slightly soluble in alcohol. It has a

molecular weight of 406.37, a molecular formula of  $C_{18}H_{21}NO_3 \cdot H_3PO_4 \cdot \frac{1}{2}H_2O$ , and the following structural formula:



Promethazine hydrochloride, a phenothiazine derivative, is chemically designated as (±)-10-[2-(Dimethylamino)propyl] phenothiazine monohydrochloride.

Promethazine hydrochloride occurs as a white to faint yellow, practically odorless, crystalline powder which slowly oxidizes and turns blue on prolonged exposure to air. It is soluble in water and freely soluble in alcohol. It has a molecular weight of 320.88, a molecular formula of  $C_{17}H_{20}N_2S \cdot HCl$ , and the following structural formula:



## CLINICAL PHARMACOLOGY

### Codeine

Narcotic analgesics, including codeine, exert their primary effects on the central nervous system and gastrointestinal tract. The analgesic effects of codeine are due to its central action; however, the precise sites of action have not been determined, and the mechanisms involved appear to be quite complex. Codeine resembles morphine both structurally and pharmacologically, but its actions at the doses of codeine used therapeutically are milder, with less sedation, respiratory depression, and gastrointestinal, urinary, and pupillary effects. Codeine produces an increase in biliary tract pressure, but less than morphine or meperidine. Codeine is less constipating than morphine. Codeine has good antitussive activity, although less than that of morphine at equal doses. It is used in preference to morphine, because side effects are infrequent at the usual antitussive dose of codeine.

Codeine in oral therapeutic dosage does not usually exert major effects on the cardiovascular system.

Narcotic analgesics may cause nausea and vomiting by stimulating the chemoreceptor trigger zone (CTZ); however, they also depress the vomiting center, so that subsequent doses are unlikely to produce vomiting. Nausea is minimal after usual oral doses of codeine.

Narcotic analgesics cause histamine release, which appears to be responsible for wheals or urticaria sometimes seen at the site of injection on parenteral administration. Histamine release may also produce dilation of cutaneous blood vessels, with resultant flushing of the face and neck, pruritus, and sweating.

Codeine and its salts are well absorbed following both oral and parenteral administration. Codeine is about 2/3 as effective orally as parenterally. Codeine is metabolized primarily in the liver by enzymes of the endoplasmic reticulum, where it undergoes O-demethylation, N-demethylation, and partial conjugation with glucuronic acid. The drug is excreted primarily in the urine, largely as inactive metabolites and small amounts of free and conjugated morphine. Negligible amounts of codeine and its metabolites are found in the feces.

Following oral or subcutaneous administration of codeine, the onset of analgesia occurs within 15 to 30 minutes and lasts for four to six hours.

The cough-depressing action, in animal studies, was observed to occur 15 minutes after oral administration of codeine, peak action at 45 to 60 minutes after ingestion. The duration of action, which is dose-dependent, usually did not exceed 3 hours.

### **Promethazine**

Promethazine is a phenothiazine derivative which differs structurally from the antipsychotic phenothiazines by the presence of a branched side chain and no ring substitution. It is thought that this configuration is responsible for its lack (1/10 that of chlorpromazine) of dopamine antagonist properties.

Promethazine is an H<sub>1</sub> receptor blocking agent. In addition to its antihistaminic action, it provides clinically useful sedative and antiemetic effects.

Promethazine is well absorbed from the gastrointestinal tract. Clinical effects are apparent within 20 minutes after oral administration and generally last four to six hours, although they may persist as long as 12 hours. Promethazine is metabolized by the liver to a variety of compounds; the sulfoxides of promethazine and N-demethylpromethazine are the predominant metabolites appearing in the urine.

## **INDICATIONS AND USAGE**

Promethazine HCl and Codeine Phosphate Oral Solution is indicated for the temporary relief of coughs and upper respiratory symptoms associated with allergy or the common cold.

## **CONTRAINDICATIONS**

The combination of promethazine hydrochloride and codeine phosphate is contraindicated in pediatric patients less than 6 years of age, because the combination may cause fatal respiratory depression in this age population.

Codeine sulfate is contraindicated for post-operative pain management in children who have undergone tonsillectomy and/or adenoidectomy. (See **WARNINGS - Death Related to Ultra-Rapid Metabolism of Codeine to Morphine**).

Codeine is contraindicated in patients with a known hypersensitivity to the drug.

Promethazine hydrochloride is contraindicated in comatose states, and in individuals known to be hypersensitive or to have had an idiosyncratic reaction to promethazine or to other phenothiazines.

Antihistamines and codeine are both contraindicated for use in the treatment of lower respiratory tract symptoms, including asthma.

## **WARNINGS**

(see **Boxed Warnings**)

### **Respiratory Depression in Children**

The combination of promethazine hydrochloride and codeine phosphate is contraindicated in pediatric patients less than 6 years of age. Concomitant administration of promethazine products with other respiratory depressants has an association with respiratory depression, and sometimes death, in pediatric patients.

Postmarketing cases of respiratory depression, including fatalities, have been reported with use of promethazine hydrochloride in pediatric patients less than 2 years of age. A wide range of weight-based doses of promethazine hydrochloride have resulted in respiratory depression in these patients.

Respiratory depression leading to arrest, coma, and death has occurred with the use of codeine antitussives in young children, particularly in the under-one-year infants whose ability to deactivate the drug is not fully developed.

### **Codeine**

- **Death Related to Ultra-Rapid Metabolism of Codeine to Morphine**

Respiratory depression and death have occurred in children who received codeine in the post-operative period following tonsillectomy and/or adenoidectomy and had evidence of being ultra-rapid metabolizers of codeine (i.e., multiple copies of the gene for cytochrome P450 isoenzyme 2D6 [CYP2D6] or high morphine concentrations). Deaths have also occurred in nursing infants who were exposed to high levels of morphine in breast milk because their mothers were ultra-rapid metabolizers of codeine. (See **PRECAUTIONS -Nursing Mothers**).

Some individuals may be ultra-rapid metabolizers because of a specific CYP2D6 genotype (gene duplications denoted as \*1/\*1xN or \*1/\*2xN). The prevalence of this CYP2D6 phenotype varies widely and has been estimated at 0.5 to 1% in Chinese and Japanese, 0.5 to

1% in Hispanics, 1 to 10% in Caucasians, 3% in African Americans, and 16 to 28% in North Africans, Ethiopians, and Arabs. Data are not available for other ethnic groups. These individuals convert codeine into its active metabolite, morphine, more rapidly and completely than other people. This rapid conversion results in higher than expected serum morphine levels. Even at labeled dosage regimens, individuals who are ultra-rapid metabolizers may have life-threatening or fatal respiratory depression or experience signs of overdose (such as extreme sleepiness, confusion, or shallow breathing). (See **OVERDOSAGE**).

Children with obstructive sleep apnea who are treated with codeine for post-tonsillectomy and/or adenoidectomy pain may be particularly sensitive to the respiratory depressant effects of codeine that has been rapidly metabolized to morphine. Codeine is contraindicated for post-operative pain management in all pediatric patients undergoing tonsillectomy and/or adenoidectomy. (See **CONTRAINDICATIONS**).

When prescribing codeine-containing drugs, healthcare providers should choose the lowest effective dose for the shortest period of time and inform patients and caregivers about these risks and the signs of morphine overdose. (See **OVERDOSAGE**).

- **Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**  
Concomitant use of opioids, including Promethazine HCl and Codeine Phosphate Oral Solution, with benzodiazepines, or other CNS depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Because of these risks, avoid use of opioid cough medications in patients taking benzodiazepines, other CNS depressants, or alcohol. (See **PRECAUTIONS - Drug Interactions**).

Observational studies have demonstrated that concomitant use of opioid analgesics and benzodiazepines increases the risk of drug-related mortality compared to use of opioids alone. Because of similar pharmacologic properties, it is reasonable to expect similar risk with concomitant use of opioid cough medications and benzodiazepines, other CNS depressants, or alcohol.

Advise both patients and caregivers about the risks of respiratory depression and sedation if Promethazine HCl and Codeine Phosphate Oral Solution is used with benzodiazepines, alcohol, or other CNS depressants. (See **PRECAUTIONS – Information for Patients**).

- Dosage of codeine **SHOULD NOT BE INCREASED** if cough fails to respond; an unresponsive cough should be reevaluated in 5 days or sooner for possible underlying pathology, such as foreign body or lower respiratory tract disease.
- Codeine may cause or aggravate constipation.
- Administration of codeine may be accomplished by histamine release and should be used with caution in atopic children.
- **Head Injury and Increased Intracranial Pressure**  
The respiratory depressant effects of narcotic analgesics and their capacity to elevate cerebrospinal fluid pressure may be markedly exaggerated in the presence of head injury, intracranial lesions, or a preexisting increase in intracranial pressure. Narcotics may produce adverse reactions which may obscure the clinical course of patients with head injuries.

- **Asthma and Other Respiratory Conditions**

Narcotic analgesics or cough suppressants, including codeine, should not be used in asthmatic patients (see **CONTRAINDICATIONS**). Nor should they be used in acute febrile illness associated with productive cough or in chronic respiratory disease where interference with ability to clear the tracheobronchial tree of secretions would have a deleterious effect on the patient's respiratory function.

- **Hypotensive Effect**

Codeine may produce orthostatic hypotension in ambulatory patients.

## **Promethazine**

- **CNS Depression**

Promethazine may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks, such as driving a vehicle or operating machinery. The impairment may be amplified by concomitant use of other central-nervous-system depressants such as alcohol, sedatives/hypnotics (including barbiturates), narcotics, narcotic analgesics, general anesthetics, tricyclic antidepressants, and tranquilizers; therefore avoid use of Promethazine HCl and Codeine Phosphate Oral Solution in patients on these medications. (See **PRECAUTIONS – Information for Patients and Drug Interactions**).

- **Respiratory Depression**

Promethazine may lead to potentially fatal respiratory depression.

Use of Promethazine in patients with compromised respiratory function (e.g. COPD, sleep apnea) should be avoided.

- **Lower Seizure Threshold**

Promethazine may lower seizure threshold. It should be used with caution in persons with seizure disorder or in persons who are using concomitant medications, such as narcotics or local anesthetics, which may also affect seizure threshold.

- **Bone-Marrow Depression**

Promethazine should be used with caution in patients with bone-marrow depression. Leukopenia and agranulocytosis have been reported, usually when promethazine HCl has been used in association with other known marrow-toxic agents.

- **Neuroleptic Malignant Syndrome**

A potentially fatal symptom complex sometimes referred to as Neuroleptic Malignant Syndrome (NMS) has been reported in association with promethazine HCl alone or in combination with antipsychotic drugs. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis and cardiac dysrhythmias).

The diagnostic evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to identify cases where the clinical presentation includes both serious medical illness (e.g., pneumonia, systemic infection, etc.) and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the

differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever and primary central nervous system (CNS) pathology.

The management of NMS should include 1) immediate discontinuation of promethazine HCl, antipsychotic drugs, if any, and other drugs not essential to concurrent therapy, 2) intensive symptomatic treatment and medical monitoring, and 3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for uncomplicated NMS.

Since recurrences of NMS have been reported with phenothiazines, the reintroduction of promethazine HCl should be carefully considered.

### **Use in Pediatric Patients**

**The combination of promethazine hydrochloride and codeine phosphate is contraindicated in pediatric patients less than 6 years of age. Concomitant administration of promethazine products with other respiratory depressants has an association with respiratory depression, and sometimes death, in pediatric patients. The association does not directly relate to individualized weight-based dosing, which might otherwise permit safe administration.**

Respiratory depression and death have occurred in children with obstructive sleep apnea who received codeine in the post-operative period following tonsillectomy and/or adenoidectomy and had evidence of being ultra-rapid metabolizers of codeine (i.e., multiple copies of the gene for CYP2D6 or high morphine concentrations). These children may be particularly sensitive to the respiratory depressant effects of codeine that has been rapidly metabolized to morphine. Codeine is contraindicated for post-operative pain management in these patients. (See **WARNINGS - Death Related to Ultra-Rapid Metabolism of Codeine to Morphine and CONTRAINDICATIONS**).

Excessively large dosages of antihistamines, including promethazine hydrochloride, in pediatric patients may cause sudden death (see **OVERDOSAGE**). Hallucinations and convulsions have occurred with therapeutic doses and overdoses of promethazine hydrochloride in pediatric patients. In pediatric patients who are acutely ill associated with dehydration, there is an increased susceptibility to dystonias with the use of promethazine HCl.

### **Other Considerations**

Administration of promethazine has been associated with reported cholestatic jaundice.

## **PRECAUTIONS**

### **General**

Narcotic analgesics, including codeine, should be administered with caution and the initial dose reduced in patients with acute abdominal conditions, convulsive disorders, significant hepatic or renal impairment, fever, hypothyroidism, Addison's disease, ulcerative colitis, prostatic hypertrophy, in patients with recent gastrointestinal or urinary tract surgery, and in the very young or elderly or debilitated patients.

Drugs having anticholinergic properties should be used with caution in patients with narrow-angle glaucoma, prostatic hypertrophy, stenosing peptic ulcer, pyloroduodenal obstruction, and bladder-neck obstruction.

Promethazine should be used cautiously in persons with cardiovascular disease or with impairment of liver function.

### **Information for Patients**

Patients should be advised to measure Promethazine HCl and Codeine Phosphate Oral Solution with an accurate measuring device. A household teaspoon is not an accurate measuring device and could lead to overdose, especially when a half a teaspoon is measured. A pharmacist can recommend an appropriate measuring device and can provide instructions for measuring the correct dose.

Promethazine and codeine may cause marked drowsiness or may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks, such as driving a vehicle or operating machinery. Ambulatory patients should be told to avoid engaging in such activities until it is known that they do not become drowsy or dizzy from promethazine and codeine therapy. Pediatric patients should be supervised to avoid potential harm in bike riding or in other hazardous activities.

Inform patients and caregivers that potentially fatal additive effects may occur if Promethazine HCl and Codeine Phosphate Oral Solution is used with benzodiazepines or other CNS depressants, including alcohol. Because of this risk, patients should avoid concomitant use of Promethazine HCl and Codeine Phosphate Oral Solution with benzodiazepines or other CNS depressants, including alcohol. (See **WARNINGS - Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**).

Patients should be advised to report any involuntary muscle movements.

Avoid prolonged exposure to the sun.

Codeine, like other narcotic analgesics, may produce orthostatic hypotension in some ambulatory patients. Patients should be cautioned accordingly.

Advise patients that some people have a genetic variation that results in codeine changing into morphine more rapidly and completely than other people. Most people are unaware of whether they are an ultra-rapid codeine metabolizer or not. These higher-than-normal levels of morphine in the blood may lead to life-threatening or fatal respiratory depression or signs of overdose such as extreme sleepiness, confusion, or shallow breathing. Children with this genetic variation who were prescribed codeine after tonsillectomy and/or adenoidectomy for obstructive sleep apnea may be at greatest risk based on reports of several deaths in this population due to respiratory depression. As a result, codeine is contraindicated in children who undergo tonsillectomy and/or adenoidectomy. Advise caregivers of children receiving codeine for other reasons to monitor for signs of respiratory depression. (See **WARNINGS - Death Related to Ultra-Rapid Metabolism of Codeine to Morphine**).

Nursing mothers taking codeine can also have higher morphine levels in their breast milk if they are ultra-rapid metabolizers. These higher levels of morphine in breast milk may lead to life-

threatening or fatal side effects in nursing babies. Instruct nursing mothers to watch for signs of morphine toxicity in their infants including increased sleepiness (more than usual), difficulty breastfeeding, breathing difficulties, or limpness. Instruct nursing mothers to talk to the baby's doctor immediately if they notice these signs and, if they can not reach the doctor right away, to take the baby to an emergency room or call 911 (or local emergency services).

## **Drug Interactions**

### **Codeine**

In patients receiving MAO inhibitors, an initial small test dose is advisable to allow observation of any excessive narcotic effects or MAOI interaction.

The use of benzodiazepines, opioids, antihistamines, antipsychotics, anti-anxiety agents, or other CNS depressants (including alcohol) concomitantly with Promethazine HCl and Codeine Phosphate Oral Solution may cause an additive CNS depressant effect, profound sedation, respiratory depression, coma, and death and should be avoided. (See **WARNINGS - Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**).

### **Promethazine**

*Epinephrine:* Because of the potential for promethazine to reverse epinephrine's vasopressor effect, epinephrine should NOT be used to treat hypotension associated with promethazine overdose.

*Anticholinergics:* Concomitant use of other agents with anticholinergic properties should be undertaken with caution.

*Monoamine Oxidase Inhibitors (MAOI):* Drug interactions, including an increased incidence of extrapyramidal effects, have been reported when some MAOI and phenothiazines are used concomitantly.

### **Drug/Laboratory Test Interactions**

Because narcotic analgesics may increase biliary tract pressure, with resultant increase in plasma amylase or lipase levels, determination of these enzyme levels may be unreliable for 24 hours after a narcotic analgesic has been given.

The following laboratory tests may be affected in patients who are receiving therapy with promethazine hydrochloride:

*Pregnancy Tests:* Diagnostic pregnancy tests based on immunological reactions between HCG and anti-HCG may result in false-negative or false-positive interpretations.

*Glucose Tolerance Test:* An increase in blood glucose has been reported in patients receiving promethazine.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term animal studies have not been performed to assess the carcinogenic potential of codeine or of promethazine, nor are there other animal or human data concerning carcinogenicity, mutagenicity, or impairment of fertility with these agents. Codeine has been reported to show no evidence of carcinogenicity or mutagenicity in a variety of test systems,

including the micronucleus and sperm abnormality assays and the *Salmonella* assay. Promethazine was nonmutagenic in the *Salmonella* test system of Ames.

## **Pregnancy**

### **Teratogenic Effects**

*Pregnancy Category C:*

Codeine: A study in rats and rabbits reported no teratogenic effect of codeine administered during the period of organogenesis in doses ranging from 5 to 120 mg/kg. In the rat, doses at the 120-mg/kg level, in the toxic range for the adult animal were associated with an increase in embryo resorption at the time of implantation. In another study a single 100-mg/kg dose of codeine administered to pregnant mice reportedly resulted in delayed ossification in the offspring.

There are no studies in humans, and the significance of these findings to humans, if any, is not known.

Promethazine: Teratogenic effects have not been demonstrated in rat-feeding studies at doses of 6.25 and 12.5 mg/kg of promethazine HCl. These doses are from approximately 2.1 to 4.2 times the maximum recommended total daily dose of promethazine for a 50-kg subject depending upon the indication for which the drug is prescribed. Daily doses of 25 mg/kg intraperitoneally have been found to produce fetal mortality in rats.

Specific studies to test the action of the drug on parturition, lactation, and development of the animal neonate were not done, but a general preliminary study in rats indicated no effect on these parameters. Although antihistamines have been found to produce fetal mortality in rodents, the pharmacological effects of histamine in the rodent do not parallel those in man. There are no adequate and well-controlled studies of promethazine in pregnant women.

Animal reproduction studies have not been conducted with the drug combination – promethazine and codeine. It is not known whether this drug combination can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Promethazine HCl and Codeine Phosphate Oral Solution should be given to a pregnant woman only if clearly needed.

Promethazine HCl and Codeine Phosphate Oral Solution should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

### **Nonteratogenic Effects**

Dependence has been reported in newborns whose mothers took opiates regularly during pregnancy. Withdrawal signs include irritability, excessive crying, tremors, hyperreflexia, fever, vomiting, and diarrhea. Signs usually appear during the first few days of life.

Promethazine administered to a pregnant woman within two weeks of delivery may inhibit platelet aggregation in the newborn.

### **Labor and Delivery**

Narcotic analgesics cross the placental barrier. The closer to the delivery and the larger the dose used, the greater the possibility of respiratory depression in the newborn. Narcotic analgesics should be avoided during labor if delivery of a premature infant is anticipated. If the mother has received narcotic analgesics during labor, newborn infants should be observed closely for signs

of respiratory depression. Resuscitation may be required (see **OVERDOSAGE**). Limited data suggests that use of promethazine hydrochloride during labor and delivery does not have an appreciable effect on the duration of labor or delivery and does not increase the risk of need for intervention in the newborn.

The effect of promethazine and/or codeine on later growth and development of the newborn is unknown.

### **Nursing Mothers**

Codeine is secreted into human milk. In women with normal codeine metabolism (normal CYP2D6 activity), the amount of codeine excreted into human milk is low and dose-dependent. Despite the common use of codeine products to manage postpartum pain, reports of adverse events in infants are rare. However, some women are ultra-rapid metabolizers of codeine. These women achieve higher-than-expected serum levels of codeine's active metabolite, morphine, leading to higher-than-expected levels of morphine in breast milk and potentially dangerously high serum morphine levels in their breastfed infants. Therefore, maternal use of codeine can potentially lead to serious adverse reactions, including death, in nursing infants.

The risk of infant exposure to codeine and morphine through breast milk should be weighed against the benefits of breastfeeding for both the mother and baby. Caution should be exercised when codeine is administered to a nursing woman. If a codeine containing product is selected, the lowest dose should be prescribed for the shortest period of time to achieve the desired clinical effect. Mothers using codeine should be informed about when to seek immediate medical care and how to identify the signs and symptoms of neonatal toxicity, such as drowsiness or sedation, difficulty breastfeeding, breathing difficulties, and decreased tone, in their baby. Nursing mothers who are ultra-rapid metabolizers may also experience overdose symptoms such as extreme sleepiness, confusion or shallow breathing. Prescribers should closely monitor mother-infant pairs and notify treating pediatricians about the use of codeine during breastfeeding. (See **WARNINGS - Death Related to Ultra-Rapid Metabolism of Codeine to Morphine.**)

### **Pediatric Use**

**The combination of promethazine hydrochloride and codeine phosphate is contraindicated in pediatric patients less than 6 years of age, because the combination may cause fatal respiratory depression in this age population (see WARNINGS – Boxed Warning and Use in Pediatric Patients).**

Respiratory depression and death have occurred in children with obstructive sleep apnea who received codeine in the post-operative period following tonsillectomy and/or adenoidectomy and had evidence of being ultra-rapid metabolizers of codeine (i.e., multiple copies of the gene for CYP2D6 or high morphine concentrations). These children may be particularly sensitive to the respiratory depressant effects of codeine that has been rapidly metabolized to morphine. Codeine is contraindicated for post-operative pain management in all pediatric patients undergoing tonsillectomy and/or adenoidectomy. (See **WARNINGS - Death Related to Ultra-Rapid Metabolism of Codeine to Morphine** and **CONTRAINDICATIONS**).

### **Geriatric Use**

Clinical studies of Promethazine HCl and Codeine Phosphate Oral Solution did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in

responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

Sedating drugs may cause confusion and over-sedation in the elderly; elderly patients generally should be started on low doses of Promethazine Hydrochloride and Codeine Phosphate Oral Solution and observed closely.

## **ADVERSE REACTIONS**

### **Codeine**

*Central Nervous System:* CNS depression, particularly respiratory depression, and to a lesser extent circulatory depression; light-headedness, dizziness, sedation, euphoria, dysphoria, headache, transient hallucination, disorientation, visual disturbances, and convulsions.

*Cardiovascular:* Tachycardia, bradycardia, palpitation, faintness, syncope, orthostatic hypotension (common to narcotic analgesics).

*Gastrointestinal:* Nausea, vomiting, constipation, and biliary tract spasm. Patients with chronic ulcerative colitis may experience increased colonic motility; in patients with acute ulcerative colitis, toxic dilation has been reported.

*Genitourinary:* Oliguria, urinary retention, antidiuretic effect has been reported (common to narcotic analgesics).

*Allergic:* Infrequent pruritus, giant urticaria, angioneurotic edema, and laryngeal edema.

*Other:* Flushing of the face, sweating and pruritus (due to opiate-induced histamine release); weakness.

### **Promethazine**

*Central Nervous System:* Drowsiness is the most prominent CNS effect of this drug. Sedation, somnolence, blurred vision, dizziness, confusion, disorientation and extrapyramidal symptoms such as oculogyric crisis, torticollis, and tongue protrusion; lassitude, tinnitus, incoordination, fatigue, euphoria, nervousness, diplopia, insomnia, tremors, convulsive seizures, excitation, catatonic-like states, hysteria. Hallucinations have also been reported.

*Cardiovascular:* Increased or decreased blood pressure, tachycardia, bradycardia, faintness.

*Dermatologic:* Dermatitis, photosensitivity, urticaria.

*Hematologic:* Leukopenia, thrombocytopenia, thrombocytopenic purpura, agranulocytosis.

*Gastrointestinal:* Dry mouth, nausea, vomiting, jaundice.

*Respiratory:* Asthma, nasal stuffiness, respiratory depression (potentially fatal) and apnea (potentially fatal) (see **WARNINGS – Promethazine; Respiratory Depression**).

*Other:* Angioneurotic edema. Neuroleptic malignant syndrome (potentially fatal) has also been reported (see **WARNINGS – Promethazine; Neuroleptic Malignant Syndrome**).

*Paradoxical Reactions:* Hyperexcitability and abnormal movements have been reported in patients following a single administration of promethazine HCl. Consideration should be given to the discontinuation of promethazine HCl and to the use of other drugs if these reactions occur. Respiratory depression, nightmares, delirium and agitated behavior have also been reported in some of these patients.

## **DRUG ABUSE AND DEPENDENCE**

### **Controlled Substance**

Promethazine HCl and Codeine Phosphate Oral Solution is a Schedule V Controlled Substance.

### **Abuse**

Codeine is known to be subject to abuse; however, the abuse potential of oral codeine appears to be quite low. Even parenteral codeine does not appear to offer the psychic effects sought by addicts to the same degree as heroin or morphine. However, codeine must be administered only under close supervision to patients with a history of drug abuse or dependence.

### **Dependence**

Psychological dependence, physical dependence, and tolerance are known to occur with codeine.

## **OVERDOSAGE**

### **Codeine**

Serious overdose with codeine is characterized by respiratory depression (a decrease in respiratory rate and/or tidal volume, Cheyne-stokes respiration, cyanosis), extreme somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, and sometimes bradycardia and hypotension. The triad of coma, pinpoint pupils, and respiratory depression is strongly suggestive of opiate poisoning. In severe overdose, particularly by the intravenous route, apnea, circulatory collapse, cardiac arrest, and death may occur. Promethazine is additive to the depressant effects of codeine.

It is difficult to determine what constitutes a standard toxic or lethal dose. However, the lethal oral dose of codeine in an adult is reported to be in the range of 0.5 to 1 gram. Infants and children are believed to be relatively more sensitive to opiates on a body-weight basis. Elderly patients are also comparatively intolerant to opiates.

### **Promethazine**

Signs and symptoms of overdose with promethazine range from mild depression of the central nervous system and cardiovascular system to profound hypotension, respiratory depression, unconsciousness and sudden death. Other reported reactions include hyperreflexia, hypertonia, ataxia, athetosis and extensor-plantar reflexes (Babinski reflex).

Stimulation may be evident, especially in children and geriatric patients. Convulsions may rarely occur. A paradoxical reaction has been reported in children receiving single doses of 75 mg to 125 mg orally, characterized by hyperexcitability and nightmares.

Atropine-like signs and symptoms – dry mouth, fixed dilated pupils, flushing, as well as gastrointestinal symptoms, may occur.

### **Treatment**

Treatment of overdosage with promethazine and codeine is essentially symptomatic and supportive. Only in cases of extreme overdosage or individual sensitivity do vital signs including respiration, pulse, blood pressure, temperature, and EKG need to be monitored. Activated charcoal orally or by lavage may be given, or sodium or magnesium sulfate orally as a cathartic. Attention should be given to the reestablishment of adequate respiratory exchange through provision of a patent airway and institution of assisted or controlled ventilation. The narcotic antagonist, naloxone hydrochloride, may be administered when significant respiratory depression occurs with promethazine and codeine; any depressant effects of promethazine are not reversed with naloxone. Diazepam may be used to control convulsions. Avoid analeptics, which may cause convulsions. Acidosis and electrolyte losses should be corrected. A rise in temperature or pulmonary complications may signal the need for institution of antibiotic therapy.

Severe hypotension usually responds to the administration of norepinephrine or phenylephrine. EPINEPHRINE SHOULD NOT BE USED, since its use in a patient with partial adrenergic blockade may further lower the blood pressure.

Limited experience with dialysis indicates that it is not helpful.

## **DOSAGE AND ADMINISTRATION**

*It is important that Promethazine HCl and Codeine Phosphate Oral Solution is measured with an accurate measuring device (see **PRECAUTIONS-Information for Patients**). A household teaspoon is not an accurate measuring device and could lead to overdosage, especially when half a teaspoon is to be measured. It is strongly recommended that an accurate measuring device be used. A pharmacist can provide an appropriate device and can provide instructions for measuring the correct dose.*

**The combination of promethazine hydrochloride and codeine phosphate is contraindicated in pediatric patients less than 6 years of age, because the combination may cause fatal respiratory depression in this age population.**

The average effective dose is given in the following table:

<b>Adults (12 years of age and over)</b>	5 mL (1 teaspoonful) every 4 to 6 hours, not to exceed 30.0 mL in 24 hours.
<b>Children 6 years to under 12 years</b>	2.5 mL to 5 mL (½ to 1 teaspoonful) every 4 to 6 hours, not to exceed 30.0 mL in 24 hours.

## **HOW SUPPLIED**

Promethazine HCl and Codeine Phosphate Oral Solution is a clear, purple solution supplied in bottles of 4 fl. oz. (118 mL) and 16 fl. oz. (473 mL).

**Keep bottles tightly closed.**

**Store at 20° to 25° C (68° to 77° F) [See USP Controlled Room Temperature.]**

**Protect from light.**

**Dispense in tight, light-resistant container (USP/NF) with a child-resistant closure.**

**DEA Order Form Required**

Manufactured for:  
ANI Pharmaceuticals, Inc.  
Baudette, MN 56623



9576 Rev 01/17

## MEDICATION GUIDE

### Promethazine Hydrochloride and Codeine Phosphate Oral Solution, C-V

#### What is the most important information I should know about Promethazine Hydrochloride and Codeine Phosphate Oral Solution?

- Taking Promethazine Hydrochloride and Codeine Phosphate Oral Solution with benzodiazepines, or other central nervous system depressants, including alcohol can cause severe drowsiness, breathing problems (respiratory depression), coma, and death.
- Promethazine Hydrochloride and Codeine Phosphate Oral Solution can cause you to be drowsy. Avoid driving a car or operating machinery during treatment with Promethazine Hydrochloride and Codeine Phosphate Oral Solution.
- **Do not** give Promethazine Hydrochloride and Codeine Phosphate Oral Solution to children who are under 6 years of age. It may harm them.
- **Do not** give Promethazine Hydrochloride and Codeine Phosphate Oral Solution to a child to treat pain after tonsillectomy or adenoidectomy surgery.
  - When you take Promethazine Hydrochloride and Codeine Phosphate Oral Solution, some of it changes into morphine in your body.
  - In some children and adults this happens very quickly, and can cause you to stop breathing and cause death due to an overdose.
- Women who breastfeed should talk to their healthcare provider before taking Promethazine Hydrochloride and Codeine Phosphate Oral Solution.
  - When you take Promethazine Hydrochloride and Codeine Phosphate Oral Solution, some of it changes into morphine in your body.
  - In some women, this happens very quickly. Codeine and morphine pass into your breast milk. A large amount of morphine can cause your baby to die.
- Call your healthcare provider or get emergency medical help right away if anyone taking Promethazine Hydrochloride and Codeine Phosphate Oral Solution, or your breastfeeding baby has any of the symptoms below:
  - increased sleepiness
  - confusion
  - difficulty breathing
  - shallow breathing
  - limpness
  - your baby has difficulty breastfeeding
- Keep Promethazine Hydrochloride and Codeine Phosphate Oral Solution in a safe place away from children. Accidental use by a child is a medical emergency and can cause death. If a child accidentally takes Promethazine Hydrochloride and Codeine Phosphate Oral Solution, get emergency medical help right away.
- Promethazine Hydrochloride and Codeine Phosphate Oral Solution can cause serious side effects including death.
- Take Promethazine Hydrochloride and Codeine Phosphate Oral Solution exactly as prescribed by your healthcare provider. If you take the wrong dose of Promethazine Hydrochloride and Codeine Phosphate Oral Solution, you could overdose and die.
- It is especially important when you take Promethazine Hydrochloride and Codeine Phosphate Oral Solution that you know exactly what dose to take.

#### What is Promethazine Hydrochloride and Codeine Phosphate Oral Solution?

- Promethazine Hydrochloride and Codeine Phosphate Oral Solution is a prescription medicine used to temporarily treat cough and upper respiratory symptoms that you can have with allergies or a common cold. Promethazine Hydrochloride and Codeine Phosphate Oral Solution contains 2 medicines, promethazine and codeine. Promethazine is an antihistamine. Codeine is a narcotic cough suppressant.
- **Promethazine Hydrochloride and Codeine Phosphate Oral Solution is a federal controlled substance (C-V) because it contains codeine that can be abused or lead to dependence.** Keep Promethazine Hydrochloride and Codeine Phosphate Oral Solution in a safe place to prevent misuse and abuse. Selling or giving away Promethazine Hydrochloride and Codeine Phosphate Oral Solution may harm others, and is against the law. Tell your healthcare provider if you have ever abused or been dependent on alcohol, prescription medicines or street drugs.
- Promethazine Hydrochloride and Codeine Phosphate Oral Solution is not for children under 6 years of age.

#### Who should not take Promethazine Hydrochloride and Codeine Phosphate Oral Solution?

- **Do not** give Promethazine Hydrochloride and Codeine Phosphate Oral Solution to children who are under 6 years of age.
- **Do not** give Promethazine Hydrochloride and Codeine Phosphate Oral Solution to a child to treat pain after tonsillectomy or adenoidectomy surgery.
- **Do not** take Promethazine Hydrochloride and Codeine Phosphate Oral Solution if you are allergic to any of the ingredients in Promethazine Hydrochloride and Codeine Phosphate Oral Solution. See the end of this Medication Guide for a complete list of ingredients in Promethazine Hydrochloride and Codeine Phosphate Oral Solution.
- **Do not** take Promethazine Hydrochloride and Codeine Phosphate Oral Solution if you have asthma.

#### Before you take Promethazine Hydrochloride and Codeine Phosphate Oral Solution, tell your healthcare provider

**about all of your medical conditions, including if you:**

- have a drug dependence
- have lung or breathing problems including chronic obstructive pulmonary disease (COPD), or sleep apnea
- have had a head injury
- have pain in your stomach-area (abdomen)
- have a history of severe or persistent cough
- have prostate problems
- have problems with your urinary tract (urethral stricture) or difficulty urinating
- have had surgery recently or plan to have surgery
- are pregnant or plan to become pregnant. It is not known if Promethazine Hydrochloride and Codeine Phosphate Oral Solution will harm your unborn baby. You and your healthcare provider should decide if you should take Promethazine Hydrochloride and Codeine Phosphate Oral Solution while you are pregnant.
- are breastfeeding or plan to breastfeed. Codeine Phosphate passes into your breast milk and may harm your baby. You and your healthcare provider should discuss whether you should take Promethazine Hydrochloride and Codeine Phosphate Oral Solution or breastfeed. You should not do both.
- drink alcohol
- have kidney or liver problems
- have thyroid problems including hypothyroidism
- have Addison's disease
- have glaucoma (increased pressure in eyes)
- have diabetes
- have seizures
- have heart problems
- have blood disorders including low white blood cells

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Taking Promethazine Hydrochloride and Codeine Phosphate Oral Solution with certain other medicines can cause side effects or affect how well Promethazine Hydrochloride and Codeine Phosphate Oral Solution or the other medicines work. Do not start or stop other medicines without talking to your healthcare provider.

Especially tell your healthcare provider if you:

- take pain medicines such as narcotics
- take cold or allergy medicines that contain antihistamines or cough suppressants
- take medicines for mental illness (anti-psychotics, anti-anxiety)
- drink alcohol
- take medicines for depression, including monoamine oxidase inhibitors (MAOIs)
- take medicines for stomach or intestinal problems

Ask your healthcare provider if you are not sure if you take one of these medicines.

**How should I take Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**

- Take Promethazine Hydrochloride and Codeine Phosphate Oral Solution exactly as your healthcare provider tells you to take it.
- Your healthcare provider will tell you how much Promethazine Hydrochloride and Codeine Phosphate Oral Solution to take and when to take it. Do not change your dose without talking to your healthcare provider.
- Ask your pharmacist to give you a measuring device to help you measure the correct amount of Promethazine Hydrochloride and Codeine Phosphate Oral Solution. Do not use a household teaspoon to measure your medicine. You may accidentally take too much.
- **Do not** take more than 30 mL of Promethazine Hydrochloride and Codeine Phosphate Oral Solution in 24 hours.
- If you take too much Promethazine Hydrochloride and Codeine Phosphate Oral Solution, call your healthcare provider or go to the nearest hospital emergency room right away.

**What should I avoid while taking Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**

- Promethazine Hydrochloride and Codeine Phosphate Oral Solution can cause you to be drowsy. Avoid driving a car or operating machinery during treatment with Promethazine Hydrochloride and Codeine Phosphate Oral Solution. Also, watch your child closely when riding a bike.
- Avoid drinking alcohol during treatment with Promethazine Hydrochloride and Codeine Phosphate Oral Solution. Drinking alcohol can increase your chances of having serious side effects.
- Avoid spending a long time in sunlight.

**What are the possible side effects of Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**

**Promethazine Hydrochloride and Codeine Phosphate Oral Solution may cause serious side effects, including:**

- **See "What is the most important information I should know about Promethazine Hydrochloride and Codeine Phosphate Oral Solution?"**
- **Breathing problems (respiratory depression)** which can lead to death. Call your healthcare provider or get emergency treatment right away if you have excessive sleepiness, shallow or slow breathing, or confusion.
- **Physical dependence or abuse.** Take Promethazine Hydrochloride and Codeine Phosphate Oral Solution exactly as your healthcare provider tells you to take it. Stopping Promethazine Hydrochloride and Codeine Phosphate Oral Solution suddenly could cause withdrawal symptoms.
- **Bowel problems** including constipation or stomach pain.

- **Increased intracranial pressure.**
- **Feeling dizzy, faint or light-headed, especially when you stand up (orthostatic hypotension).** Light-headedness or fainting may happen when getting up too quickly from a sitting or lying position.
- **Neuroleptic malignant syndrome (NMS) which can lead to death.** Tell your healthcare provider right away if you have any of the following symptoms of NMS:
  - high fever
  - confusion
  - changes in pulse, heart rate, and blood pressure
  - stiff muscles
  - sweating

The most common side effects of Promethazine Hydrochloride and Codeine Phosphate Oral Solution include:

- sleepiness
- nausea and vomiting
- high blood pressure
- confusion
- difficulty urinating
- restlessness
- dryness of mouth, nose, throat
- trouble breathing
- fast heart beat

These are not all the possible side effects of Promethazine Hydrochloride and Codeine Phosphate Oral Solution.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

#### **How should I store Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**

- Store Promethazine Hydrochloride and Codeine Phosphate Oral Solution at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep Promethazine Hydrochloride and Codeine Phosphate Oral Solution in a tightly closed, child-resistant container and out of the light.
- Safely throw away medicine that is out of date or no longer needed.
- **Keep Promethazine Hydrochloride and Codeine Phosphate Oral Solution and all medicines out of the reach of children.**

#### **General information about the safe and effective use of Promethazine Hydrochloride and Codeine Phosphate Oral Solution.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Promethazine Hydrochloride and Codeine Phosphate Oral Solution for a condition for which it was not prescribed. Do not give Promethazine Hydrochloride and Codeine Phosphate Oral Solution to other people, even if they have the same symptoms that you have. It may harm them.

You can ask your pharmacist or healthcare provider for information about Promethazine Hydrochloride and Codeine Phosphate Oral Solution that is written for health professionals.

#### **What are the ingredients in Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**

**Active ingredients:** promethazine hydrochloride and codeine phosphate

**Inactive ingredients:** artificial and natural flavors, citric acid, D&C Red 33, FD&C Blue 1, FD&C Yellow 6, glycerin, saccharin sodium, sodium benzoate, sodium citrate, sodium propionate, water, and other ingredients.

Manufactured for: ANI Pharmaceuticals, Inc., Baudette, MN 56623

For more information, go to [www.anipharmaceuticals.com](http://www.anipharmaceuticals.com) or call 1-800-308-6755.



This Medication Guide has been approved by the U.S. Food and Drug Administration

Issued: January 2017



Rx Only

**WARNING: RESPIRATORY DEPRESSION IN CHILDREN; DEATH RELATED TO ULTRA RAPID METABOLISM OF CODEINE TO MORPHINE and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS**

**Respiratory Depression in Children**

The combination of promethazine hydrochloride and codeine phosphate is contraindicated in pediatric patients less than 6 years of age. Concomitant administration of promethazine products with other respiratory depressants has an association with respiratory depression, and sometimes death, in pediatric patients.

Postmarketing cases of respiratory depression, including fatalities, have been reported with use of promethazine hydrochloride in pediatric patients less than 2 years of age. A wide range of weight-based doses of promethazine hydrochloride have resulted in respiratory depression in these patients.

**Death Related to Ultra-Rapid Metabolism of Codeine to Morphine**

Respiratory depression and death have occurred in children who received codeine following tonsillectomy and/or adenoidectomy and had evidence of being ultra-rapid metabolizers of codeine due to a CYP2D6 polymorphism.

**Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death (see WARNINGS, PRECAUTIONS - Drug Interactions). Avoid use of opioid cough medications in patients taking benzodiazepines, other CNS depressants, or alcohol.


**DESCRIPTION**

Each 5 mL (one teaspoonful), for oral administration contains: Codeine phosphate 10 mg; promethazine hydrochloride 6.25 mg; phenylephrine hydrochloride 5 mg in a flavored syrup base with a pH between 4.8 and 5.4. Alcohol 7%.

Inactive ingredients: Artificial and natural flavors, citric acid, D&C Red 33, FD&C Yellow 6, glycerin, saccharin sodium, sodium benzoate, sodium citrate, sodium propionate, water, and other ingredients.


Codeine is one of the naturally occurring phenanthrene alkaloids of opium derived from the opium poppy, it is classified pharmacologically as a narcotic analgesic. Codeine phosphate may be chemically designated as 7,8-Didehydro-4, 5 $\alpha$ -epoxy-3-methoxy-17-methylmorphinan-6 $\alpha$ -ol phosphate (1:1) (salt) hemihydrate.

The phosphate salt of codeine occurs as white, needle-shaped crystals or white crystalline powder. Codeine phosphate is freely soluble in water and slightly soluble in alcohol. It has a molecular weight of 406.37, a molecular formula of  $C_{18}H_{21}NO_3 \cdot H_3PO_4 \cdot \frac{1}{2} H_2O$  and the following structural formula:




Promethazine hydrochloride, a phenothiazine derivative, is chemically designated as  $(\pm)$ -10-[2-(Dimethylamino)propyl] phenothiazine monohydrochloride.

Promethazine hydrochloride occurs as a white to faint yellow, practically odorless, crystalline powder which slowly oxidizes and turns blue on prolonged exposure to air. It is soluble in water and freely soluble in alcohol. It has a molecular weight of 320.88, a molecular formula of  $C_{17}H_{20}N_2S \cdot HCl$ , and the following structural formula:



Phenylephrine hydrochloride is a sympathomimetic amine salt which is chemically designated as  $(-)$ -*m*-hydroxy- $\alpha$ -[(methyl-amino)methyl] benzyl alcohol hydrochloride. It occurs as white or nearly white crystals, having a bitter taste. It is freely soluble in water and alcohol.

Phenylephrine hydrochloride is subject to oxidation and must be protected from light and air. It has a molecular weight of 203.67, a molecular formula of  $C_9H_{13}NO_2 \cdot HCl$  and the following structural formula:



## CLINICAL PHARMACOLOGY

### Codeine

Narcotic analgesics, including codeine, exert their primary effects on the central nervous system and gastrointestinal tract. The analgesic effects of codeine are due to its central action; however, the precise sites of action have not been determined, and the mechanisms involved appear to be quite complex. Codeine resembles morphine both structurally and pharmacologically, but its actions at the doses of codeine used therapeutically are milder, with less sedation, respiratory depression and gastrointestinal, urinary, and pupillary effects. Codeine produces an increase in biliary tract pressure, but less than morphine or meperidine. Codeine is less constipating than morphine.

Codeine has good antitussive activity, although less than that of morphine at equal doses. It is used in preference to morphine, because side effects are infrequent at the usual antitussive dose of codeine.

Codeine in oral therapeutic dosage does not usually exert major effects on the cardiovascular system. Narcotic analgesics may cause nausea and vomiting by stimulating the chemoreceptor trigger zone (CTZ); however, they also depress the vomiting center, so that subsequent doses are unlikely to produce vomiting. Nausea is minimal after usual oral doses of codeine.

Narcotic analgesics cause histamine release, which appears to be responsible for wheals or urticaria sometimes seen at the site of injection on parenteral administration. Histamine release may also produce dilation of cutaneous blood vessels, with resultant flushing of the face and neck, pruritus, and sweating.

Codeine and its salts are well absorbed following both oral and parenteral administration. Codeine is about 2/3 as effective orally as parenterally. Codeine is metabolized primarily in the liver by enzymes of the endoplasmic reticulum, where it undergoes O-demethylation, N-demethylation, and partial conjugation with glucuronic acid. The drug is excreted primarily in the urine, largely as inactive metabolites and small amounts of free and conjugated morphine. Negligible amounts of codeine and its metabolites are found in the feces.

Following oral or subcutaneous administration of codeine, the onset of analgesia occurs within 15 to 30 minutes and lasts for four to six hours.

The cough-depressing action, in animal studies, was observed to occur 15 minutes after oral administration of codeine, peak action at 45 to 60 minutes after ingestion. The duration of action, which is dose-dependent, usually did not exceed 3 hours.

### **Promethazine**

Promethazine is a phenothiazine derivative which differs structurally from the antipsychotic phenothiazines by the presence of a branched side chain and no ring substitution. It is thought that this configuration is responsible for its relative lack (1/10 that of chlorpromazine) of dopamine antagonist properties.

Promethazine is an H<sub>1</sub> receptor blocking agent. In addition to its antihistaminic action, it provides clinically useful sedative and antiemetic effects.

Promethazine is well absorbed from the gastrointestinal tract. Clinical effects are apparent within 20 minutes after oral administration and generally last four to six hours, although they may persist as long as 12 hours. Promethazine is metabolized by the liver to a variety of compounds;

the sulfoxides of promethazine and N-demethylpromethazine are the predominant metabolites appearing in the urine.

### **Phenylephrine**

Phenylephrine is a potent postsynaptic- $\alpha$ -receptor agonist with little effect on  $\beta$  receptors of the heart. Phenylephrine has no effect on  $\beta$ -adrenergic receptors of the bronchi or peripheral blood vessels. A direct action at receptors accounts for the greater part of its effects, only a small part being due to its ability to release norepinephrine.

Therapeutic doses of phenylephrine mainly cause vasoconstriction. Phenylephrine increases resistance and, to a lesser extent, decreases capacitance of blood vessels. Total peripheral resistance is increased, resulting in increased systolic and diastolic blood pressure. Pulmonary arterial pressure is usually increased, and renal blood flow is usually decreased. Local vasoconstriction and hemostasis occur following topical application or infiltration of phenylephrine into tissues.

The main effect of phenylephrine on the heart is bradycardia; it produces a positive inotropic effect on the myocardium in doses greater than those usually used therapeutically. Rarely, the drug may increase the irritability of the heart, causing arrhythmias. Cardiac output is decreased slightly. Phenylephrine increases the work of the heart by increasing peripheral arterial resistance. Phenylephrine has a mild central stimulant effect.

Following oral administration or topical application of phenylephrine to the mucosa, constriction of blood vessels in the nasal mucosa relieves nasal congestion associated with allergy or head colds. Following oral administration, nasal decongestion may occur within 15 or 20 minutes and may persist for up to 4 hours.

Phenylephrine is irregularly absorbed from and readily metabolized in the gastrointestinal tract. Phenylephrine is metabolized in the liver and intestine by monoamine oxidase. The metabolites and their route and rate of excretion have not been identified. The pharmacologic action of phenylephrine is terminated at least partially by uptake of the drug into tissues.

### **INDICATIONS AND USAGE**

Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution is indicated for the temporary relief of coughs and upper respiratory symptoms, including nasal congestion, associated with allergy or the common cold.

### **CONTRAINDICATIONS**

The combination of promethazine hydrochloride, phenylephrine hydrochloride and codeine phosphate is contraindicated in pediatric patients less than 6 years of age, because the combination may cause fatal respiratory depression in this age population.

Codeine sulfate is contraindicated for post-operative pain management in children who have undergone tonsillectomy and/or adenoidectomy. (See **WARNINGS - Death Related to Ultra-Rapid Metabolism of Codeine to Morphine**).

Codeine is contraindicated in patients with a known hypersensitivity to the drug.

Promethazine is contraindicated in comatose states, and in individuals known to be hypersensitive or to have had an idiosyncratic reaction to promethazine or to other phenothiazines.

Antihistamines and codeine are both contraindicated for use in the treatment of lower respiratory tract symptoms, including asthma.

Phenylephrine is contraindicated in patients with hypertension or with peripheral vascular insufficiency (ischemia may result with risk of gangrene or thrombosis of compromised vascular beds). Phenylephrine should not be used in patients known to be hypersensitive to the drug or in those receiving a monoamine oxidase inhibitor (MAOI).

## **WARNINGS**

(see **Boxed Warnings**)

### **Respiratory Depression in Children**

The combination of promethazine hydrochloride and codeine phosphate is contraindicated in pediatric patients less than 6 years of age. Concomitant administration of promethazine products with other respiratory depressants has an association with respiratory depression, and sometimes death, in pediatric patients.

Postmarketing cases of respiratory depression, including fatalities, have been reported with use of promethazine hydrochloride in pediatric patients less than 2 years of age. A wide range of weight-based doses of promethazine hydrochloride have resulted in respiratory depression in these patients.

Respiratory depression leading to arrest, coma, and death has occurred with the use of codeine antitussives in young children, particularly in the under-one-year infants whose ability to deactivate the drug is not fully developed.

### **Codeine**

- **Death Related to Ultra-Rapid Metabolism of Codeine to Morphine**

Respiratory depression and death have occurred in children who received codeine in the post-operative period following tonsillectomy and/or adenoidectomy and had evidence of being ultra-rapid metabolizers of codeine (i.e., multiple copies of the gene for cytochrome P450 isoenzyme 2D6 [CYP2D6] or high morphine concentrations). Deaths have also occurred in nursing infants who were exposed to high levels of morphine in breast milk because their mothers were ultra-rapid metabolizers of codeine. (See **PRECAUTIONS -Nursing Mothers**).

Some individuals may be ultra-rapid metabolizers because of a specific CYP2D6 genotype (gene duplications denoted as \*1/\*1xN or \*1/\*2xN). The prevalence of this CYP2D6 phenotype varies widely and has been estimated at 0.5 to 1% in Chinese and Japanese, 0.5 to 1% in Hispanics, 1 to 10% in Caucasians, 3% in African Americans, and 16 to 28% in North Africans, Ethiopians, and Arabs. Data are not available for other ethnic groups. These individuals convert codeine into its active metabolite, morphine, more rapidly and completely than other people. This rapid conversion results in higher than expected serum morphine levels. Even at labeled dosage regimens, individuals who are ultra-rapid metabolizers may have life-threatening or fatal respiratory depression or experience signs of overdose (such as extreme sleepiness, confusion, or shallow breathing). (See **OVERDOSAGE**).

Children with obstructive sleep apnea who are treated with codeine for post-tonsillectomy and/or adenoidectomy pain may be particularly sensitive to the respiratory depressant effects of codeine that has been rapidly metabolized to morphine. Codeine is contraindicated for post-operative pain management in all pediatric patients undergoing tonsillectomy and/or adenoidectomy. (See **CONTRAINDICATIONS**).

When prescribing codeine-containing drugs, healthcare providers should choose the lowest effective dose for the shortest period of time and inform patients and caregivers about these risks and the signs of morphine overdose. (See **OVERDOSAGE**).

- **Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**  
Concomitant use of opioids, including Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution, with benzodiazepines, or other CNS depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Because of these risks, avoid use of opioid cough medications in patients taking benzodiazepines, other CNS depressants, or alcohol. (See **PRECAUTIONS - Drug Interactions**).

Observational studies have demonstrated that concomitant use of opioid analgesics and benzodiazepines increases the risk of drug-related mortality compared to use of opioids alone. Because of similar pharmacologic properties, it is reasonable to expect similar risk with concomitant use of opioid cough medications and benzodiazepines, other CNS depressants, or alcohol.

Advise both patients and caregivers about the risks of respiratory depression and sedation if Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution is used with benzodiazepines, alcohol, or other CNS depressants. (See **PRECAUTIONS – Information for Patients**).

- Dosage of codeine **SHOULD NOT BE INCREASED** if cough fails to respond; an unresponsive cough should be re-evaluated in 5 days or sooner for possible underlying pathology, such as a foreign body or lower respiratory tract disease.
- Codeine may cause or aggravate constipation.
- Administration of codeine may be accomplished by histamine release and should be used with caution in atopic children.

- **Head Injury and Increased Intracranial Pressure**

The respiratory-depressant effects of narcotic analgesics and their capacity to elevate cerebrospinal fluid pressure may be markedly exaggerated in the presence of head injury, intracranial lesions, or a pre-existing increase in intracranial pressure. Narcotics may produce adverse reactions which may obscure the clinical course of patients with head injuries.

- **Asthma and Other Respiratory Conditions**

Narcotic analgesics or cough suppressants, including codeine, should not be used in asthmatic patients (see **CONTRAINDICATIONS**). Nor should they be used in acute febrile illness associated with productive cough or in chronic respiratory disease where interference with ability to clear the tracheobronchial tree of secretions would have a deleterious effect on the patient's respiratory function.

- **Hypotensive Effect**

Codeine may produce orthostatic hypotension in ambulatory patients.

### **Promethazine**

- **CNS Depression**

Promethazine may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks, such as driving a vehicle or operating machinery. The impairment may be amplified by concomitant use of other central nervous system depressants such as alcohol, sedatives/hypnotics (including barbiturates), narcotics, narcotic analgesics, general anesthetics, tricyclic antidepressants, and tranquilizers; therefore avoid use of Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution in patients on these medications. (See **PRECAUTIONS – Information for Patients and Drug Interactions**).

- **Respiratory Depression**

Promethazine may lead to potentially fatal respiratory depression.

Use of promethazine in patients with compromised respiratory function (e.g. COPD, sleep apnea) should be avoided.

- **Lower Seizure Threshold**

Promethazine may lower seizure threshold. It should be used with caution in persons with seizure disorders or in persons who are using concomitant medications, such as narcotics or local anesthetics, which may also affect seizure threshold.

- **Bone-Marrow Depression**

Promethazine should be used with caution in patients with bone marrow depression. Leukopenia and agranulocytosis have been reported, usually when promethazine HCl has been used in association with other known marrow toxic agents.

- **Neuroleptic Malignant Syndrome**

A potentially fatal symptom complex sometimes referred to as Neuroleptic Malignant Syndrome (NMS) has been reported in association with promethazine HCl alone or in combination with antipsychotic drugs. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis and cardiac dysrhythmias).

The diagnostic evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to identify cases where the clinical presentation includes both serious medical illness (e.g., pneumonia, systemic infection, etc.) and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever and primary central nervous system (CNS) pathology.

The management of NMS should include 1) immediate discontinuation of promethazine HCl, antipsychotic drugs, if any, and other drugs not essential to concurrent therapy, 2) intensive symptomatic treatment and medical monitoring, and 3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for uncomplicated NMS.

Since recurrences of NMS have been reported with phenothiazines, the reintroduction of promethazine HCl should be carefully considered.

### **Use in Pediatric Patients**

**The combination of promethazine hydrochloride, phenylephrine hydrochloride and codeine phosphate is contraindicated in pediatric patients less than 6 years of age. Concomitant administration of promethazine products with other respiratory depressants has an association with respiratory depression, and sometimes death, in pediatric patients. The association does not directly relate to individualized weight-based dosing, which might otherwise permit safe administration.**

Respiratory depression and death have occurred in children with obstructive sleep apnea who received codeine in the post-operative period following tonsillectomy and/or adenoidectomy and had evidence of being ultra-rapid metabolizers of codeine (i.e., multiple copies of the gene for CYP2D6 or high morphine concentrations). These children may be particularly sensitive to the respiratory depressant effects of codeine that has been rapidly metabolized to morphine. Codeine is contraindicated for post-operative pain management in these patients. (See **WARNINGS - Death Related to Ultra-Rapid Metabolism of Codeine to Morphine and CONTRAINDICATIONS**).

Excessively large dosages of antihistamines, including promethazine hydrochloride, in pediatric patients may cause sudden death (see **OVERDOSAGE**). Hallucinations and convulsions have occurred with therapeutic doses and overdoses of promethazine hydrochloride in pediatric patients. In pediatric patients who are acutely ill associated with dehydration, there is an increased susceptibility to dystonias with the use of promethazine HCl.

### **Other Considerations**

Administration of promethazine has been associated with reported cholestatic jaundice.

### **Phenylephrine**

Because phenylephrine is an adrenergic agent, it should be given with caution to patients with thyroid diseases, diabetes mellitus and heart diseases or those receiving tricyclic antidepressants.

Men with symptomatic, benign prostatic hypertrophy can experience urinary retention when given oral nasal decongestants.

Phenylephrine can cause a decrease in cardiac output, and extreme caution should be used when administering the drug parenterally or orally to patients with arteriosclerosis, to elderly individuals, and/or to patients with initially poor cerebral or coronary circulation.

Phenylephrine should be used with caution in patients taking diet preparations, such as amphetamines or phenylpropanolamine, because synergistic adrenergic effects could result in serious hypertensive response and possible stroke.

## **PRECAUTIONS**

### **General**

Narcotic analgesics, including codeine, should be administered with caution and the initial dose reduced in patients with acute abdominal conditions, convulsive disorders, significant hepatic or renal impairment, fever, hypothyroidism, Addison's disease, ulcerative colitis, prostatic hypertrophy, in patients with recent gastrointestinal or urinary tract surgery and in the very young or elderly or debilitated patients.

Drugs having anticholinergic properties should be used with caution in patients with narrow-angle glaucoma, prostatic hypertrophy, stenosing peptic ulcer, pyloroduodenal obstruction, and bladder-neck obstruction.

Promethazine should be used cautiously in persons with cardiovascular disease, or with impairment of liver function.

Phenylephrine should be used with caution in patients with cardiovascular disease.

### **Information for Patients**

Patients should be advised to measure Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution with an accurate measuring device. A household teaspoon is not an accurate measuring device and could lead to overdosage, especially when a half a teaspoon is measured. A pharmacist can recommend an appropriate measuring device and can provide instructions for measuring the correct dose.

Promethazine, phenylephrine and codeine may cause marked drowsiness or may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks, such as driving a vehicle or operating machinery. Ambulatory patients should be told to avoid engaging in such activities until it is known that they do not become drowsy or dizzy from promethazine, phenylephrine and codeine therapy. Pediatric patients should be supervised to avoid potential harm in bike riding or in other hazardous activities.

Inform patients and caregivers that potentially fatal additive effects may occur if Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution is used with benzodiazepines or other CNS depressants, including alcohol. Because of this risk, patients should avoid concomitant use of Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution with benzodiazepines or other CNS depressants, including alcohol. (See **WARNINGS - Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**).

Patients should be advised to report any involuntary muscle movements.

Avoid prolonged exposure to the sun.

Codeine, like other narcotic analgesics, may produce orthostatic hypotension in some ambulatory patients. Patients should be cautioned accordingly.

Advise patients that some people have a genetic variation that results in codeine changing into morphine more rapidly and completely than other people. Most people are unaware of whether they are an ultra-rapid codeine metabolizer or not. These higher-than-normal levels of morphine in the blood may lead to life-threatening or fatal respiratory depression or signs of overdose such as extreme sleepiness, confusion, or shallow breathing. Children with this genetic variation who were prescribed codeine after tonsillectomy and/or adenoidectomy for obstructive sleep apnea may be at greatest risk based on reports of several deaths in this population due to respiratory depression. As a result, codeine is contraindicated in children who undergo tonsillectomy and/or adenoidectomy. Advise caregivers of children receiving codeine for other reasons to monitor for signs of respiratory depression. (See **WARNINGS - Death Related to Ultra-Rapid Metabolism of Codeine to Morphine**).

Nursing mothers taking codeine can also have higher morphine levels in their breast milk if they are ultra-rapid metabolizers. These higher levels of morphine in breast milk may lead to life-threatening or fatal side effects in nursing babies. Instruct nursing mothers to watch for signs of morphine toxicity in their infants including increased sleepiness (more than usual), difficulty breastfeeding, breathing difficulties, or limpness. Instruct nursing mothers to talk to the baby's doctor immediately if they notice these signs and, if they can not reach the doctor right away, to take the baby to an emergency room or call 911 (or local emergency services).

### **Drug Interactions**

#### **Codeine**

In patients receiving MAO inhibitors, an initial small test dose is advisable to allow observation of any excessive narcotic effects or MAOI interaction.

The use of benzodiazepines, opioids, antihistamines, antipsychotics, anti-anxiety agents, or other CNS depressants (including alcohol) concomitantly with Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution may cause an additive CNS depressant effect, profound sedation, respiratory depression, coma, and death and should be avoided. (See **WARNINGS - Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**).

#### **Promethazine**

*Epinephrine:* Because of the potential for promethazine to reverse epinephrine's vasopressor effect, epinephrine should NOT be used to treat hypotension associated with promethazine overdose.

*Anticholinergics:* Concomitant use of other agents with anticholinergic properties should be undertaken with caution.

*Monoamine Oxidase Inhibitors (MAOI):* Drug interactions, including an increased incidence of extrapyramidal effects, have been reported when some MAOI and phenothiazines are used concomitantly.

## Phenylephrine

Drug	Effect
Phenylephrine with prior administration of monoamine oxidase inhibitors (MAOI).	Cardiac pressor response potentiated. May cause acute hypertensive crisis.
Phenylephrine with tricyclic antidepressants.	Pressor response increased.
Phenylephrine with ergot alkaloids.	Excessive rise in blood pressure.
Phenylephrine with bronchodilator sympathomimetic agents and with epinephrine or other sympathomimetics.	Tachycardia or other arrhythmias may occur.
Phenylephrine with prior administration of propranolol or other $\beta$ -adrenergic blockers.	Cardiostimulating effects blocked.
Phenylephrine with atropine sulfate.	Reflex bradycardia blocked; pressor response enhanced.
Phenylephrine with prior administration of phentolamine or other $\alpha$ -adrenergic blockers.	Pressor response decreased.
Phenylephrine with diet preparations, such as amphetamines or phenylpropanolamine.	Synergistic adrenergic response.

### Drug/Laboratory Test Interactions

Because narcotic analgesics may increase biliary tract pressure, with resultant increase in plasma amylase or lipase levels, determination of these enzyme levels may be unreliable for 24 hours after a narcotic analgesic has been given.

The following laboratory tests may be affected in patients who are receiving therapy with promethazine hydrochloride.

**Pregnancy Tests:** Diagnostic pregnancy tests based on immunological reactions between HCG and anti-HCG may result in false-negative or false-positive interpretations.

**Glucose Tolerance Test:** An increase in blood glucose has been reported in patients receiving promethazine.

### Carcinogenesis, Mutagenesis, Impairment of Fertility

**Codeine and Promethazine:** Long-term animal studies have not been performed to assess the carcinogenic potential of codeine or of promethazine, nor are there other animal or human data concerning carcinogenicity, mutagenicity, or impairment of fertility with these agents. Codeine has been reported to show no evidence of carcinogenicity or mutagenicity in a variety of test systems, including the micronucleus and sperm abnormality assays and the *Salmonella* assay. Promethazine was nonmutagenic in the *Salmonella* test system of Ames.

**Phenylephrine:** A study which followed the development of cancer in 143,574 patients over a four-year period indicated that in 11,981 patients who received phenylephrine (systemic or topical), there was no statistically significant association between the drug and cancer at any or all sites.

Long-term animal studies have not been performed to assess the carcinogenic potential of phenylephrine, nor are there other animal or human data concerning mutagenicity.

A study of the effects of adrenergic drugs on ovum transport in rabbits indicated that treatment with phenylephrine did not alter incidence of pregnancy; the number of implantations was significantly reduced when high doses of the drug were used.

## **Pregnancy**

### **Teratogenic Effects**

*Pregnancy Category C:*

Codeine: A study in rats and rabbits reported no teratogenic effect of codeine administered during the period of organogenesis in doses ranging from 5 to 120 mg/kg. In the rat, doses at the 120-mg/kg level, in the toxic range for the adult animal, were associated with an increase in embryo resorption at the time of implantation. In another study a single 100-mg/kg dose of codeine administered to pregnant mice reportedly resulted in delayed ossification in the offspring.

There are no studies in humans, and the significance of these findings to humans, if any, is not known.

Promethazine: Teratogenic effects have not been demonstrated in rat-feeding studies at doses of 6.25 and 12.5 mg/kg of promethazine HCl. These doses are from approximately 2.1 to 4.2 times the maximum recommended total daily dose of promethazine for a 50-kg subject, depending on the indication for which the drug is prescribed. Daily doses of 25 mg/kg intraperitoneally have been found to produce fetal mortality in rats.

Specific studies to test the action of the drug on parturition, lactation, and development of the animal neonate were not done, but a general preliminary study in rats indicated no effect on these parameters. Although antihistamines have been found to produce fetal mortality in rodents, the pharmacological effects of histamine in the rodent do not parallel those in man. There are no adequate and well-controlled studies of promethazine in pregnant women.

Phenylephrine: A study in rabbits indicated that continued moderate overexposure to phenylephrine (3 mg/day) during the second half of pregnancy (22<sup>nd</sup> day of gestation to delivery) may contribute to perinatal wastage, prematurity, premature labor, and possibly fetal anomalies; when phenylephrine (3 mg/day) was given to rabbits during the first half of pregnancy (3<sup>rd</sup> day after mating for seven days), a significant number gave birth to litters of low birth weight. Another study showed that phenylephrine was associated with anomalies of aortic arch and with ventricular septal defect in the chick embryo.

Animal reproduction studies have not been conducted with the drug combination – promethazine, phenylephrine and codeine. It is not known whether this drug combination can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution should be given to a pregnant woman only if clearly needed.

Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

### **Nonteratogenic Effects**

Dependence has been reported in newborns whose mothers took opiates regularly during pregnancy. Withdrawal signs include irritability, excessive crying, tremors, hyperreflexia, fever, vomiting and diarrhea. Signs usually appear during the first few days of life.

Promethazine administered to a pregnant woman within two weeks of delivery may inhibit platelet aggregation in the newborn.

### **Labor and Delivery**

Narcotic analgesics cross the placental barrier. The closer to delivery and the larger the dose used, the greater the possibility of respiratory depression in the newborn. Narcotic analgesics should be avoided during labor if delivery of a premature infant is anticipated. If the mother has received narcotic analgesics during labor, newborn infants should be observed closely for signs of respiratory depression. Resuscitation may be required (see **OVERDOSAGE**).

Limited data suggest that use of promethazine hydrochloride during labor and delivery does not have an appreciable effect on the duration of labor or delivery and does not increase the risk of need for intervention in the newborn.

The effect of promethazine and/or codeine on later growth and development of the newborn is unknown.

Administration of phenylephrine to patients in late pregnancy or labor may cause fetal anoxia or bradycardia by increasing contractility of the uterus and decreasing uterine blood flow.

See also **Nonteratogenic Effects**.

### **Nursing Mothers**

Codeine is secreted into human milk. In women with normal codeine metabolism (normal CYP2D6 activity), the amount of codeine excreted into human milk is low and dose-dependent. Despite the common use of codeine products to manage postpartum pain, reports of adverse events in infants are rare. However, some women are ultra-rapid metabolizers of codeine. These women achieve higher-than-expected serum levels of codeine's active metabolite, morphine, leading to higher-than-expected levels of morphine in breast milk and potentially dangerously high serum morphine levels in their breastfed infants. Therefore, maternal use of codeine can potentially lead to serious adverse reactions, including death, in nursing infants.

The risk of infant exposure to codeine and morphine through breast milk should be weighed against the benefits of breastfeeding for both the mother and baby. Caution should be exercised when codeine is administered to a nursing woman. If a codeine containing product is selected, the lowest dose should be prescribed for the shortest period of time to achieve the desired clinical effect. Mothers using codeine should be informed about when to seek immediate medical care and how to identify the signs and symptoms of neonatal toxicity, such as drowsiness or sedation, difficulty breastfeeding, breathing difficulties, and decreased tone, in their baby. Nursing mothers who are ultra-rapid metabolizers may also experience overdose symptoms such as extreme sleepiness, confusion or shallow breathing. Prescribers should closely monitor mother-infant pairs and notify treating pediatricians about the use of codeine during breastfeeding. (See **WARNINGS - Death Related to Ultra-Rapid Metabolism of Codeine to Morphine**.)

It is not known whether phenylephrine or promethazine are excreted in human milk.

### **Pediatric Use**

**The combination of promethazine hydrochloride, phenylephrine hydrochloride and codeine phosphate is contraindicated in pediatric patients less than 6 years of age, because the combination may cause fatal respiratory depression in this age population (see WARNINGS – Boxed Warning and Use in Pediatric Patients).**

Respiratory depression and death have occurred in children with obstructive sleep apnea who received codeine in the post-operative period following tonsillectomy and/or adenoidectomy and had evidence of being ultra-rapid metabolizers of codeine (i.e., multiple copies of the gene for CYP2D6 or high morphine concentrations). These children may be particularly sensitive to the respiratory depressant effects of codeine that has been rapidly metabolized to morphine. Codeine is contraindicated for post-operative pain management in all pediatric patients undergoing tonsillectomy and/or adenoidectomy. (See **WARNINGS - Death Related to Ultra-Rapid Metabolism of Codeine to Morphine** and **CONTRAINDICATIONS**).

### **Geriatric Use**

Clinical studies of Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

Sedating drugs may cause confusion and over-sedation in the elderly; elderly patients generally should be started on low doses of Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution and observed closely.

## **ADVERSE REACTIONS**

### **Codeine**

*Central Nervous System:* CNS depression, particularly respiratory depression, and to a lesser extent circulatory depression; light-headedness, dizziness, sedation, euphoria, dysphoria, headache, transient hallucination, disorientation, visual disturbances and convulsions.

*Cardiovascular:* Tachycardia, bradycardia, palpitation, faintness, syncope, orthostatic hypotension (common to narcotic analgesics).

*Gastrointestinal:* Nausea, vomiting, constipation, and biliary tract spasm. Patients with chronic ulcerative colitis may experience increased colonic motility; in patients with acute ulcerative colitis, toxic dilation has been reported.

*Genitourinary:* Oliguria, urinary retention; antidiuretic effect has been reported (common to narcotic analgesics).

*Allergic:* Infrequent pruritus, giant urticaria, angioneurotic edema, and laryngeal edema.

*Other:* Flushing of the face, sweating and pruritus (due to opiate-induced histamine release); weakness.

### **Promethazine**

*Central Nervous System:* Drowsiness is the most prominent CNS effect of this drug. Sedation, somnolence, blurred vision, dizziness; confusion, disorientation and extrapyramidal symptoms such as oculogyric crisis, torticollis, and tongue protrusion; lassitude, tinnitus, incoordination, fatigue, euphoria, nervousness, diplopia, insomnia, tremors, convulsive seizures, excitation, catatonic-like states, hysteria. Hallucinations have also been reported.

*Cardiovascular:* Increased or decreased blood pressure, tachycardia, bradycardia, faintness.

*Dermatologic:* Dermatitis, photosensitivity, urticaria.

*Hematologic:* Leukopenia, thrombocytopenia, thrombocytopenic purpura, agranulocytosis.

*Gastrointestinal:* Dry mouth, nausea, vomiting, jaundice.

*Respiratory:* Asthma, nasal stuffiness, respiratory depression (potentially fatal) and apnea (potentially fatal) (see **WARNINGS – Promethazine; Respiratory Depression**).

*Other:* Angioneurotic edema. Neuroleptic malignant syndrome (potentially fatal) has also been reported (see **WARNINGS – Promethazine; Neuroleptic Malignant Syndrome**).

*Paradoxical Reactions:* Hyperexcitability and abnormal movements have been reported in patients following a single administration of promethazine HCl. Consideration should be given to the discontinuation of promethazine HCl and to the use of other drugs if these reactions occur. Respiratory depression, nightmares, delirium and agitated behavior have also been reported in some of these patients.

### **Phenylephrine**

*Central Nervous System:* Restlessness, anxiety, nervousness and dizziness.

*Cardiovascular:* Hypertension (see **WARNINGS**).

*Other:* Primordial pain, respiratory distress, tremor and weakness.

## **DRUG ABUSE AND DEPENDENCE**

### **Controlled Substance**

Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution is a Schedule V Controlled Substance.

### **Abuse**

Codeine is known to be subject to abuse; however, the abuse potential of oral codeine appears to be quite low. Even parenteral codeine does not appear to offer the psychic effects sought by addicts to the same degree as heroin or morphine. However, codeine must be administered only under close supervision to patients with a history of drug abuse or dependence.

## **Dependence**

Psychological dependence, physical dependence and tolerance are known to occur with codeine.

## **OVERDOSAGE**

### **Codeine**

Serious overdose with codeine is characterized by respiratory depression (a decrease in respiratory rate and/or tidal volume, Cheyne-Stokes respiration, cyanosis), extreme somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin and sometimes bradycardia and hypotension. The triad of coma, pinpoint pupils and respiratory depression is strongly suggestive of opiate poisoning. In severe overdose, particularly by the intravenous route, apnea, circulatory collapse, cardiac arrest, and death may occur. Promethazine is additive to the depressant effects of codeine.

It is difficult to determine what constitutes a standard toxic or lethal dose. However, the lethal oral dose of codeine in an adult is reported to be in the range of 0.5 to 1.0 gram. Infants and children are believed to be relatively more sensitive to opiates on a body-weight basis. Elderly patients are also comparatively intolerant to opiates.

### **Promethazine**

Signs and symptoms of overdose with promethazine HCl range from mild depression of the central nervous system and cardiovascular system to profound hypotension, respiratory depression, unconsciousness and sudden death. Other reported reactions include hyperreflexia, hypertonia, ataxia, athetosis and extensor-plantar reflexes (Babinski reflex).

Stimulation may be evident, especially in children and geriatric patients. Convulsions may rarely occur. A paradoxical reaction has been reported in children receiving single doses of 75 mg to 125 mg orally, characterized by hyperexcitability and nightmares.

Atropine-like signs and symptoms – dry mouth, fixed dilated pupils, flushing, as well as gastrointestinal symptoms, may occur.

### **Phenylephrine**

Signs and symptoms of overdose with phenylephrine include hypertension, headache, convulsions, cerebral hemorrhage, and vomiting. Ventricular premature beats and short paroxysms of ventricular tachycardia may also occur. Headache may be a symptom of hypertension. Bradycardia may also be seen early in phenylephrine overdose through stimulation of baroreceptors.

### **Treatment**

The treatment of overdose with promethazine, phenylephrine and codeine is essentially symptomatic and supportive. Only in cases of extreme overdose or individual sensitivity do vital signs including respiration, pulse, blood pressure, temperature, and EKG need to be monitored. Activated charcoal orally or by lavage may be given, or sodium or magnesium sulfate orally as a cathartic. Attention should be given to the re-establishment of adequate respiratory exchange through provision of a patent airway and institution of assisted or controlled ventilation. The narcotic antagonist, naloxone hydrochloride, may be administered when

significant respiratory depression occurs with promethazine, phenylephrine and codeine; any depressant effects of promethazine are not reversed by naloxone. Diazepam may be used to control convulsions. Avoid analeptics, which may cause convulsions. Acidosis and electrolyte losses should be corrected. A rise in temperature or pulmonary complications may signal the need for institution of antibiotic therapy.

Severe hypotension usually responds to the administration of norepinephrine or phenylephrine. EPINEPHRINE SHOULD NOT BE USED, since its use in a patient with partial adrenergic blockade may further lower the blood pressure.

Limited experience with dialysis indicates that it is not helpful.

## DOSAGE AND ADMINISTRATION

*It is important that Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution is measured with an accurate measuring device (see **PRECAUTIONS-Information for Patients**). A household teaspoon is not an accurate measuring device and could lead to overdosage, especially when half a teaspoon is to be measured. It is strongly recommended that an accurate measuring device be used. A pharmacist can provide an appropriate device and can provide instructions for measuring the correct dose.*

**The combination of promethazine hydrochloride, phenylephrine hydrochloride and codeine phosphate is contraindicated in pediatric patients less than 6 years of age, because the combination may cause fatal respiratory depression in this age population.**

The average effective dose is given in the following table:

<b>Adults (12 years of age and over)</b>	5 mL (1 teaspoonful) every 4 to 6 hours, not to exceed 30.0 mL in 24 hours.
<b>Children 6 years to under 12 years</b>	2.5 mL to 5 mL (½ to 1 teaspoonful) every 4 to 6 hours, not to exceed 30.0 mL in 24 hours.

## HOW SUPPLIED

Promethazine HCl, Phenylephrine HCl and Codeine Phosphate Oral Solution is a clear, reddish-orange solution supplied in bottles of 4 fl. oz. (118 mL) and 16 fl. oz. (473 mL).

**Keep bottles tightly closed.**

**Store at 20° to 25° C (68° to 77° F) [See USP Controlled Room Temperature.]**

**Protect from light.**

**Dispense in tight, light-resistant container (USP/NF) with a child-resistant closure.**

**DEA Order Form Required**

Manufactured for:  
ANI Pharmaceuticals, Inc.  
Baudette, MN 56623



9577 Rev 01/17

## MEDICATION GUIDE

Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution, C-V

### What is the most important information I should know about Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution?

- Taking Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution with benzodiazepines, or other central nervous system depressants, including alcohol can cause severe drowsiness, breathing problems (respiratory depression), coma, and death.
- Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution can cause you to be drowsy. Avoid driving a car or operating machinery during treatment with Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution.
- **Do not** give Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution to children who are under 6 years of age. It may harm them.
- **Do not** give Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution to a child to treat pain after tonsillectomy or adenoidectomy surgery.
  - When you take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution, some of it changes into morphine in your body.
  - In some children and adults this happens very quickly, and can cause you to stop breathing and cause death due to an overdose.
- Women who breastfeed should talk to their healthcare provider before taking Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution.
  - When you take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution, some of it changes into morphine in your body.
  - In some women, this happens very quickly. Codeine and morphine pass into your breast milk. A large amount of morphine can cause your baby to die.
- Call your healthcare provider or get emergency medical help right away if anyone taking Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution, or your breastfeeding baby has any of the symptoms below:
  - increased sleepiness
  - confusion
  - difficulty breathing
  - shallow breathing
  - limpness
  - your baby has difficulty breastfeeding
- Keep Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution in a safe place away from children. Accidental use by a child is a medical emergency and can cause death. If a child accidentally takes Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution, get emergency medical help right away.
- Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution can cause serious side effects including death.
- Take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution exactly as prescribed by your healthcare provider. If you take the wrong dose of Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution, you could overdose and die.
- It is especially important when you take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution that you know exactly what dose to take.

### What is Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution?

- Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution is a prescription medicine used to temporarily treat cough and upper respiratory symptoms, including a stuffy nose (nasal congestion) that you can have with allergies or a common cold. Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution contains 3 medicines, promethazine, phenylephrine and codeine. Promethazine is an antihistamine. Phenylephrine is a decongestant. Codeine is a narcotic cough suppressant.
- **Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution is a federal controlled substance (C-V) because it contains codeine that can be abused or lead to dependence.** Keep Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution in a safe place to prevent misuse and abuse. Selling or giving away Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution may harm others, and is against the law. Tell your healthcare provider if you have ever abused or been dependent on alcohol, prescription medicines or street drugs.
- Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution is not for children under 6 years of age.

### Who should not take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution?

- **Do not** give Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution to children who are under 6 years of age.
- **Do not** give Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution to a

child to treat pain after tonsillectomy or adenoidectomy surgery.

- **Do not** take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution if you are allergic to any of the ingredients in Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution. See the end of this Medication Guide for a complete list of ingredients in Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution.
- **Do not** take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution if you have asthma.
- **Do not** take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution if you have high blood pressure or narrowing of blood vessels to your legs or arms (peripheral vascular insufficiency).
- **Do not** take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution if you take a medicine called a Monoamine Oxidase Inhibitor (MAOI).

**Before you take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution, tell your healthcare provider about all of your medical conditions, including if you:**

- have a drug dependence
- have lung or breathing problems including chronic obstructive pulmonary disease (COPD), or sleep apnea
- have had a head injury
- have pain in your stomach-area (abdomen)
- have a history of severe or persistent cough
- have prostate problems
- have problems with your urinary tract (urethral stricture) or difficulty urinating
- have had surgery recently or plan to have surgery
- drink alcohol
- have kidney or liver problems
- are pregnant or plan to become pregnant. It is not known if Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution will harm your unborn baby. You and your healthcare provider should decide if you should take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution while you are pregnant.
- are breastfeeding or plan to breastfeed. Codeine Phosphate passes into your breast milk and may harm your baby. It is not known if Promethazine Hydrochloride or Phenylephrine Hydrochloride can pass into your breast milk. You and your healthcare provider should discuss whether you should take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution or breastfeed. You should not do both.
- have thyroid problems including hypothyroidism
- have Addison's disease
- have glaucoma (increased pressure in eyes)
- have diabetes
- have seizures
- have heart or blood vessel (cardiovascular) problems including high blood pressure or peripheral vascular insufficiency See **“Who should not take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution?”**
- have blood disorders including low white blood cells

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Taking Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution with certain other medicines can cause side effects or affect how well Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution or the other medicines work. Do not start or stop other medicines without talking to your healthcare provider.

Especially tell your healthcare provider if you:

- take pain medicines such as narcotics
- take cold or allergy medicines that contain antihistamines or cough suppressants
- take medicines for mental illness (anti-psychotics, anti-anxiety)
- drink alcohol
- take medicines for depression, including monoamine oxidase inhibitors (MAOIs) or tricyclic antidepressants (TCAs)
- take medicines for stomach or intestinal problems
- take ergot containing medicines
- take medicines called alpha blockers
- take medicines called beta blockers
- take stimulants or uppers (amphetamines)

Ask your healthcare provider if you are not sure if you take one of these medicines.

**How should I take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution?**

- Take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution exactly as your healthcare provider tells you to take it.
- Your healthcare provider will tell you how much Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution to take and when to take it. Do not change your dose without talking to your healthcare provider.
- Ask your pharmacist to give you a measuring device to help you measure the correct amount of Promethazine

Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution. Do not use a household teaspoon to measure your medicine. You may accidentally take too much.

- **Do not** take more than 30 mL of Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution in 24 hours.
- If you take too much Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution, call your healthcare provider or go to the nearest hospital emergency room right away.

### **What should I avoid while taking Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution?**

- Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution can cause you to be drowsy. Avoid driving a car or operating machinery during treatment with Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution. Also, watch your child closely when riding a bike.
- Avoid drinking alcohol during treatment with Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution. Drinking alcohol can increase your chances of having serious side effects.
- Avoid spending a long time in sunlight.

### **What are the possible side effects of Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution?**

Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution may cause serious side effects, including:

- **See “What is the most important information I should know about Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution?”**
- **Breathing problems (respiratory depression)** which can lead to death. Call your healthcare provider or get emergency treatment right away if you have excessive sleepiness, shallow or slow breathing, or confusion.
- **Physical dependence or abuse.** Take Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution exactly as your healthcare provider tells you to take it. Stopping Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution suddenly could cause withdrawal symptoms.
- **Bowel problems** including constipation or stomach pain.
- **Increased intracranial pressure.**
- **Feeling dizzy, faint or light-headed, especially when you stand up (orthostatic hypotension).** Light-headedness or fainting may happen when getting up too quickly from a sitting or lying position.
- **Neuroleptic malignant syndrome (NMS) which can lead to death.** Tell your healthcare provider right away if you have any of the following symptoms of NMS:
  - high fever
  - confusion
  - changes in pulse, heart rate, and blood pressure
  - stiff muscles
  - sweating

The most common side effects of Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution include:

- |                                  |                        |                       |
|----------------------------------|------------------------|-----------------------|
| • sleepiness                     | • nausea and vomiting  | • high blood pressure |
| • confusion                      | • difficulty urinating | • restlessness        |
| • dryness of mouth, nose, throat | • trouble breathing    | • fast heart beat     |

These are not all the possible side effects of Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution?**

- Store Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution in a tightly closed, child-resistant container and out of the light.
- Safely throw away medicine that is out of date or no longer needed.
- **Keep Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution and all medicines out of the reach of children.**

### **General information about the safe and effective use of Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution for a condition for which it was not prescribed. Do not give Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution to other people, even if they have the same symptoms that you have. It may harm them.

You can ask your pharmacist or healthcare provider for information about Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution that is written for health professionals.

**What are the ingredients in Promethazine Hydrochloride, Phenylephrine Hydrochloride and Codeine Phosphate Oral Solution?**

**Active ingredients:** codeine phosphate, promethazine hydrochloride, and phenylephrine hydrochloride

**Inactive ingredients:** artificial and natural flavors, citric acid, D&C Red 33, FD&C Yellow 6, glycerin, saccharin sodium, sodium benzoate, sodium citrate, sodium propionate, water, and other ingredients.

Manufactured for: ANI Pharmaceuticals, Inc., Baudette, MN 56623  
For more information, go to [www.anipharma.com](http://www.anipharma.com) or call 1-800-308-6755.



This Medication Guide has been approved by the U.S. Food and Drug Administration

Issued: January/2017