

4-ounce Dyna-Hex 4

▲ Lift Here

NDC 0116-1061-04

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic
Non-sterile Solution

Contains: 4% Chlorhexidine Gluconate
Distributed By: Xttrium Laboratories, Inc.
Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

For Single-Use



0 11610 61040 4

Net Contents: 4 fl oz (118 mL)

Drug Facts	Purpose
Active ingredient Chlorhexidine gluconate 4% solution	Antiseptic
Uses <ul style="list-style-type: none">■ surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care■ healthcare personnel handwash: helps reduce bacteria that potentially can cause disease■ patient preoperative skin preparation: for the preparation of the patient's skin prior to surgery■ skin wound and general skin cleansing	

4-ounce Dyna-Hex 4 (Continued)

Drug Facts (continued)

Warnings

For external use only

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area
- as a preoperative skin preparation of the head or face

When using this product

- keep out of the eyes, ears and mouth. May cause serious and permanent eye injury if placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Drug Facts (continued)

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Patient preoperative skin preparation:

- apply product liberally to surgical site and swab for at least 2 minutes and dry with a sterile towel

Drug Facts (continued)

- repeat procedure for an additional 2 minutes and dry with a sterile towel

Skin wound and general skin cleaning:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol

Questions or comments?

call 1-800-587-3721

Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions:

Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

8-ounce Dyna-Hex 4

NDC 0116-1061-08

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic

Contains: 4% Chlorhexidine Gluconate

Distributed By: Xttrium Laboratories, Inc.
Mount Prospect, IL 60056



FOR EXTERNAL USE ONLY

Net Contents: 8 fl oz (237 mL)

4DY08BTLBLD

Drug Facts	Purpose
Active ingredient Chlorhexidine gluconate 4% solution	Antiseptic
Uses <ul style="list-style-type: none">■ surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care■ healthcare personnel handwash: helps reduce bacteria that potentially can cause disease■ skin wound and general skin cleansing	
Warnings For external use only	
Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: <ul style="list-style-type: none">■ wheezing/difficulty breathing■ shock	

▲ Lift Here

8-ounce Dyna-Hex 4 (Continued)

Drug Facts (continued)

- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area

When using this product

- keep out of eyes, ears and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly
- repeat procedure for an additional 2 minutes and dry with a sterile towel

Drug Facts (continued)

Skin wound and general skin cleaning:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol

Questions or comments?

call 1-800-587-3721

Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions:

Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

16-ounce Dyna-Hex 4

NDC 0116-1061-16

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic

Contains: 4% Chlorhexidine Gluconate

Distributed By: Xttrium Laboratories, Inc.
Mount Prospect, IL 60056



0 11610 61160 9

FOR EXTERNAL USE ONLY

Net Contents: 16 fl oz (473 mL)

Drug Facts

Active ingredient
Chlorhexidine gluconate 4% solution Antiseptic

Purpose
Antiseptic

Uses

- surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- healthcare personnel hand wash: helps reduce bacteria that potentially can cause disease
- skin wound and general skin cleansing

Warnings
For external use only

Allergy alert:
This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

▲ Lift Here

4DY16BTL1BLD

16-ounce Dyna-Hex 4 (Continued)

Drug Facts (continued)

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area

When using this product

- keep out of eyes, ears and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- If solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Skin wound and general skin cleansing:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol.

Drug Facts (continued)

Questions or comments?

call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions:

Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

NDC 0116-1061-30

DYNA-HEX 4[®]

Chlorhexidine Gluconate 4% Solution

Antiseptic

Contains: 4% Chlorhexidine Gluconate
 Distributed By: Xttrium Laboratories, Inc.
 Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 30 fl oz (1 qt) (887 mL)

To Install

1. Twist to remove plug from back of dispenser spout.
2. Seat dispenser into bracket, taking care to insert the open air inlet into proper receptacle.
3. Twist to remove plug from end of spout.

To Remove

1. Hold bracket against the wall and pull the empty dispenser straight forward.
2. Discard entire bottle and spout.



Drug Facts

Active ingredient
 Chlorhexidine gluconate 4% solution Antiseptic

Purpose

Uses

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease

Warnings

For external use only

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you are allergic to chlorhexidine gluconate or any other ingredient in this product

When using this product

- Keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- If solution should contact these areas, rinse out promptly and thoroughly with water
- do not use routinely if you have wounds which involve more than the superficial layers of the skin

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol.

Questions or comments?

call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions: Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

4DY30BTLBL

NDC 0116-1061-32

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic

Contains: 4% Chlorhexidine Gluconate
 Distributed By: Xttrium Laboratories, Inc.
 Mount Prospect, IL 60056



FOR EXTERNAL USE ONLY

Net Contents: 32 fl oz (1 qt) (946 mL)

4DY32BTLLBLB

Drug Facts

Active ingredient Chlorhexidine gluconate 4% solution Antiseptic
Purpose Antiseptic

Uses
 ■ surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
 ■ healthcare personnel handwash: helps reduce bacteria that potentially can cause disease
 ■ skin wound and general skin cleansing

Warnings
 For external use only
Allergy alert:
 This product may cause a severe allergic reaction. Symptoms may include:
 ■ wheezing/difficulty breathing
 ■ shock
 ■ facial swelling
 ■ hives
 ■ rash
 If an allergic reaction occurs, stop use and seek medical help right away.

Do not use
 ■ if you or the patient are allergic to chlorhexidine gluconate or any other ingredient in this product
 ■ in contact with meninges
 ■ in the genital area

When using this product
 ■ keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
 ■ if solution should contact these areas, rinse out promptly and thoroughly with water
 ■ wounds which involve more than the superficial layers of the skin should not be routinely treated
 ■ repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Drug Facts (continued)

Keep out of reach of children.
 If swallowed, get medical help or contact a Poison Control Center right away.

Directions
 ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
Surgical hand scrub:
 ■ wet hands and forearms with water
 ■ scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces
 ■ a separate nail cleaner may be used
 ■ rinse thoroughly
 ■ wash for an additional 3 minutes with 5 mL of product and rinse under running water
 ■ dry thoroughly
Healthcare personnel handwash:
 ■ wet hands with water
 ■ dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
 ■ rinse and dry thoroughly
Skin wound and general skin cleansing:
 ■ thoroughly rinse the area to be cleaned with water
 ■ apply the minimum amount of product necessary to cover the skin or wound area and wash gently
 ■ rinse again thoroughly

Other information
 ■ store at 20-25°C (68-77°F)
 ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol.

Questions or comments?
 call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions: Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

128-ounce (1-gallon) Dyna-Hex 4

▲ Lift Here for Drug Facts NDC 0116-1061-01

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic

Contains: 4% Chlorhexidine Gluconate

Distributed By: Xttrium Laboratories, Inc.
Mount Prospect, IL 60056


0 11610 61010 7

FOR EXTERNAL USE ONLY

Net Contents: 128 fl oz (1 gal) (3.785 L)

Drug Facts

Active ingredient	Purpose
Chlorhexidine gluconate 4% solution	Antiseptic

Uses

- surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- healthcare personnel handwash: helps reduce bacteria that potentially can cause disease
- skin wound and general skin cleansing

128-ounce (1-gallon) Dyna-Hex 4 (Continued)

Drug Facts (continued)

Warnings

For external use only

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area

When using this product

- keep out of eyes, ears and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Drug Facts (continued)

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Skin wound and general skin cleansing:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tricetyl alcohol


Questions or comments?

call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions:

Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

15-mL Dyna-Hex 4 (carton)-Full view



FOR EXTERNAL USE ONLY

Dyna-Hex 4
Antiseptic
Non-Sterile Solution

4% Chlorhexidine Gluconate
Xtrium Laboratories, Inc.
Mount Prospect, IL 60056

NDC 0116-1061-15

Drug Facts

Active ingredient: Chlorhexidine gluconate

Purpose: Antiseptic

Uses:

- surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- healthcare personnel hand scrub: helps reduce bacteria that ordinarily can cause illness
- patient preoperative skin preparation: for the preparation of the patient's skin prior to surgery
- skin wound and general skin cleansing

Warnings:

For external use only

Always alert:

This product may cause a severe allergic reaction. Symptoms may include:

- severe, generalized itching
- shock
- facial swelling
- hives
- rash

If a severe reaction occurs, stop use and seek medical help right away.

Directions:

- If you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with piercings
- in the genital area
- as a preoperative skin preparation of the head or face

Drug Facts (continued)

When using this product:

- keep out of the eyes, ears and mouth. It may cause irritation and permanent eye injury if placed or kept in the eye during surgical procedures or may cause mechanical injury if used in the mouth or through perforated eardrums.
- if chlorine alcohol contact free areas, rinse and promptly and thoroughly with water
- wear a skin lotion over the top superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and use brush paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- scrub for an additional 2 minutes with 5 mL of product and use brush paying close attention to the nails, cuticles, and interdigital spaces

Healthcare personnel hand scrub:

- wet hands with water
- dispense about 5 mL of product into support hands and scrub in a vigorous manner for 15 seconds
- clean and dry thoroughly

Drug Facts (continued)

Patient preoperative skin preparation:

- apply product liberally to surgical site and neck for at least 2 minutes
- dry with a sterile towel
- repeat procedure for an additional 2 minutes and dry with a sterile towel
- if chlorine alcohol contact free areas, rinse and promptly and thoroughly with water
- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- wear again liberally

Other information:

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients: carbamide DEA, hydrogen peroxide, polyoxyethylene-10 lauryl ether, polyoxyethylene-20 lauryl ether, polyoxyethylene-42 lauryl ether, purified water, alcohol denat.

Questions or comments?

call 1-800-537-8771
Monday through Friday
8:00 AM to 4:00 PM CST

4DY15C7N15L1A

Warnings/Chemical Instructions:

Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

Lot number and expiration date printed on side of box.

NDC 0116-1061-15

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic
Non-Sterile Solution

Contains: 4% Chlorhexidine Gluconate
Distributed By: Xtrium Laboratories, Inc.
Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 50 Single-Use 0.5 fl oz (15-mL) Packettes



NDC 0116-1061-15

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic
Non-Sterile Solution

Contains: 4% Chlorhexidine Gluconate
Xtrium Laboratories, Inc.
Distributed By: Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 50 Single-Use 0.5 fl oz (15-mL) Packettes

NDC 0116-1061-15

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic
Non-Sterile Solution

Contains: 4% Chlorhexidine Gluconate
Distributed By: Xtrium Laboratories, Inc.
Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 50 Single-Use 0.5 fl oz (15-mL) Packettes

LOT# 1061470B
EXP 01/2018

15-mL Dyna-Hex 4 (carton)-Enlargement of Drug Facts panel (Continued)

<p>Drug Facts</p> <p>Active ingredient Purpose</p> <p>Chlorhexidine gluconate 4% solution Antiseptic</p>	<p>Drug Facts (continued)</p> <p>When using this product</p> <ul style="list-style-type: none"> ■ keep out of the eyes, ears and mouth. May cause serious and permanent eye injury if placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums. ■ if solution should contact these areas, rinse out promptly and thoroughly with water ■ wounds which involve more than the superficial layers of the skin should not be routinely treated ■ repeated general skin cleansing of large body areas should not be done except when advised by a health care provider 	<p>Drug Facts (continued)</p> <p>Patient preoperative skin preparation:</p> <ul style="list-style-type: none"> ■ apply product liberally to surgical site and swab for at least 2 minutes ■ dry with a sterile towel ■ repeat procedure for an additional 2 minutes and dry with a sterile towel
<p>Uses</p> <ul style="list-style-type: none"> ■ surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care ■ healthcare personnel handwash: helps reduce bacteria that potentially can cause disease ■ patient preoperative skin preparation: for the preparation of the patient's skin prior to surgery ■ skin wound and general skin cleansing 	<p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>	<p>Skin wound and general skin cleansing:</p> <ul style="list-style-type: none"> ■ thoroughly rinse the area to be cleaned with water ■ apply the minimum amount of product necessary to cover the skin or wound area and wash gently ■ rinse again thoroughly
<p>Warnings</p> <p>For external use only</p> <p>Allergy alert:</p> <p>This product may cause a severe allergic reaction. Symptoms may include:</p> <ul style="list-style-type: none"> ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash <p>If an allergic reaction occurs, stop use and seek medical help right away.</p>	<p>Directions</p> <ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. <p>Surgical hand scrub:</p> <ul style="list-style-type: none"> ■ wet hands and forearms with water ■ scrub for 3 minutes with about 5 mL of product and a wet brush paying close attention to the nails, cuticles, and interdigital spaces ■ a separate nail cleaner may be used ■ rinse thoroughly ■ wash for an additional 3 minutes with 5 mL of product and rinse under running water ■ dry thoroughly <p>Healthcare personnel handwash:</p> <ul style="list-style-type: none"> ■ wet hands with water ■ dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds ■ rinse and dry thoroughly 	<p>Other information</p> <ul style="list-style-type: none"> ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F) <p>Inactive ingredients</p> <p>cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol</p>
<p>Do not use</p> <ul style="list-style-type: none"> ■ if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product ■ in contact with meninges ■ in the genital area ■ as a preoperative skin preparation of the head or face 	<p>Questions or comments?</p> <p>call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST</p> <p>4DY15CTNLBLA</p> <p>Laundering/Cleaning Instructions: Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.</p> <p>Lot number and expiration date printed on side of box</p>	

NDC 0116-1061-15

DYNA-HEX 4[®]
Chlorhexidine Gluconate 4% Solution
Antiseptic
Non-Sterile Solution

Contains: 4% Chlorhexidine Gluconate
Xttrium Laboratories, Inc.
Distributed By: Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 50 Single-Use 0.5 fl oz (15-mL) Packettes

15-mL Dyna-Hex 4 (packet)

NDC 0116-1061-15 Compare to the active ingredient in Hibiclens®*

DYNA-HEX 4®
Chlorhexidine Gluconate 4% Solution
Antiseptic
Non-Sterile Solution

FOR EXTERNAL USE ONLY

Dyna-Hex 4® reduces bacteria on the skin

Dye-Free



Distributed by
Xttrium Laboratories, Inc.
Mount Prospect, IL 60056

Net Contents: 15 mL
Single Use



3 0116106115 3

Please refer to outer packaging for directions for use and product warnings

*This product is not manufactured or distributed by M&Alycke Healthcare, owner of the registered trademark Hibiclens®.



Item Code: 4DYN15INNLEJL

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
09/14/2017