00000000
Drug Facts (continued)
Stomach bleeding warning: This product contains an NSAID, which may cause a stomach bleeding. The chance is higher if you
■ are age 60 or over ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing ibuprofen (NSAIDs), aspirin, diflunisal, or salsalate
■ have a bleeding disorder ■ have cancer ■ have diabetes ■ have heart disease ■ have high blood pressure ■ have liver disease ■ have stomach pain that does not get better
Heart attack and stroke warning: NSAIDs increase the risk of heart failure, heart attack, and stroke. These can be fatal. The risk is higher if you use more than the recommended dosage, for longer than directed
Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery
■ if you are now taking a prescription or nonprescription drug (NSAID) (for blood clots, gout, inflammation, pain, or fever), or have taken one within the last 24 hours (for ibuprofen or naproxen sodium), or within the last 8 hours (for ketoprofen), unless directed by a doctor
■ if you are allergic to any of the following: aspirin, other NSAIDs, or any other product in this family of medicines
■ if you have had asthma, hives, or other allergic reaction after taking aspirin or any other NSAID
■ if you have had a recent or upcoming coronary artery bypass graft (CABG) surgery
Ask a doctor before use if you have
■ heart disease ■ high blood pressure ■ heart failure
■ thyroid disease ■ diabetes
■ trouble or difficulty urinating due to an enlarged prostate gland
■ liver disease or cirrhosis ■ asthma
■ if you are taking a blood thinning (anticoagulant) or steroid drug
Ask a doctor or pharmacist before use if you are
■ taking any other drug containing an NSAID (prescription or nonprescription) and/or a nasal decongestant
■ taking a blood thinning (anticoagulant) or steroid drug
Drug Facts (continued)
Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding
■ vomit blood or have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ symptoms can raise a fever
■ any new symptoms appear
If pregnant or breast feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless clearly directed by a doctor because it may cause a problem in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1 800 222 1222)
Dosage
■ do not exceed maximum dosage
■ take with food or milk if stomach upset occurs
Directions
■ do not take more than directed
■ the smallest effective dose should be used
■ do not take longer than 10 days, unless directed by a doctor (see Warnings)
adults and children 12 years of age and older
■ take 1 caplet every 4 to 6 hours while symptoms persist
■ if symptoms do not respond to 1 caplet, 2 caplets may be used
■ do not use more than 6 caplets in any 24 hour period unless directed by a doctor
Drug Facts (continued)
Other ingredients
■ do not use if blister is open
■ store between 20 - 25C (68 - 77F) ■ avoid excessive heat above 40C (104F)
■ keep out of reach of children
Questions? 1 877 895 3665 (M-F 9am-5pm EST)
Distributed by JOHNSON & JOHNSON CONSUMER INC.
Skillman, NJ 08558 USA
© JJCI 2016
Drug Facts (continued)
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include
■ hives ■ facial swelling (angioedema) ■ shock
■ asthma (wheezing) ■ skin reddening ■ rash
■ blisters
If an allergic reaction occurs, stop use and seek medical help. Keep out of reach of children.
Drug Facts (continued)
Uses
■ temporarily relieves minor aches and pains due to the common cold or flu
■ headache ■ nasal congestion ■ fever ■ minor body aches and pains
Drug Facts (continued)
Active ingredients (in each caplet)
Ibuprofen (NSAID)* 200 mg
Pseudoephedrine HCl 30 mg
Purpose
Pain reliever/fever reducer
Nasal decongestant
*nonsteroidal anti-inflammatory drug
Warnings
Liver warning: This product contains ibuprofen. Ibuprofen may cause severe liver damage. Do not exceed recommended dosage.
Allergy alert: Ibuprofen may cause severe allergic reactions, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help. Keep out of reach of children.
Uses
■ temporarily relieves minor aches and pains due to the common cold or flu
■ headache ■ nasal congestion ■ fever ■ minor body aches and pains
Directions
■ do not take more than directed
■ the smallest effective dose should be used
■ do not take longer than 10 days, unless directed by a doctor (see Warnings)
adults and children 12 years of age and older
■ take 1 caplet every 4 to 6 hours while symptoms persist
■ if symptoms do not respond to 1 caplet, 2 caplets may be used
■ do not use more than 6 caplets in any 24 hour period unless directed by a doctor
Other ingredients
■ do not use if blister is open
■ store between 20 - 25C (68 - 77F) ■ avoid excessive heat above 40C (104F)
■ keep out of reach of children
Questions? 1 877 895 3665 (M-F 9am-5pm EST)
Distributed by JOHNSON & JOHNSON CONSUMER INC.
Skillman, NJ 08558 USA
© JJCI 2016
00000000
See New Warnings
NDC 50580-146-01
Motrin®
Cold & Sinus
contains
Ibuprofen 200 mg/Pseudoephedrine HCl 30 mg
Pain Reliever/Fever Reducer (NSAID)
Nasal Decongestant
▸ Sinus Pressure
▸ Nasal Congestion
20 Coated Caplets*
*Capsule Shaped Tablets
NON-DROWSY FORMULA
Motrin® Cold & Sinus
Ibuprofen 200 mg/Pseudoephedrine HCl 30 mg
EXP:
LOT:
Motrin® Cold & Sinus
Ibuprofen 200 mg/Pseudoephedrine HCl 30 mg
00000000
3 0045-0738-01 1