

X = 4.97      Y = 7.10  
 WGN10231D2  
 01/14/2014  
 PRINTED SIDE RT MTL FOLDING



**Drug Facts**

**Active ingredient Purpose**  
 Adapalene 0.1% (retinoid) Acne treatment  
 \*read consumer information leaflet.

**Use**  
 ■ For the treatment of acne

**Warnings**  
 For external use only

**Do not use**  
 ■ on damaged skin (cuts, abrasions, eczema, sunburn)  
 ■ if you are allergic to adapalene or any of the ingredients in this product.

**If pregnant or breast-feeding**, ask a doctor before use.

**When using this product**  
 ■ limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors  
 ■ do not wax to remove hair in areas where the product has been applied  
 ■ during the early weeks of use, your acne may appear to worsen before it improves (this is normal); continue using as directed, unless you get irritation that becomes severe

■ irritation (redness, itching, dryness, burning) is more likely to occur:  
 ■ in the first few weeks of use  
 ■ if using more than one topical acne medication at a time  
 ■ but irritation usually lessens with continued use of this product

■ it may take up to 3 months of once daily use to see results  
 ■ avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water.  
 ■ wash hands after use

**Stop use and ask a doctor if**  
 ■ you become pregnant, or are planning to become pregnant, while using the product  
 ■ you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)  
 ■ irritation becomes severe

**Drug Facts (continued)**

■ you see no improvement after 3 months of once daily use

**Keep out of reach of children.**  
 If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

Adults and children 12 years of age and older:

- use once daily
- clean the skin gently and pat dry before applying the product
- cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face.
- do not use more than one time a day. Applying more than directed will not provide faster or better results, but may worsen skin irritation.

**Children under 12 years of age:**  
 ask a doctor

**Other information**

- store at room temperature 68°-77°F
- protect from freezing

**Inactive ingredients**

carbomer 940, edetate disodium, methylparaben, poloxamer 182, propylene glycol, purified water and sodium hydroxide.  
 May contain hydrochloric acid to adjust pH.

**Questions?**  
 1-866-735-4137

Distributed by:  
 Galderma Laboratories, L.P.  
 Fort Worth, TX 76177 USA  
 Made in Canada  
 All trademarks are the property of their respective owners

**DIFFERIN®  
 GEL**

ADAPALENE GEL 0.1%  
 ACNE TREATMENT

**DIFFERIN®  
 GEL**

ADAPALENE GEL 0.1%  
 ACNE TREATMENT

- FIRST FDA-APPROVED OVER-THE-COUNTER TOPICAL RETINOID\* FOR ACNE TREATMENT
- PREVIOUSLY AVAILABLE ONLY BY PRESCRIPTION
- DERMATOLOGIST DEVELOPED AND TESTED


\*Read carton and enclosed consumer information leaflet before using this product. Keep this carton and consumer information leaflet. They contain important information.

Once Daily  
 Topical retinoid\*

\*Read consumer information leaflet before use.

Oil Free  
 Fragrance Free  
 FDA Approved

000000-0



ACTUAL SIZE

GALDERMA

NET WT 0.5 OZ (15 g)

WGN10231D2

X = 10.69 Y = 10.53  
WGN10288D12  
05/07/2014  
PRINTED SIDE RT MTL FOLDING



Galderma Laboratories, L.P.

Differin (adapalene) Gel 0.1% - NDA 020380 Seq 0041

Draft Labeling

### Drug Facts

Active ingredient	Purpose
Adapalene 0.1% (retinoid)*	Acne treatment

\*read consumer information leaflet

#### Use

- For the treatment of acne

#### Warnings

##### For external use only

##### Do not use

- on damaged skin (cuts, abrasions, eczema, sunburn)
- if you are allergic to adapalene or any of the ingredients in this product.

If pregnant or breast-feeding, ask a doctor before use.

##### When using this product

- limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors
- do not wax to remove hair in areas where the product has been applied
- during the early weeks of use, your acne may appear to worsen before it improves (this is normal); continue using as directed, unless you get irritation that becomes severe
- irritation (redness, itching, dryness, burning) is more likely to occur:
  - in the first few weeks of use
  - if using more than one topical acne medication at a time
  - but irritation usually lessens with continued use of this product
- it may take up to 3 months of once daily use to see results
- avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water.
- wash hands after use

##### Stop use and ask a doctor if

- you become pregnant, or are planning to become pregnant, while using the product
- you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)
- irritation becomes severe
- you see no improvement after 3 months of once daily use

##### Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

##### Directions

##### Adults and children 12 years of age and older:

- use once daily
- clean the skin gently and pat dry before applying the product
- cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face.
- do not use more than one time a day. Applying more than directed will not provide faster or better results, but may worsen skin irritation.

##### Children under 12 years of age: ask a doctor

##### Other information

- store at room temperature 68°-77°F
- protect from freezing

##### Inactive ingredients

carbomer 940, edetate disodium, methylparaben, poloxamer 182, propylene glycol, purified water and sodium hydroxide.  
May contain hydrochloric acid to adjust pH.

##### Questions? 1-866-735-4137

(b) (4)