

Active ingredient  
(in each tablet)

Loperamide HCl 2 mg..... Anti-diarrheal

Use controls symptoms of diarrhea,  
including Travelers' Diarrhea

LOT

EXP.

XXXXXXXX

Purpose

A-D Loperamide HCl/  
Anti-Diarrheal

Imodium®

A-D Loperamide HCl/  
Anti-Diarrheal

EZ CHEWS

20 Chewable

Cool Mint Flavored Tablets

Warning

Allergy alert: Do not use if you have ever  
had a rash or other allergic reaction to loperamide  
HCl. Heart alert: Taking more than directed can  
cause serious heart problems or death. Do not  
use if you have bloody or black stool. Ask a doctor

before use if you have • fever • mucus in the stool • a history of liver disease. Ask a doctor or pharmacist  
before use if you are taking antibiotics. When using this product • tiredness, drowsiness or dizziness may  
occur. Be careful when driving or operating machinery. Stop use and ask a doctor if • symptoms get worse  
• diarrhea lasts for more than 2 days • you get abdominal swelling or bulging. These may be signs of a  
serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach  
of children. In case of overdose, get medical help or call a Poison Control Center right away.  
(1-800-222-1222). Directions • drink plenty of clear fluids to help prevent dehydration caused by  
diarrhea • take only on an empty stomach (1 hour before or 2 hours after a meal) • if possible, use weight to  
dose; otherwise, use age. Adults and children 12 years and over: chew 2 tablets after the first loose stool;  
chew 1 tablet after each subsequent loose stool; but no more than 4 tablets in 24 hours. Children 9-11 years  
(60-95 lbs): chew 1 tablet after the first loose stool; chew 1/2 tablet af er each subsequent loose stool; but  
(60-95 lbs): chew 1 tablet after the first loose stool; chew 1/2 tablet af er each subsequent loose stool; but  
no more than 3 ablets in 24 hours. Children 6-8 years (48-59 lbs): chew 1 tablet after the first loose stool;  
chew 1/2 tablet after each subsequent loose stool; but no more than 2 tablets in 24 hours. Children 2-5  
years (34 to 47 lbs): ask a doctor. Children under 2 years (up to 33 lbs): do not use. Other information  
• store between 20-25°C (68-77°F) • do not use if printed foil seal under bottle cap is open or torn  
Questions or comments? call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish)  
DISTRIBUTED BY: MCNIEL PPC, INC., FORT WASHINGTON, PA 19034 USA © MCN-PPC, Inc. '06

Reference ID: 4083268



---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/

---

VALERIE S PRATT

04/12/2017