

(b) (4)

Children's Motrin Ibuprofen Oral Suspension (NSAID)

Distributed by: JOHNSON & JOHNSON CONSUMER INC.

Original Berry Flavor

rt Washington, PA 19034

NDC 50580-601-22 For Ages 2 to 11 Years **Children's** Motrin contains **Ibuprofen** Oral Suspension (NSAID) 100 mg per 5 mL Pain Reliever / Fever Reducer Lasts up to 8 hours Alcohol Free A BARA

original Berry Flavor

LOT:

EXP:

fl oz (30 mL)

Important Read all product information before using. Keep this box for important information. This product is intended for use in children. Drug Facts Purpose Uses temporarly: Telleves minor a hes and pains due to he common cold, flu, sore hroat, headache and too hache reduces fever

001

Teduces fever

Wernings
Allergy alert lbuprofen mey cause a severe allergic reaction,
especel y impeople alle gic to aspin. Symptoms may include:

Inse Elicial seveling as time whereing
shoch skin reddning as time whereing
shoch skin reddning as time whereing
shoch skin reddning the skin allolstes
If an alle gic react n occurs, stop use and seek medica helpright
away.
Stomach bleeting warning This product contains an NSAD,
which may cause severe stomach bleeding. The hance is higher if
takes here to uschaining presing in or nonprescription
NSADE (seprin, buprofen, naryor, n, or birs)
takes and to a containing treasition or nonprescription
NSADE (seprin, buprofen, naryor, n, or birs)
takes mere or a longer time than direded
Heart attack and stroke warning, NSAD sevegat aspin,
nonrease her isk is higher if you use more han direct d or for
longer hand fred.
Or strok hard attack, heart bank, heart bane, heart bank. These
can be fall. The risk is higher if you use more han direct d or for
longer hand fred.
Or stroke warning, Severe or persistent sore hroat
or stroke hoat accompanied by high liver, headache, nausea, and
winting may be series. Consult doctor promp Iy.Don to use
more than 2 days or administer to hid in under 3 years of age
unless directed by doctor.



Drug Facts (continued) Drug Facts (chntrued) Do not use in if he ch that ever had an allergic react in to burd nor any other pain reliever fever fuce infort before or after helpt surgery Ast a doctor before use if is man had beeding waning applies to your of it is that problems, such as hearth un is that problems is obtained problems, such as hearth un is that problems. If the substruction finaling parameters or hear reduces is of the final that not been drinking that is in high data state in the substruction for there is which high hold metaway. heard datase, liver that hearth is high hold metaway. heard datase, liver diar hea ■ child has high blood pressure, heart disease, liver cir hosis, k dney disease or had a stroke ■ hild has as hma ■ hild is taking a diuretic Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious c ndition taking any o he d ug When using this product ■ take wih food or m lk if stomach upset ocurs Lake with hisod or mik is bonnaru typestoc urs Stop use and ask a doctor if mich id experiences any of the following signs of the mich hielending: The feels faint wornts blo d "heatbid vyor blads storis". These stimes have in the desender better mich hich has symptoms on heat problems or storie. "Check pain "through threa hing "weakness" in one part or side thoody "suited spee "In leg seeling". The hid does not get any retriev in hing star year of the site of the see or optimum star worden the site of the site of the site of the see of the site of thoody in the site of the site of the site of the see of the site of the see of the site of thoody in the site of Keep out of reach of children. In case of ove dose, get medical help or contact a Poison C introl Center right away. (1-800 222-1222) Directions or complete warnings for adult use ■dd not give more than directed ■ shake wel before using ■ mL = mi liter ■ find right dose n chart. If possible, use weight to dose; →



o herwise use age. Use only enclosed dosing cup. Do not use any o her dosing device. Uf needed, repeat dose every 6-8 hours do not use more han 4 times a day replace original bottle cap to maintain child resistance

Drug Facts (continued)

Dose (mL)*

ask a doctor 5 mL

75 mL

10 mL

3



 Important: Read all product information before using. Keep the box for important information. This product is intended for use in children... Active ing edient(free 1.5 mL)
 Purpose

 top den 0 mg(NAD)
 Painrelever fiver reducer fronze oidlamb in memortry of ug
 Painrelever fiver reducer fiver eligible in memortry of ug

 Uses tempo ally ill releves mino a hes and pains due to the common old, lu, sore h eat, heads he and too to be ill excess feer
 We intes
 he pan ul area ill any rener symp om sappear. Keip out of leach of all dien. n asie of ove dose, ge med die helpfor contact a Poison Cont ol Center right away (800 222-1222). Di ectors ill his p dotact does not contain di ectio s or complete warni gs for adult use ill don dig ill emo e han di ectod il state ve before using ill nit. – mili ter il ni bit dotasen in hetbow. Horosche use welford bose, hore intere use ill evond to diverso director adult use and on dig ill dorice. Ill freeded, speal dose every 6-0 hou s ill donot use more han 4 limes a day ill replace original bolle aplomanitain hit esistance Dosing Chart 50580-601-20 Children's Motrin[®]



- Ask a doctor before use if stomach bleeding warning applies to your child child has a history of stomach problems, such as beauthur.
- child has drawing of standar prosterior, solar heartbun
 child has problems or serious side effects from taking pain relievers or fever reducers child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diar hea
 child has high blood pressure, heart disease, liver cir hosis, kidney disease, or had a stroke

child has asthma

- child is taking a diuretic
- Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition taking any other drug

- When using this product take with food or milk if stomach upset occurs Stop use and ask a doctor if
- child experiences any of the following signs of stomach bleeding:
 feels faint
- vomits blood
- has bloody or black stools
 has stomach pain that does not get better
 child has symptoms of heart problems or stroke:
- chest pain
- trouble breathing
 weakness in one part or side of body
 slurred speech
- leg swelling
- the child does not get any relief w thin first day (24 hours) of treatment
 fever or pain gets worse or lasts more than 3 days redness or swelling is present in the painful area any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions this product does not contain direct complete warnings for adult use
 do not give more than directed ons o

	 shake well before using mL = milliliter find right dose on chart. If possible, use weight to dose; otherwise use age. use only enclosed dosing cup. Do not use any other dosing device. if needed, repeat dose every 6-8 hours do not use more than 4 times a day replace original bottle cap to maintain child resistance Dosing Chart 				
	Weight (lb)	Age (yr)	Dose (mL)*		
	under 24	under 2 years	ask a doctor		
24-35 lbs 2-3 years 5 mL					
	36-47 lbs	7.5 mL			
	48-59 lbs	6-8 years	10 mL		
	60-71 lbs	9-10 years	12.5 mL		

Drug Facts (continued)

or as directed by a doctor Other information each 5 mL contains: sodium 2 mg
store between 20-25'C (68-77'F)
do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

15 mL

72-95 lbs 11 years

Inactive ingredients acesulfame potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium oate, sucrose, xanthan gum

Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by: JOHNSON & JOHNSON CONSUMER INC. McNeil Consumer Healthcare Division Fort Washington, PA 19034 USA ©J&JCI 2016 www.motrin.com 00000000

Children's Motrin buprofen Oral Suspension (NSA

Original Berry Flavor



^{contains} Ibuprofen Oral Suspension (NSAID) *100 mg per 5 mL*

Pain Reliever / Fever Reducer





Original **Berry Flavor**



3 0045- 0192- 04 2

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts Active ingredient Purpose (in each 5 mL) buprofen 100 mg (NSAID)*.....Pain reliever/ *nonste oidal anti-inflammatory drug ever reducer

Uses temporarily:

relieves minor aches and pains due to the common cold, flu, sore th oat, headache and toothache

Warnings Allergy alert: buprofen may cause a seve e allergic reaction, especially in people allergic to aspi in. Symptoms may include:

- thives
 Initial states
 Initinity states
 Initinity
 Initial states
 Initi
- help right away. Stomach bleeding warning: This p oduct dontains an
- NSAID, which may cause severe stomach bleeding. The
- chance is higher if your child:
 has had stomach ulcers or bleeding prob
 takes a blood thinning (anticoagulant) of
- ste oid drug takes other drugs containing prescription or

 a case other of use containing prescription of nonprescription NSAIDs (aspirin, buprolen, naproxen, or others)
 a takes more or for a longer time than directed
 Heart attack and stroke warning NSAIDs, except aspi in, increase the risk of heart attack, heart failure, and st oke. These can be fatal. The risk is higher if you use more than directed tail. In this is figure in you see note than directed Sore threat warning: Severe or persistent one th oat or sore throat accompanied by high fever, heacache, nausea, and vomiting may be serious. Consult doctor p omptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use if the child has ever had an alle gic reaction to bup ofen or any other pain reliever/fever educer right before or after heart su gery







Important: Read all product information before using. Keep the box for important information. This product is intended for use in children. Activing edient/ine a fismultant for the product is intended for use in children. Purpose buy den dimp(RND)" Purpose buy den dimp(RND)" Parelever fevereducer "ronet oddartin-inametry dug Udes tempo arty. Eleises mino a hes and pairs due to the common old, lu, sore host, heada he and loota by a ectores fever Ware the edieses mino a hes and pairs due to the common old, lu, sore host, heada he and loota by a ectores fever Edieses mino a hes and pairs due to the common old, lu, sore host, heada he and loota by a ectores fever War ings Alle qu'after thur ofon may asse a severa elergic eadion, especially in people	Children's Motrino"	hepor contact a Poison Cont of C Di ect ons ■ his p oduct does not contai han di ected ■ shake we befor If poss ble, use weight to dose; o h	lenter right away. (* 800 222-1222). In diectios or complete warni we using ■ mL = m liter ■ inde nerwise use age. ■ use only enclos every 6-8 hous ■ do not use mor every 6-8 hous ■ do not use mor	gstor adult use ■ do not gi e mo e htdose on hart below. ed dosing up. Do not use any o he dosing re han 4 times a day ■ replace original
alle gic to aspi in. Symptoms may include: III hives III facial swe ling	^{contains} Ibuprofen Oral Suspension (NSAID)	Weight (Ib)	Age (yr)	Dose (mL)*
as hma (wheezing) Shock Skin reddening Ash biste s. If an alle gic	Oral Suspension (NSAID)	under 24	under 2 years	ask a doctor
eaction oc u s, stop use and seek medi a help light away. Stomach bleeding warning: This p oduct contains an NSAID, which may ause severe stoma h	100 mg per 5 mL 4 fl oz	24-35 lbs	2-3 years	5 mL
	Pain Reliever/Fever Reducer (120 mL)	36-47 lbs	4-5 years	7 5 mL
bleeding. The chance is higher if your hid III has had stome hudes so theleding polemis III takes a blood himing (antii cagudart) or steroid dug III takes o he dugs during a sociation or noncrescrip ion NAS Ds kasi in. buo fen. nao own. or	Non-Staining Dye-Free • Original Berry Flavor	48-59 lbs	6-8 years	10 mL
ontaining p escription or nonprescrip ion NSA Ds (aspi in, bup ofen, nap oxen, or or line of the s) at takes more or fo a longer time han directed. Heart attack a d stroke	Non-Staining Dyerriee • Original Berry Plavor	60-71 lbs	9-10 years	12 5 mL
warning: NSAIDs, except aspi in, inc ease the risk o heart attack, heart failure, and	has no been drinking fluids 🔳 child has lost a lot of fluid due to omiting o diar hea 🔳 chi d has	72-95 lbs	11 years	15 mL
4 One These can be teld. The is is higher fyour serve now han directed or for longer han directed. So le all variantific server or sets settors bed or so is had to be all variantific server or sets settors bed or so is had to be all variantific server or sets settors bed or so is had to be all variantific set of the set of th	high blod pressue, hærd folseae, her och nosis, körey dissae, o hed at ske ∎ hå håns sam ∎ i hå tisk ska or dure å kå at doert om hærna ska bled nosis en lit het bå til sunder a doctor's care for an sersons cond fon i til sking any o her dug. When uslag hås p dukt til slav in hörd or mi i försmich upset occur s. Styrup se and ask at doctor i til i hå superincera any för hölloning sjord stranstruktioner. Die blod mission and start of blod stranstruktion at til stores notge better i til håd sa syntoms of heart o delman stra kare. I het spån i til obser hørge better i til håd sa syntoms of heart o delman stra kare. I het spån i til obser hørge heart og men stranstruktion som part o s ded boly i skured speech i ling senling in til hådes notget anv eller tilt nis stag (24 hou s) oft estmet i liver orgingets no se orlats more han 3 døy ill redness o sve ling i presentin	do not use if bottle wrap in Inactive ingredients acesu far p egela inized starch, puri ied wa Questions or comments? Ca	ater, sodium benzoate, sucrose, xa I 1-877-895-3665 (to I-free) or & JOHNSON CONSUMER are Division	is broken or missing id, flavo s, glyce in polyso bate 80, an han gum 215-273-8755 (co lect)

- Ask a doctor before use if stomach bleeding warning applies to your child child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diar hea
 child has high blood pressure, heart disease, liver cir hosis, kidney disease, or had a stroke
- child has asthma
 child is taking a diuretic
- Ask a doctor or pharmacist before use if the
- child is under a doctor's care for any serious condition
 taking any other drug
- When using this product take with food or milk if stomach upset occurs
- Stop use and ask a doctor if
- child experiences any of the following signs of stomach bleeding:
 feels faint
 - vomits blood
- has bloody or black stools
 has stomach pain that does not get better
 child has symptoms of heart problems or stroke:
- chest pain
- crouble breathing
 weakness in one part or side of body
- slurred speech
- leg swelling
- the child does not get any relief w thin first day (24 hours) of treatment
 fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
 any new symptoms appear
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions this product does not contain directions or complete warnings for adult use do not give more than directed

Drug Facts (continued)

- shake well before using
 mL = milliliter
 find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any
- other dosing device.
 if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
 replace original bottle cap to maintain child resistance

 Dosing	Chart

Weight (lb)	Age (yr)	Dose (mL)*		
under 24	under 2 years	ask a doctor		
24-35 lbs	2-3 years	5 mL		
36-47 lbs	4-5 years	7.5 mL		
48-59 lbs	6-8 years	10 mL		
60-71 lbs	9-10 years	12.5 mL		
72-95 lbs	11 years	15 mL		
* or as directed by a doctor				

Other information

each 5 mL contains: sodium 2 mg store between 20-25'C (68-77'F) do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by: JOHNSON & JOHNSON CONSUMER INC. McNeil Consumer Healthcare Division Fort Washington, PA 19034 USA ©J&JCI 2016 www.motrin.com 0000000

Children's Motrin DYE-FREE tontains Ibuprofen Oral Suspension (NSAID)

Non-Staining **Original Berry** Flavor




DYE - FREE





Non-Staining Original Berry Flavor



3 0045-0184-047

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient Purpose (in each 5 mL) Ibup ofen 100 mg (NSAID)*.....Pain relieve *nonsteroidal anti-inflammatory drug ver reducer

Uses temporarily:

 relieves minor aches and pains due to the offu, sore throat, headache and toothache non cold. ■ reduces feve

Warnings Allergy alert: buprofen may cause a seve e allergic reaction, especially in people alle gic to asp rin. Symptoms may include:

■ hives ■ asthma (wheezing) ■ shock ■ skin reddening If an allergic reaction occurs, stop use and seek me be arbitvare

help right away. Stomach bleeding warning: This p oduct contains an

NSAID, which may cause severe stomach bleeding. The chance is higher if your child: has had stomach ulcers or bleeding p oblems takes a blood thinning (anticoagulant) of


ste oid drug takes other drugs containing prescription or

nonprescription NSAIDs (aspirin, bup ofen,

aproxen, or others) takes more or for a longer time than directed Heart attack and stroke warning NSAIDs, except aspi in, increase the risk of heart attack, heart failure, and st oke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, heacache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use if the child has ever had an alle gic reaction to buprofen or any other pain reliever/fever reducer right before or after heart su gery







Important: Read all product information before using, Keep the box fer important information. This product is intended for use in children. Active lag edient(ince h 5 mJ) Purpose Lop den: 00mg(NSAU) Pain relever forer retacter "rontee dial minimitury" dig Uses tempo arity ill relevers mino a heard pains due to the common oid, lu, sore hoat, heada he and too ta bail is coales feer Was ings Winnigs Nin rigs	Children's Motrin®	he gân ul arsu E airyneu sympo om sopser Keëp out of each ol chi d en . n asë of ore dose, ga med cai her phonormata Poison Cont ol Center right away. (800 222-1222). Di echans E hisp oud does not contain di ectio sor complete warming si or adult use E do not gi e mo e hand i det d'asta esta be bersus gin ar militer II ni ght dose on hat telow. If possible, use weight to dose, or hermiseus es aga E. Lus en of verdosed dosing up. Do not use any o he dosing locia. El fresde, adde see eys 6 - Bous 8 do not use more hand times a day E replace original bo fe aptomartian hid esistance Dosing Chart		
	^{contains} Ibuprofen Oral Suspension (NSAID)	Weight (lb)	Age (yr)	Dose (mL)*
ale gc to aspi n. Symptoms may include: II hves II tacal swe ing II as hma (wheezing) II shock II skin reddening II ash II biste s. If an alle gic		under 24	under 2 years	ask a doctor
eaction oc u s, stop use and seek medi a help ight away. Stomach bleeding warning: This p oduct contains an NSAID, which may ause severe stoma h	100 mg per 5 mL 4 fl oz	24-35 lbs	2-3 years	5 mL
bleeding. The chance is higher if your hid: I has had stoma hulce s or bleeding	Pain Reliever/Fever Reducer (120 mL)	36-47 lbs	4-5 years	7 5 mL
p oblems ∎ takes a blood hinning (anti-oagulant) or steroid d ug ∎ takes o he d ugs	Bubble Gum Flavor	48-59 lbs	6-8 years	10 mL
ontaining pescription or nonprescription NSA Ds (aspi in, bup ofen, nap oxen, or or ball of the s) at takes more or fo a longer time han directed. Heart attack a d stroke	δάρριε σαιτί Γιάνοι	60-71 lbs	9-10 years	12 5 mL
	as no been drinking fluids 🔳 child has lost a lot of fluid due to omiting o diar hea 🔳 child has	72-95 lbs	11 years	15 mL
A compared by the few heater news and voming may be sense. Crown in the sense of consistent sense and voming may be sense. Crown in the sense and voming may be sense. Crown in the sense and voming may be sense. Crown in the sense fract and e.g. a compared by choice. Due totake and the sense heat and e.g. a set of a set of the sense fract and e.g. a set of the sense fract and e.g. a set of the sense fract sense fract set. Set of the s	lip blod pressue, heart diesse, lie orch host, k dwy diesse, o had at die ∎ hich has hen m hich klaine guiente, kak a doere orpharmacisk betreus ei hie hoch die under a doctor's care for any serious cond tion ∎ taking any oher die, When using his obsch "bie hie hie hodor mik f stomach upset cours. Stop use and ask a doter 11 m hid generises any hie hodoring signs of stomach diedering "biefs aftin "biefstoms of a professione the hodor signs of stomach upset cours. Stop use and ask a doter 11 m hid stodory. Bieds kotos ≣ his stomach pain had does not ge beet ≡ childhas symptoms of and polemos of side. His stomach pain had does not ge beet ≡ childhas symptoms of and polemos storiks. En his stomach pain had does not ge beet ≡ childhas symptoms of and polemos storiks. His stomach pole taken beet beet might be the stop of the stop of the side of the stop of the sto	∎do ot use if bottle wrap i	mprinted "SAFETY SEAL®" ne polassium, anhydrous c tric ac arch, pur fied water, sodium benz II 1-877-895-3665 (to I-free) o & JOHNSON CONSUMER re Division	: d, FD&C red no. 40, flavo s, glycerin, cate, sucrose, anthan gum r 215-273-8755 (collect)

- Ask a doctor before use if stomach bleeding warning applies to your child child has a history of stomach problems, such as hearthu
- child has drawing of standar prosterior, solar heartbun
 child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids child has lots a lot of fluid due to vomiting or diar hea
 child has high blood pressure, heart disease, liver cir hosis, kidney disease, or had a stroke

child has asthma

- child is taking a diuretic
- Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition

taking any other drug

- When using this product take with food or milk if stomach upset occurs
- Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding:
- feels faint vomits blood
- has bloody or black stools
 has stomach pain that does not get better
 child has symptoms of heart problems or stroke:
- chest pain
- trouble breathing
 weakness in one part or side of body
 slurred speech
- leg swelling
- the child does not get any relief w thin first day (24 hours) of treatment
 fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ this product does not contain directions or complete warnings for adult use do not give more than directed

Drug Facts (continued)

shake well before using
■ mL = milliliter
■ find right dose on chart. If possible, use weight
to dose; otherwise use age.
use only enclosed dosing cup. Do not use any
other dosing device.
if needed, repeat dose every 6-8 hours

 do not use more than 4 times a day
 replace original bottle cap to maintain child resistance

Dosing Chart					
Weight (lb)	Age (yr)	Dose (mL)*			
under 24	under 2 years	ask a doctor			
24-35 lbs	2-3 years	5 mL			
36-47 lbs	4-5 years	7.5 mL			
48-59 lbs	6-8 years	10 mL			
60-71 lbs	9-10 years	12.5 mL			
72-95 lbs	11 years	15 mL			
* or as directed by a doctor					

Other information each 5 mL contains: sodium 2 mg
 store between 20-25°C (68-77°F)
 do not use if carton is opened or if carton tape
 or bottle wrap imprinted "SAFETY SEAL®" is
 broken or missing


Inactive ingredients acesulfame potassium, anhydrous citric acid, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose.xanthan gum

Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by: JOHNSON & JOHNSON CONSUMER INC. McNeil Consumer Healthcare Division Fort Washington, PA 19034 USA ©J&JCI 2016 www.motrin.com 00000000

Children's Motrin buprofen Oral Suspension (NSAID)

Bubble Gum Flavor



^{contains} Ibuprofen Oral Suspension (NSAID) 100 mg per 5 mL

Pain Reliever / Fever Reducer





Bubble Gum Flavor



Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts Active ingredient Purpose (in each 5 mL) bup ofen 100 mg (NSAID)*.....Pain reliever/ *nonste oidal anti-inflammatory drug ever reducer

Uses temporarily:

relieves minor aches and pains due to the common cold, flu, sore th oat, headache and toothache

Warnings Allergy alert: buprofen may cause a seve e allergic reaction, especially in people allergic to aspi in. Symptoms may include:

- thives
 Eacial swelling
 Estimate (wheezing)
 Estimate (whee
- help right away. Stomach bleeding warning: This p oduct dontains an
- NSAID, which may cause severe stomach bleeding. The
- chance is higher if your child:
 has had stomach ulcers or bleeding prob
 takes a blood thinning (anticoagulant) of
- ste oid drug takes other drugs containing prescription or

 takes unler bruge containing prescription on nonprescription NSAIDs (aspirin, buprolen, naproxen, or others)
 takes more or for a longer time than directed
 Heart attack and stroke warning NSAIDs, except aspi in, increase the risk of heart attack, heart failure, and st oke. These can be fatal. The risk is higher if you use more Sore throat warning: Severe or persistent for the directed. Sore throat warning: Severe or persistent fore th oat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor

p omptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. Do not use if the child has ever had an alle gic reaction to

bup ofen or any other pain reliever/fever educer right before or after heart su gery







Important: Read all product information before using. Keep the box for important information. This product is intended for use in children. Active big edient (in ex h 5 ml) Active big edient (in ex h 5 ml) Purpose bup den (in ex h 5 ml) Pain relever fever reducer Trade ediant in memotry of g Uses tempo ariv) = releves minor to g Uses tempo ariv) = releves minor to g Was nigs Vanigs Wa nigs Wa nigs Wa nigs Wa nigs	Children's Motrin°s	her or contact a Poison Cont of C Di ectons his p oduct does not contail han di ected II shake we befor If poss ble, use weight to dose; o hi device. II if needed, epeat dose ef	hisp oduct does not contain di ectio sor complete warni gsfor adult uss∎ do not gi e handi ected ≣ state use beforessong Int = nilitær in ni djrtdosen hartbelow. De sobels use weight does, o hervieus eage useroji wende does useroji verde does useroji wende does useroji verd keixe. Ei meded, epad does evey 6-8 bou s≣ do not use more han 4 timesa day ≣ nepice o ofe ap homentain hid estatave Dosing Chart			
Alle gy allert: bup den may ause a severe a lengt caction, especially n people alle gic to aspin. Symptoms may include: III hives III facial swe ling III as inma (viheezing) III should III shine and dening III ash III she s. If an alle gic	^{contains} Ibuprofen Oral Suspension (NSAID)	Weight (lb)	Age (yr)	Dose (mL)*		
as hma (wheezing) I shock I skin reddening I ash I b iste s. If an alle gic		under 24	under 2 years	ask a doctor		
eaction oc u s, stop use and seek medi a help ight away. Stomach bleeding warning: This p oduct contains an NSAID, which may ause severe stoma h	100 mg per 5 mL 4 fl oz	24-35 lbs	2-3 years	5 mL		
	Pain Reliever/Fever Reducer (120 mL)	36-47 lbs	4-5 years	7 5 mL		
bleeding. The chance is higher if your hid t has had stoma huces or bleeding. p oblems t takes a blood himing fand cagular() or steroid d ug t takes 0 he d ugs or taking no escription or concression in NSA DS (say ii h, buo fan ang one or	Crano Elavor	48-59 lbs	6-8 years	10 mL		
ontaining p escription or nonprescrip ion NSA Ds (aspi in, bup ofen, nap oxen, or o he s) atkes more or fo a longer time han directed. Heart attack a d stroke	Grape Flavor	60-71 lbs	9-10 years	12 5 mL		
warning: NSAIDs, except aspi in, inc ease the risk o heart attack, heart failure, and	has no been drinking fluids E child has lost a lot of fluid due to omiting o diar hea E chi d has	72-95 lbs	11 years	15 mL 🔪		
comparised by high feer, headen, names, and von timp be service. Crossit, doctro pompty. Do not use more han 2 days or administe to children under 3 yeas of age unless directably bolok. Ob not user III to hit has vere had naile gis each to buy often car any be regime isserflever educer III is that has vere had naile give and to buy often car any be regime isserflever educer III is that has vere had naile give a structure buy often car any be regime isserflever educer III is that has vere had naile give the hast sugery Ask a doctor belo e use III is shown bibeding use ing applies to had not be done and the had naile give in the service of the service of the hast sugery Ask a doctor belo e use III is shown bibeding use ing applies to you in the life that has history of some in the potenties, such as keep on III bit Algas?	high blod greessum, haard telesse, keira ih nasis, king dissase, a hafa si kei a lihitas as hme ■ hi di s haking ad wetic. Aak a doctor or pharmacist before use i the chi di si undre a doctor si care for any serious condition. It taking any o her di ugi When using hat p duct "Liste with hold or mit. Homsch uses toccus. Stop use and ask a doctor 11 m. hi degreences any of her blonning sings of sharmable beefing. He list sim is under the sime blood bloody ob back stobs. It has stomach pain hat ches not pe better. ■ chi blas symptoms of herat o dolms or si che. It hespan it m blood beefing in well same nor period so de body to back stobs. It has stomach pain hat ches not pe better. ■ chi blas symptoms of harat o dolms or si che Hespan it m bloot beefing in weakens nor period so de body it sizer speech. ■ leg seeling it he hi does not pet any refir whith is dap (A bou s) of t estimet. ■ leg see ingin. He hind does not pet extense o swe ling is present in	or as directed by a doctor Oher kind main if a each sm. Loontains sodium 2 mg if sore between 20 35 C (68-77 F) Oher kind main if a each sm. Loontains sodium 2 mg if sore between 20 35 C (68-77 F) Image if any solid s				

- Ask a doctor before use if stomach bleeding warning applies to your child child has a history of stomach problems, such as
- child has drawing of standar prosterior, solar heartbun
 child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids child has lots a lot of fluid due to vomiting or diar hea
 child has high blood pressure, heart disease, liver cir hosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic
- Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition
- taking any other drug
- When using this product take with food or milk if stomach upset occurs
- Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding:
 feels faint
- vomits blood
- has bloody or black stools
 has stomach pain that does not get better
 child has symptoms of heart problems or stroke:
- chest pain
- trouble breathing
 weakness in one part or side of body
 slurred speech leg swelling
- the child does not get any relief w thin first day (24 hours) of treatment
 fever or pain gets worse or lasts more than 3 days redness or swelling is present in the painful area any new symptoms appear
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions this product does not contain directions or complete warnings for adult use do not give more than directed

Drug Facts (continued)

shake well before using
mL = milliliter
■ find right dose on chart. If possible, use weight
to dose; otherwise use age.
■ use only enclosed dosing cup. Do not use any
other dosing device.

 if needed, repeat dose every 6-8 hours
 do not use more than 4 times a day
 replace original bottle cap to maintain child resistance

Tesistarice	Dosing Chart				
Weight (lb)	Age (yr)	Dose (mL)*			
under 24	under 2 years	ask a doctor			
24-35 lbs	2-3 years	5 mL			
36-47 lbs	4-5 years	7.5 mL			
48-59 lbs	6-8 years	10 mL			
60-71 lbs	9-10 years	12.5 mL			
72-95 lbs	11 years	15 mL			
* or as directed by a doctor					

Other information each 5 mL contains: sodium 2 mg
store between 20-25'C (68-77'F)
do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing


Inactive ingredients acesulfame potassium, anhydrous citric acid, D&C red no. 33, FD&C blue no.1, FD&C red no. 40, flavors, glycerir polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by: JOHNSON & JOHNSON CONSUMER INC. McNeil Consumer Healthcare Division Fort Washington, PA 19034 USA ©J&JCI 2016 www.motrin.com 00000000

Children's Motrin buprofen Oral Suspension (NSAID)

Grape Flavor



^{contains} Ibuprofen Oral Suspension (NSAID) 100 mg per 5 mL

Pain Reliever / Fever Reducer







Grape Flavor 4 fl oz (120 mL)

0045-0603-40

3



Drug Facts Active ingredient Purpose (in each 5 mL) bup ofen 100 mg (NSAID)*.....Pain reliever/ *nonste oidal anti-inflammatory drug ever reducer

Uses temporarily: relieves minor aches and pains due to the common cold, flu, sore th oat, headache and toothache

Warnings Allergy alert: buprofen may cause a seve e allergic reaction, especially in people allergic to aspi in. Symptoms may include:

- thives
 Eacial swelling
 Estimate (wheezing)
 Estimate (whee
- help right away. Stomach bleeding warning: This p oduct dontains an
- NSAID, which may cause severe stomach bleeding. The
- chance is higher if your child:
 has had stomach ulcers or bleeding prob
 takes a blood thinning (anticoagulant) of
- ste oid drug takes other drugs containing prescription or

 a case other of use containing prescription of nonprescription NSAIDs (aspirin, buprolen, naproxen, or others)
 a takes more or for a longer time than directed
 Heart attack and stroke warning NSAIDs, except aspi in, increase the risk of heart attack, heart failure, and st oke. These can be fatal. The risk is higher if you use more Sore throat warning: Severe or persistent for the directed. Sore throat warning: Severe or persistent fore th oat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor

p omptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. Do not use if the child has ever had an alle gic reaction to

bup ofen or any other pain reliever/fever educer right before or after heart su gery










Important: Read all product information before using. Keep the box for important information. This product is intended for use in children. Advering edint (ine a 5 mL) Purpose Active ing edint (ine a 5 mL) Pain relever for reducer for rouse odd all-of immetry of u Pain relever for reducer for rouse in the immetry of u Uses the more advectors for each one base for each	For HOSPITAL/GOVERNMENT USE ONLY Net for Retail Sale NDC 50580-601-21 Children's Motrino®	help or contact a Poison Cont ol C Di ect ons his p oduct does not contai	enter right away. (* 800 222-1222). n di ectio s or complete warni re using ■ mL = m lifter ■ ind ig erwise use age. ■ use only enclos every 6-8 hou s ■ do not use mor	gs for adult use ∎ do not gi e mo e
Alle gy alert: bup ofen may ause a severe a lergic eaction, especially in people alle gic to aspi in. Symptoms may include: II hives II facial swe ling II as hma (wheezing) III shock III skin reddening III ash III biste s. If an alle gic	contains Ibuprofen Oral Suspension (NSAID)	Weight (lb)	Age (yr)	Dose (mL)*
as hma (wheezing) I shock I skin reddening I ash I biste s. If an alle gic	Oral Suspension (NSAID)	under 24	under 2 years	ask a doctor
eaction oc u s, stop use and seek medi a help light away. Stomach bleeding warning: This p oduct contains an NSAID, which may ause severe stoma h	100 mg per 5 mL	24-35 lbs	2-3 years	5 mL
bleeding. The chance is higher if your hid: I has had stoma huice s or bleeding	Pain Reliever/Fever Reducer 4 fl oz (120 mL)	36-47 lbs	4-5 years	7 5 mL
bleeding. The chance is higher if your hid: It has had stoma h uice s or bleeding p oblems I takes a blood himning (anti cegulant) or stervid ug I takes o he d ugs or takes a blood himning (anti cegulant) or stervid up (anticegulant) or den on or one or takes of the den or takes of the den or takes of the den of takes of the den or takes of the den of takes of the den or takes of the den of takes of takes of takes of takes of the den of takes of take	Original Berry Flavor	48-59 lbs	6-8 years	10 mL
ontaining p escription or nonprescription NSA Ds (aspi in, bup ofen, nap oxen, or the secription of a longer time han directed. Heart attack a d stroke	Original berry Flavor	60-71 lbs	9-10 years	12 5 mL
warning: NSAIDs, except aspi in, inc ease the risk o heart attack, heart failure, and	has no been drinking fluids 🔳 child has lost a lot of fluid due to omiting o diar hea 🔳 chi d has	72-95 lbs	11 years	15 mL
st ole. These can be teld. The sits in biter 'you use more han directed or for longer han directed. So is the all varianticity Server or set sets too hand or sore har a comparied by high lever, headbate, nausea, and vom ling may be serious. Consult doubt promptly. Donot use more than 2 days or admissible to bitter mode 3 yee as 0 agauness directed by door to any one prainne levelfineer advoor. It will be those eardon to buy den or any or brainne levelfineer advoor. It will be those hearts user, Mack addoor their loss at its strong blestop work in an ago to potens or serious side effects of making pain telever ear our is of these potens or serious side effects of making pain telever advoor.	high blod pressue, hard disease, lever in hosis, king disease, i had ast kie ∎ hit has sam ■ hit skie kan durine, kast addurer on patematich bled use use it the all dis ∎ under a doctor's care for any serious cond tion ■ taking any o her dug. When using his p dugt it take with hood rm iik formach upset cours. S Dig use and ask a doctor if ■ hit experimence any of her bloining signs of stransmitheding. It will be in the mission of the blody or blad stock. When some house how the bloin the sign the sign of the signal patient of the sis a signal patient of the	do not use if bottle wrap in Inactive ingredients acesu far flavo s, glyce in polyso bate 80 gum. Questions or comments	pregelatinized star h purified wat ? Cal 1-877-895-3665 (toll-fro & JOHNSON CONSUMER re Division	is broken or missing id, D&C yellow no. 10, FD&C ed no. 40, er, sodium benzoate, sucrose, xan han ee) or 215-273-8755 (co lect)



do not give more than directed

©J&JCI 2016 www.motrin.com 00000000 bup ofen or any other pain reliever/fever educer ■ right before or after heart su gery





N 50580 60121





(b) (4)