Do not use if seal under cap is broken or missing.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Uses ■ **relieves minor aches and pains due to the common cold, flu, sore throat, headache, toothache, backache, muscle aches, menstrual cramps, minor joint pain, arthritis pain** ■ **reduces fever** ■ **reduces inflammation** ■ **reduces swelling associated with sinus congestion and NSAID-induced edema** ■ **reduces swelling associated with inflammation and pain from bursitis, tendinitis, and other soft tissue (muscle, tendon, ligament) disorders** ■ **reduces pain associated with the flare-up of osteoarthritis** ■ **reduces pain associated with the flare-up of rheumatoid arthritis** ■ **reduces pain associated with the flare-up of juvenile rheumatoid arthritis** ■ **reduces pain associated with the flare-up of ankylosing spondylitis** ■ **reduces pain associated with the flare-up of gout** ■ **reduces pain associated with the flare-up of bursitis** ■ **reduces pain associated with the flare-up of tendinitis** ■ **reduces pain associated with the flare-up of other musculoskeletal disorders** ■ **reduces pain associated with the flare-up of other inflammatory disorders** ■ **reduces pain associated with the flare-up of other joint disorders** ■ **reduces pain associated with the flare-up of other soft tissue

Reference ID: 4098075

3 00045-0909-24 4

EXP:

LOT:

See New Warnings

Motrin® Junior Strength

contains Ibuprofen Tablets (NSAID), 100 mg

Pain Reliever / Fever Reducer

24 Grape-Flavored Chewable Tablets

For Ages 2 to 11 Years

Lasts up to 8 hours

NDC 50580-909-24

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**

Dosing Chart

Weight (lb)	Age (yr)	Tablets
under 24	under 2	ask a doctor
24–35	2–3	1
36–47	4–5	1 1/2
48–59	6–8	2
60–71	9–10	2 1/2
72–95	11	3

Other information

- phenylketonurics: contains phenylalanine 2.8 mg per tablet
- store between 20–25°C (68-77°F)
- **do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing**

Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, D&C red no. 7 calcium lake, D&C red no. 30 aluminum lake, FD&C blue no. 1 aluminum lake, flavor, fumaric acid, hydroxyethyl cellulose, hypromellose, magnesium stearate, mannitol, microcrystalline cellulose, povidone, sodium lauryl sulfate, sodium starch glycolate

Important: Read all product information before using. Keep this box for important information. This product is for children only.

Drug Facts

Active ingredient (in each tablet) — Purpose

Ibuprofen 100 mg (NSAID)* ..... Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Drug Facts (continued)

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- mouth or throat burning may occur; give with food or water
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
  - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Drug Facts (continued)

Questions or comments?

Call **1-877-895-3665** (toll-free) or **215-273-9755** (collect)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA

©&℠ JCI 2016

www.motrin.com

US5.320.855

0000000

0000000

0000000

Do not use if stock seal is broken or missing.

Keep out of reach of children.

Read all warnings and directions before use.

Store at 20-25°C (68-77°F).

Tamper Resistant

Motrin®

Junior Strength

(ibuprofen) **Tablets 100 mg** (NSAID)

Fruit Punch Flavored Medicine

24 chewable Tablets; 100 mg each

More than four important formulas.

Ac | **4-hrs relief** | **24-hrs relief** | **Pa** *pain*

Rx: Order 60 mg NSAID QD - Pain relief while or reduce fever.

Uses immediate pain ■ discomfort for ■ minor or minor a has ■ headache, upset stomach or ■ muscle aches ■ backache ■ toothache ■ menstrual cramps ■ common cold. flu, sore throat pain ■ pain due to the common cold, flu, sore throat ■ headache ■ minor aches and pains due to the common cold, flu, sore throat ■ headache ■ toothache ■ muscle aches ■ backache ■ menstrual cramps ■ minor pain ■ reduce fever

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer. ■ stomach bleeding ■ peptic ulcer ■ severe liver disease ■ severe kidney disease ■ severe heart failure ■ **if you are taking a prescription for** ■ **blood thinning drugs** ■ **anticoagulant or anti-platelet drugs** ■ **ulcer or stomach bleeding** ■ **severe liver disease** ■ **severe kidney disease** ■ **severe heart failure**

Reference ID: 4098075

3 0045-0494-245

EXP:

LOT:

See New Warnings

Motrin®

Junior Strength

contains Ibuprofen Tablets (NSAID), 100 mg

Pain Reliever / Fever Reducer

24 Orange-Flavored Chewable Tablets

For Ages 2 to 11 Years

Lasts up to 8 hours

NDC 50580-207-24

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2016
www.motrin.com
US5.320.855

Drug Facts (continued)

Questions or comments?
Call **1-877-895-3655** (toll-free) or **215-273-8755** (collect)

Important: Read all product information before using. Keep this box for important information. This product is for children only.

Drug Facts

Active ingredient (in each tablet) — **Purpose**
Ibuprofen 100 mg (NSAID)* ..... Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses
temporarily:
- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:
- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Drug Facts (continued)

Do not use
- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if
- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is
- under a doctor's care for any serious condition
- taking any other drug

When using this product
- mouth or throat burning may occur; give with food or water
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if
- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
  - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800 222-1222)

Directions
- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**

Dosing Chart

Weight (lb)	Age (yr)	Tablets
under 24	under 2	ask a doctor
24–35	2–3	1
36–47	4–5	1 1/2
48–59	6–8	2
60–71	9–10	2 1/2
72–95	11	3

Other information
- phenylketonurics: contains phenylalanine 2.8 mg per tablet
- store between 20–25°C (68-77°F)
- **do not use if neck wrap or foil inner seal imprinted "Safety Seal®" is broken or missing**

Inactive ingredients
acesulfame potassium, anhydrous citric acid, aspartame, FD&C yellow no. 6 aluminum lake, flavor, fumaric acid, hydroxyethyl cellulose, hypromellose, magnesium stearate, mannitol, microcrystalline cellulose, povidone, sodium lauryl sulfate, sodium starch glycolate

0000000

0000000

0000000