



**Drug Facts (continued)**

**Warnings**  
**Allergy alert** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child: has had stomach ulcers or bleeding problems; takes a blood thinning (anticoagulant) or steroid drug; takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); takes more or for a longer time than directed.  
**Heart attack and stroke warning** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**Sore throat warning** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.  
**Do not use**  
 if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer right before or after heart surgery.  
**Ask a doctor before use if**  
 stomach bleeding warning applies to your child; child has a history of stomach problems, such as heartburn; child has problems or serious side effects from taking pain relievers or fever reducers; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke; child has asthma; child is taking a diuretic.  
**Ask a doctor or pharmacist before use if the child is**  
 under a doctor's care for any serious condition; taking any other drug.  
**When using this product**  
 take with food or milk if stomach upset occurs.  
**Stop use and ask a doctor if**  
 child experiences any of the following signs of stomach bleeding: feels faint; vomits blood; has bloody or black stools; has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling; the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear.

**Drug Facts (continued)**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)  
**Directions**  
 This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. Find right dose on chart below. If possible, use weight to dose; otherwise use age.  
 mL = milliliter. Measure with the dosing device provided. Do not use with any other device. Dispense liquid slowly into the child's mouth, toward the inner cheek. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day.  

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL








**Other information**  
 Store between 20-25°C (68-77°F). Do not use if carton top is imprinted "SAFETY SEAL" or bottle wrap is imprinted "SAFETY SEAL" and "USE WITH ENCLOSED DOSING DEVICE ONLY" is broken or missing.  
**Inactive ingredients** anhydrous citric acid, carmellose, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum.  
**Questions or comments?**  
 Call 1-877-895-3665 (toll-free) or 215-273-8755 (colect).

See New Warnings  
 NDC 50580-100-18  
**For Ages 6 Mos. to 23 Mos.**  
**Concentrated Motrin Infants' Drops**  
 contains Ibuprofen Oral Suspension (NSAID)  
 50 mg per 1.25 mL  
**Pain Reliever/Fever Reducer**  
**Lasts up to 8 hours**  
 Use only with enclosed syringe.  
**Original Berry Flavor**  
 1/2 fl oz (15 mL)



LOT: \_\_\_\_\_  
 EXP: \_\_\_\_\_  
 Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**  
 McNeil Consumer Healthcare Division  
 Fort Washington, PA 19034 USA ©J&JCI 2016  
 www.motrin.com

USA  
 000000  
 891



(b) (4)



**Drug Facts**  
**Active ingredient (in each 1.25 mL)**  
 Ibuprofen 50 mg (NSAID)  
**Purpose**  
 Pain reliever/fever reducer  
**Uses**  
 Temporarily relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches  
 Relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

See New Warnings  
 NDC 50580-198-50  
**For Ages 6 Mos. to 23 Mos.**  
**Concentrated Motrin® Infants' Drops**  
**DYE-FREE**  
 contains Ibuprofen Oral Suspension (NSAID)  
 50 mg per 1.25 mL  
**Pain Reliever/Fever Reducer**  
**Lasts up to 8 hours**  
 Use only with enclosed syringe.  
**Non-Staining Original Berry Flavor**  
**1 fl oz (30 mL)**

**Concentrated Motrin® Infants' Drops**  
**DYE-FREE**  
**Concentrated Motrin® Infants' Drops**  
 contains Ibuprofen Oral Suspension (NSAID)  
 50 mg per 1.25 mL  
**Pain Reliever/Fever Reducer**  
**Non-Staining Original Berry Flavor**  
**1 fl oz (30 mL)**  
 Actual Size

**Drug Facts (continued)**  
**Warnings**  
**Allergy alert** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 hives, facial swelling, rash, blisters, wheezing, shock, skin redness, dizziness, difficulty breathing, stomach pain, or vomiting.  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:  
 has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.  
**Heart attack and stroke warning** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**Sore throat warning** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.  
**Do not use**  
 if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer right before or after heart surgery.  
**Ask a doctor before use if**  
 stomach bleeding warning applies to your child, child has a history of stomach problems, such as heartburn, child has problems or serious side effects from taking pain relievers or fever reducers, child has not been drinking fluids, child has lost a lot of fluid due to vomiting or diarrhea, child has high blood pressure, heart disease, liver cirrhosis, kidney disease or had a stroke, child has asthma, child is taking a diuretic.  
**Ask a doctor or pharmacist before use if the child is**  
 under a doctor's care for any serious condition, taking any other drug.  
**When using this product** take with food or milk if stomach upset occurs.  
**Stop use and ask a doctor if**  
 child experiences any of the following signs of stomach bleeding: feels faint, vomits blood, has bloody or black stools, has stomach pain that does not get better, child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, child does not get any relief within first day (24 hours) of treatment, fever or pain gets worse or lasts more than 3 days, dizziness or swelling is present in the painful area, any new symptoms appear.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (800) 222-1222.  
**Directions**  
 this product does not contain directions or complete warnings for adult use

**Drug Facts (continued)**  
 do not give more than directed, shake well before using, find right dose on chart below. If possible, use weight to dose, or otherwise use age, mL = milliliter, measure with the dosing device provided. Do not use without any of the device, dispense liquid slowly into the child's mouth, lower to the inner cheek, needed, repeat dose every 6-8 hours, do not use more than 4 times a day.  
**Dosing Chart**

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
8-23 lbs	12-23 mos	1.875 mL

**Other information**  
 store below 20-25°C (68-77°F), do not use if carton is opened or carton tape imprinted "SAFETY SEAL" or bottle wrap imprinted "SAFETY SEAL" and "USE WITH ENCLOSED DOSING DEVICE ONLY" is broken or missing.  
**Inactive ingredients** anhydrous citric acid, caramel, levorotatory glycerin, polyorbital 80, pregelatinized starch, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum.  
**Questions or comments?** Call 1-877-895-3665 (toll-free) or 215-273-8755 (no text).  
 3 0045-0198-013

LOT: \_\_\_\_\_  
 EXP: \_\_\_\_\_  
 Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**  
 McNeil Consumer Healthcare Division  
 Fort Washington, PA 19034 USA ©J&JCI 2016  
 www.motrin.com

USA  
 000000  
 887

(b) (4)

54,0 mm 111,0 mm 57,0 mm

51,0 mm

**THE DOSING CHART AND DIRECTIONS APPEARING ON THE FIRST AND LAST PAGE ARE REPEATED FOR YOUR CONVENIENCE. PLEASE READ AND KEEP ALL WARNINGS AND DIRECTIONS TO ENSURE THE PROPER USE OF THIS CONCENTRATED INFANTS' MOTRIN® PRODUCT.**

**Important** Read all product information before using. Keep the box for important information. This product is intended for use in children ages 6 months to 23 months of age.

**Active ingredient (in each 1.25 mL)**  
 ibuprofen 50 mg (NSAID)\* Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Purpose**  
 ■ reduces fever  
 ■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Directions** ■ this product does not contain directions or complete warnings for adult use  
 ■ do not give more than directed  
 ■ shake well before using  
 ■ find right dose on chart below. If possible, use weight to dose; otherwise use age. ■ mL = milliliter  
 ■ measure with the dosing device provided. Do not use with any other device. ■ dispense liquid slowly into the child's mouth, toward the inner cheek. ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day

**Concentrated Motrin® Infants' Drops**  
 contains ibuprofen Oral Suspension (NSAID)  
 50 mg per 1.25 mL  
 Pain Reliever/Fever Reducer  
 Non-Staining Original Berry Flavor  
 1 fl oz (30 mL)

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

**Page 2** **Page 7**

57,0 mm 111,0 mm 54,0 mm

51,0 mm

**!!! OPEN to read all Warnings and Directions before use**

**Directions** ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed ■ shake well before using ■ find right dose on chart below. If possible, use weight to dose; otherwise use age. ■ mL = milliliter ■ measure with the dosing device provided. Do not use with any other device. ■ dispense liquid slowly into the child's mouth, toward the inner cheek. ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day

**Concentrated Motrin® Infants' Drops**  
 NDC 50580-198-50  
 contains ibuprofen Oral Suspension (NSAID)  
 50 mg per 1.25 mL  
 Pain Reliever/Fever Reducer  
 Non-Staining Original Berry Flavor  
 1 fl oz (30 mL)

**Glue Area (glued to base label if only 7 pages needed)**

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

**Page 1 (COVER)**

54,0 mm 108,0 mm 54,0 mm

51,0 mm

**Do not use**  
 ■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer  
 ■ right before or after heart surgery

**Ask a doctor before use if**  
 ■ stomach bleeding warning applies to your child  
 ■ child has a history of stomach problems, such as heartburn  
 ■ child has problems or serious side effects from taking pain relievers or fever reducers  
 ■ child has not been drinking fluids  
 ■ child has lost a lot of fluid due to vomiting or diarrhea  
 ■ child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke  
 ■ child has asthma ■ child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**  
 ■ under a doctor's care for any serious condition ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  
 ■ child experiences any of the following signs of stomach bleeding:  
 ■ feels faint ■ vomits blood ■ has bloody or black stools  
 ■ has stomach pain that does not get better  
 ■ child has symptoms of heart problems or stroke: ■ chest pain  
 ■ trouble breathing ■ weakness in one part or side of body  
 ■ slurred speech ■ leg swelling  
 ■ the child does not get any relief within first day (24 hours) of treatment  
 ■ fever or pain gets worse or lasts more than 3 days  
 ■ redness or swelling is present in the painful area ■ any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** ■ this product does not contain directions or complete warnings for adult use  
 ■ do not give more than directed

**Dosing Chart**

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

**Page 4** **Page 5**

54,0 mm 108,0 mm 54,0 mm

51,0 mm

■ measure with the dosing device provided. Do not use with any other device.  
 ■ dispense liquid slowly into the child's mouth, toward the inner cheek  
 ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day

**Other information**  
 ■ store between 20-25°C (68-77°F)  
 ■ do not use if bottle wrap imprinted "SAFETY SEAL®" and "USE WITH ENCLOSED DOSING DEVICE ONLY" is broken or missing

**Inactive ingredients** anhydrous citric acid, caramel, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

**Questions or comments?**  
 Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
 McNeil Consumer Healthcare Division  
 Fort Washington, PA 19034 USA  
 ©J&JCI 2016 www.motrin.com 00000000

**Warnings**  
**Allergy alert** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.  
 Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:  
 ■ has had stomach ulcers or bleeding problems  
 ■ takes a blood thinning (anticoagulant) or steroid drug  
 ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ takes more or for a longer time than directed

**Heart attack and stroke warning** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for a longer time than directed.

**Sore throat warning** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Page 6** **Page 3**

13,0 mm 90,5 mm 57,0 mm 20,5 mm

51 mm < 9,5 mm

**Cover Label FRONT**

LOT: 000000 EXP:

20,5 mm 90,5 mm 57,0 mm 13,0 mm

51 mm < 9,5 mm

**Cover Label BACK**

54,0 mm 74,5 mm 20,5 mm

51,0 mm

**Base Label**



(b) (4)