

00000000



*Mint Flavored Tablets*

**18 Chewable**

Relief of diarrhea plus:  
Gas Cramps Pressure Bloating

**Imodium**  
Anti-Gas

Loperamide HCl, 2 mg/ Simethicone, 125 mg An

**MULTI-SYMPTOM RELIEF**

See New Warning and Direction

NDC 50880-339-18

00000000

00000000

**Imodium** Loperamide HCl, 2mg/  
MULTI-SYMPTOM RELIEF Simethicone, 125mg  
Antidiarrheal/Anti-Gas Chewable

*The Makers of Imodium*  
do not manufacture store brands.

PATENTED FORMULA

XXXXXX

|  |   |
|--|---|
| <b>Drug Facts</b>                          | <b>Drug Facts (continued)</b>   |
| <b>Active ingredients (in each tablet)</b> | <b>Uses</b>   |
| Loperamide HCl 2 mg.....Anti-diarrheal     | relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas |
| Simethicone 125 mg.....Anti-gas            |   |

Distributed by: AMNUL Consumer Healthcare  
 FC03 WASHINGTON, PA 15391 USA  
 © MCH-PC, INC. 2018 www.imodium.com  
 U.S. Patent No. 5,489,436

**Imodium**  
 MULTI-SYMPTOM RELIEF  
 Loperamide HCl, 2mg/  
 Simethicone, 125mg  
 Antidiarrheal/Anti-Gas  
 Chewable

|   |  |  |
|---|--|--|
| <b>Drug Facts (continued)</b>   | <b>Drug Facts (continued)</b>  | <b>Drug Facts (continued)</b>  |
| <b>Warnings</b>   | <b>Allergy alert:</b>  | find right dose on chart below. If possible, use weight to dose; otherwise use age.  |
| Do not use if you have bloody or black stool  | Do not use if you have ever had a rash or other allergic reaction to loperamide HCl  | adults and children 12 years and over  |
| Ask a doctor before use if you have   | Heart alert: Taking more than directed can cause serious heart problems or death   | chew 2 tablets and take with water after the first loose stool; chew 1 tablet and take with water after each subsequent loose stool; but no more than 4 tablets in 24 hours  |
| Ask a doctor or pharmacist before use if you are taking antibiotics   | Do not use if you have bloody or black stool   | children 9-11 years (60-95 lbs)  |
| When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery. | Ask a doctor before use if you have  | chew 1 tablet and take with water after the first loose stool; chew 1/2 tablet and take with water after each subsequent loose stool; but no more than 3 tablets in 24 hours |
| Stop use and ask a doctor if  | fever  | children 6-8 years (48-59 lbs)   |
| diarrhea lasts for more than 2 days   | mucus in the stool   | chew 1 tablet and take with water after the first loose stool; chew 1/2 tablet and take with water after each subsequent loose stool; but no more than 2 tablets in 24 hours |
| you get abdominal swelling or bulging. These may be signs of a serious condition.                                     | a history of liver disease   | children 2-5 years (34 to 47 lbs)  |
| If pregnant or breast-feeding, ask a health professional before use.  | Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) | ask a doctor   |
| <b>Directions</b>   | Directions   | children under 2 years (up to 33 lbs)  |
| drink plenty of clear fluids to help prevent dehydration caused by diarrhea   | do not use   |  |



-0202-185

|  |   |
|--|---|
| <b>Drug Facts (continued)</b>                        | <b>Drug Facts (continued)</b>                             |
| <b>Other information</b>                             | <b>Questions or comments?</b>                             |
| each tablet contains: calcium 50 mg                  | call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish) |
| do not use if carton or blister unit is open or torn |   |
| store between 20-25 C (68-77 F)                      |   |
| see side panel for expiration date and lot number    |   |
| <b>Inactive ingredients</b> cellulose acetate,       |   |

-----  
**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**  
-----

/s/  
-----

VALERIE S PRATT  
04/12/2017