



Front



Back



Right Side View

Left Side View



Top View



Bottom View



Front




Back



Right Side View

Left Side View



Top View



Bottom View



Varnish free area  Reserved area 




(b) (4)



Bar prints 100mm @ 100%



Varnish free area  Reserved area 



(b) (4)

Bar prints 100mm @ 100%

VARIABLE DATA LINE 1
VARIABLE DATA LINE 2

RHINOCORT[®] ALLERGY SPRAY

Budesonide (budesonide)
Nasal spray, 32 mcg per spray
Nasal Allergy Symptom Reliever

Original Prescription Strength
Active ingredient made in Netherlands.
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©2016 Johnson & Johnson Consumer Inc.

60 SPRAYS 0.169 fl oz (5 mL)

Questions? call 1-800-343-7805 (toll-free) or 215-273-8755 (collect)

RHINOCORT[®] is a licensed trademark of AstraZeneca AB

children under 8 years of age **Do not use**

Do not use more than directed ■ If you forget a dose, do not double the next dose. ■ Do not spray into eyes or mouth. ■ If allergy symptoms do not improve after two weeks, stop using and talk to a doctor. ■ Do not use for the common cold. ■ Shake well before each use.

Other information ■ Do not use if aluminum seal under unit cap and spray tip is damaged/broken/unsealed ■ Keep package and insert. They contain important information. ■ Store at 20°-25° C (68°-77° F)

Inactive ingredients carbonylmethyl cellulose sodium, decazone anhydrous, disodium edetate, hydrochloric acid (for pH adjustment), microcrystalline cellulose, polyorbata 80, potassium sorbate, purified water

Questions? call 1-800-343-7805 (toll-free) or 215-273-8755 (collect)

PS08490-01
Keep package and insert. They contain important information.

RHINOCORT[®] ALLERGY SPRAY

Budesonide (budesonide)
Nasal spray, 32 mcg per spray
Nasal Allergy Symptom Reliever

60 SPRAYS 0.169 fl oz (5 mL)

PEEL BACK TAB AT ARROW TO OPEN

Active ingredient (in each spray) Purpose
Budesonide
Budesonide (budesonide) 32 mcg. ■ Nasal allergy symptom reliever
Does temporarily relieve these symptoms of hay fever or other upper respiratory allergies:
■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose
■ itchy throat

Do not use ■ In children under 8 years of age ■ If you have ever had an allergic reaction to any of the ingredients. Ask a doctor before use if you ■ Have had recent nasal ulcers or nose surgery ■ Have had a nose

injury that has not healed ■ are using a steroid medicine for asthma, allergies or skin rash ■ have an eye infection ■ have or had glaucoma or cataracts. **When using this product** ■ the growth rate of some children may be slower ■ some symptoms may get better on the first day of treatment. It may take up to two weeks of daily use to feel the most symptom relief. ■ do not share this bottle with anyone else as this may spread germs ■ remember to tell your doctor about all medicines you take, including this one

Stop use and ask a doctor if ■ you have, or come into contact with someone who has chickenpox, measles or tuberculosis ■ you have or develop symptoms of an infection such as persistent fever ■ you have any change in vision ■ you have severe or frequent nosebleeds ■ if pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions Read instructions (inside package) on how to: ■ get a new bottle ready (prime) before first use ■ prime bottle again if not used for two days ■ use the spray ■ clean the spray nozzle

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

adults and children 12 years of age and older ■ once daily, spray 2 times into each nostril while sniffing gently ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 6 TO UNDER 12 YEARS OF AGE

■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

children 6 to under 12 years of age ■ in adult dosed supervised use ■ once daily, spray 1 time into each nostril while sniffing gently ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.



(b) (4)



Bar prints 100mm @ 100%

RHINOCORT, ALLERGY SPRAY

Budesonide (budesonide)

Nasal spray, 32 mcg per spray
Nasal Allergy Symptom Reliever

120 SPRAYS 0.285 fl oz (8.43 mL)

Questions? call 1-800-343-7805 (toll-free) or 215-273-8755 (collect)

Original Prescription Strength

Active ingredient made in Netherlands.

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©2019 Johnson & Johnson Consumer Inc.

RHINOCORT is a licensed trademark of AstraZeneca AS.

children under 6 years of age **Do not use**

Do not use more than directed If you forget a dose, do not double the next dose. Do not spray into eyes or mouth. If allergy symptoms do not improve after two weeks, stop using and talk to a doctor. Do not use for the common cold. Shake well before each use.

Other information Do not use if aluminum seal under end cap and spray tip is damaged/broken/unsealed. Keep package and insert. They contain important information. Store at 20°-25° C (68°-77° F).

Inactive ingredients carbonylmethyl cellulose sodium, dextrose anhydrous, disodium edetate, hydrochloric acid (for pH adjustment), microcrystalline cellulose, polysorbate 80, potassium sorbate, purified water.

Questions? call 1-800-343-7805 (toll-free) or 215-273-8755 (collect)

RX 3050-646-01

RHINOCORT, ALLERGY SPRAY

Budesonide (budesonide)

Nasal spray, 32 mcg per spray
Nasal Allergy Symptom Reliever

120 SPRAYS 0.285 fl oz (8.43 mL)

PEEL BACK TAB AT ARROW TO OPEN

Active ingredient (in each spray) Purpose
Budesonide (budesonide) 32 mcg... Nasal allergy symptom reliever
Use temporarily to relieve these symptoms of hay fever or other upper respiratory allergies:
nasal congestion runny nose sneezing itchy nose
Itching

Do not use in children under 6 years of age if you have ever had an allergic reaction to any of the ingredients. Ask a doctor before use if you have had recent nose ulcers or nose surgery have had a nose

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

adults and children 12 years of age and older once daily, spray 2 times into each nostril while sniffing gently once your allergy symptoms improve, reduce to 1 spray in each nostril per day.

CHILDREN 6 TO UNDER 12 YEARS OF AGE

children 6 to under 12 years of age once daily, spray 1 time into each nostril while sniffing gently once your allergy symptoms improve, reduce to 1 spray in each nostril per day.


CHILDREN UNDER 6 YEARS OF AGE

children under 6 years of age Do not use. In children under 6 years of age, the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

Do not use if you have ever had an allergic reaction to any of the ingredients. Ask a doctor before use if you have had recent nose ulcers or nose surgery have had a nose

injury that has not healed are using a steroid medicine for asthma, allergies or skin rash have an eye infection have or had glaucoma or cataracts. **When using this product** the growth rate of some children may be slower. Some symptoms may get better on the first day of treatment. It may take up to two weeks of daily use to feel the most symptom relief. Do not share this bottle with anyone else as this may spread germs. Remember to tell your doctor about all medicines you take, including this one. **Stop use and ask a doctor if** you have, or come into contact with someone who has chickenpox, measles or tuberculosis. you have or develop symptoms of an infection such as persistent fever. you have any change in vision. you have severe or frequent nosebleeds. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-322-1222)

Directions Read instructions (inside package) on how to use. Get a new bottle ready (prime) before first use. Prime bottle again if not used for two days. Use the spray. Clean the spray nozzle





1. IMPORTANT INFORMATION


1. Read entire insert before using RHINOCORT® Allergy Spray for complete instructions on how to get a bottle ready for use (primed), how to use the spray bottle, and how to clean the spray nozzle.
2. Keep this insert as it contains important information.



2. STEPS TO GET A BOTTLE READY FOR USE (PRIMED)

Before first use, a new bottle must be primed. To prime RHINOCORT® Allergy Spray:

1. Shake bottle gently.
2. Remove protective cap as shown in Figure A.
3. Hold bottle firmly, as shown in Figure B.



4. Prime the pump by quickly and firmly pressing down on the white collar while holding the base of the bottle with your thumb for 8 sprays. The pump is now ready to use.
5. a. If not used for two consecutive days, reprime the pump until a fine mist appears.
- b. If not used for more than 14 days, rinse the spray tip of the pump using the cleaning steps listed in Section 4. After cleaning, reprime until a fine spray appears.


3. INSTRUCTIONS FOR USE

Follow these instructions for daily use of RHINOCORT® Allergy Spray:

1. Gently blow nose to clear nostrils.
2. Shake bottle gently and remove cap.
3. Hold bottle firmly. (See Figure B)

Insert spray tip into nostril (tip should not reach far into nose). Close other nostril with a finger and lean head slightly forward so the spray will aim toward the back of nose. (See Figure C)

DO NOT spray towards nasal septum (wall between two nostrils). (See Figure D)



DO NOT spray towards nasal septum (wall between two nostrils)

4. For each spray, press down quickly and firmly on white collar while holding the base of the bottle with your thumb. Breathe gently inward.
5. After spraying into your nostril, lean your head slightly backward for a few seconds.
6. If a second spray is needed in the same nostril, repeat steps 3 through 5.
7. Repeat steps 3 through 6, for the other nostril.
8. Avoid blowing your nose for 15 MINUTES after using RHINOCORT® Allergy Spray.
9. Wipe spray tip with a clean tissue (See Figure E), and replace outer cap. Store the bottle in an upright position.



TABLE 1: DIRECTIONS FOR USE

- Before first use, you must get a new bottle ready (primed). See Section 2.
- If bottle is not used for two consecutive days, reprime the pump until a fine spray appears.
- If bottle is not used for more than two weeks, rinse the spray tip of the pump using the cleaning steps listed in Section 4. After cleaning, reprime the pump until a fine spray appears.

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- | | |
|---|--|
| adults and children 12 years of age and older | <ul style="list-style-type: none"> ■ once daily, spray 2 times into each nostril while sniffing gently ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day |
|---|--|

CHILDREN 6 TO UNDER 12 YEARS OF AGE

- the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

- | | |
|-------------------------------------|--|
| children 6 to under 12 years of age | <ul style="list-style-type: none"> ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day. |
|-------------------------------------|--|


- | | |
|-------------------------------|--|
| children under 6 years of age | <ul style="list-style-type: none"> ■ do not use |
|-------------------------------|--|

Other important information for use

- do not use more than directed
- if you forget a dose, do not double the next dose
- do not spray into eyes or mouth
- if allergy symptoms do not improve after two weeks, stop using and talk to a doctor
- do not use for the common cold
- shake well before each use

4. HOW TO CLEAN YOUR RHINOCORT® ALLERGY SPRAY

Rinse cap and spray tip regularly. To do this:



1. Remove cap and gently pull spray tip away from the bottle. (See Figure F)
2. Wash **only cap and spray tip** in warm water and rinse in cold tap water. Shake to remove excess water.
3. Allow cap and spray tip to air dry completely before reassembling the nasal spray.
4. Press and release spray nozzle until a fine spray is produced, taking care not to spray in the face.
5. **If the spray tip becomes blocked, it can be cleared by repeating steps 1 through 3. Do not unblock the nasal applicator with a pin or other sharp object.**

INDOOR ALLERGEN TIPS

From the makers of RHINOCORT® Allergy Spray

Filter your air.
HEPA (High Efficiency Particulate Air) filters can be used in air conditioning systems and vacuums to trap small allergen particles.

Minimize dust mite matter.
Use hypoallergenic bedding with zippered dust-proof covers. Wash sheets and blankets in hot water each week.

Control pets.
Keep pets off of upholstered furniture and out of the bedrooms. It may also help to have your pets washed often to help control dander.

OUTDOOR ALLERGEN TIPS

From the makers of RHINOCORT® Allergy Spray

Keep outdoor allergens out.
On high-pollen days, keep doors and windows closed. Shower and change your clothes after you've spent time outdoors. Also, drive with the windows closed.

Stay off the grass.
If possible, exercise on asphalt or cement, at the beach or at the gym to reduce contact with pollen from grass and weeds.

Dry clothes inside.
Don't hang your clothing or linens outside to dry. Drying them inside helps avoid getting pollen on your clean laundry.

For additional information
about RHINOCORT® Allergy Spray, please call the McNeil Consumer Care Center at 1-800-343-7805 (toll-free) or 215-273-8755 (collect)

Active Ingredient Made in Netherlands. Distributed by: Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division Fort Washington, PA 19034 USA
RHINOCORT® is a licensed trademark of AstraZeneca AB.
©2016 Johnson & Johnson Consumer Inc.


Text area for artwork



(b) (4)



Bar prints 100mm @ 100%



(b) (4)



Bar prints 100mm @ 100%