


DRUG FACT FORMATTING: MODIFIED		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	



LOT/EXP AREA



Distributed by GSK Consumer Healthcare  
Warren, NJ 07059 ©2017 GSK or its licensor  
16/274-50 Visit us at www.excedrin.com



See new warnings information  
**EXCEDRIN**  
**MIGRAINE**  
Acetaminophen, Aspirin (NSAID) and Caffeine  
Pain Reliever/Pain Reliever Aid  
50 CAPLETS\*  
NDC 0067-2039-50

Glue



TAMPER-EVIDENT BOTTLE  
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH  
"SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

**Drug Facts**  
**Active ingredients (in each caplet)**  
Acetaminophen 250 mg.....Pain reliever  
Aspirin 250 mg (NSAID)\*.....Pain reliever  
Caffeine 65 mg.....Pain reliever aid  
\*nonsteroidal anti-inflammatory drug

**Use** • treats migraine

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
• skin reddening • blisters • rash  
If a skin reaction occurs, stop use and seek medical help right away.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
• more than 2 caplets in 24 hours, which is the maximum daily amount  
• with other drugs containing acetaminophen  
• 3 or more alcoholic drinks every day while using this product  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older

**Drug Facts (continued)**  
• have had stomach ulcers or bleeding problems  
• take a blood thinning (anticoagulant) or steroid drug  
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
• have 3 or more alcoholic drinks every day while using this product  
• take more or for a longer time than directed  
**Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month.  
**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

**Do not use**  
• if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer  
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

**Ask a doctor before use if**  
• you have never had migraines diagnosed by a health professional  
• you have a headache that is different from your usual migraines  
• you have the worst headache of your life  
• you have fever and stiff neck  
• you have headaches beginning after or caused by head injury, exertion, coughing or bending  
• you experienced your first headache after the age of 50  
• you have daily headaches  
• you have a migraine so severe as to require bed rest  
• you have liver disease  
• stomach bleeding warning applies to you  
• you have a history of stomach problems, such as heartburn  
• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Drug Facts (continued)**  
• you are taking a diuretic • you have asthma  
• you have problems or serious side effects from taking pain relievers or fever reducers  
• you have vomiting with your migraine headache

**Ask a doctor or pharmacist before use if you are**  
• taking a prescription drug for:  
• diabetes • gout • arthritis  
• under a doctor's care for any serious condition  
• taking any other drug  
• taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

**Stop use and ask a doctor if**  
• an allergic reaction occurs. Seek medical help right away.  
• you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools  
• have stomach pain that does not get better  
• your migraine is not relieved or worsens after first dose  
• new or unexpected symptoms occur  
• ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.**  
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** • do not use more than directed  
• adults: take 2 caplets with a glass of water  
• if symptoms persist or worsen, ask your doctor  
• do not take more than 2 caplets in 24 hours, unless directed by a doctor  
• under 18 years of age: ask a doctor

**Drug Facts (continued)**  
**Other information**  
• store at room temperature 20°-25°C (68°-77°F)  
• use cap lightly after use  
• read all product information before using. Keep this box for important information.

**Inactive ingredients** benzoic acid, carnaliba wax, FD&C Blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polyvinylpyrrolidone, polyethylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

**Questions? 1-800-452-0851**

DRUG FACT FORMATTING- MODIFIED		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.3 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	



DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	



DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

LOT/EXP AREA



Distributed by GSK Consumer Healthcare, Warren, NJ 07059 ©2017 GSK group of companies or its licensor. All rights reserved. 16/274-200 Visit us at www.excedrin.com

46170720

CODE 128 FPO

46170720

CODE 128 FPO

See new warnings information

NDC 0067-2039-92

EXCEDRIN

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine Pain Reliever/Pain Reliever Aid

MIGRAINE Acetaminophen, Aspirin (NSAID) and Caffeine Pain Reliever/Pain Reliever Aid

EXCEDRIN

See new warnings information

See new warnings information

NDC 0067-2039-92

TAMPER-EVIDENT BOTTLE DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet) Purposes Acetaminophen 250 mg Pain reliever Aspirin 250 mg (NSAID) Pain reliever Caffeine 65 mg Pain reliever aid \*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away. Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 caplets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older

• have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

• have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if • you have never had migraines diagnosed by a health professional

Drug Facts (continued)

• you have a headache that is different from your usual migraines • you have the worst headache of your life • you have fever and stiff neck • you have headaches beginning after or caused by head injury, exertion, coughing or bending • you experienced your first headache after the age of 50 • you have daily headaches • you have a migraine so severe as to require bed rest • you have liver disease • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have problems or serious side effects from taking pain relievers or fever reducers • you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

• taking a prescription drug for: • diabetes • gout • arthritis • under a doctor's care for any serious condition • taking any other drug • taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if

• an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • your migraine is not relieved or worsens after first dose • new or unexpected symptoms occur • ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed

• adults: take 2 caplets with a glass of water • if symptoms persist or worsen, ask your doctor • do not take more than 2 caplets in 24 hours, unless directed by a doctor • under 18 years of age: ask a doctor

Other information • store at controlled room temperature 20°-25°C (68°-77°F) • close cap tightly after use • read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

Questions? 1-800-468-7746

Glue

46170720

CODE 128 FPO

46170720

CODE 128 FPO

Table with 3 columns: DRUG FACTS TEXT, TYPE SIZE, TYPE STYLE. Rows include DRUG FACTS TITLE (9 pt, Helvetica Cond Bold Obli), DRUG FACTS CONT. (8 pt, Helvetica Cond Bold Obli), HEADINGS (8 pt, Helvetica Cond Bold), SUBHEADS (6 pt, Helvetica Cond), BODY TEXT (6 pt, Helvetica Cond), TRACKING (0), CONDENSATION (83%), LEADING (6.5 pt), BARLINES, HAIRLINES (1.5, 0.5 pt), CHARACTERS/INCH (<39), BULLETS (5 pt).

LOT/EXP AREA



Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
©2017 GSK group of companies or its licensor. All rights reserved.  
167274-250 Visit us at www.excedrin.com



46170647



46170647



See new warnings information

NDC 0067-2039-77

# EXCEDRIN

## MIGRAINE

Acetaminophen,  
Aspirin (NSAID) and Caffeine  
Pain Reliever/Pain Reliever Aid

**25%  
MORE  
FREE**

**250  
200**  
CAPLETS<sup>™</sup>  
\*CAPSULE-SHAPED TABLETS

See new warnings information

# EXCEDRIN

MIGRAINE  
Acetaminophen, Aspirin (NSAID)  
and Caffeine  
Pain Reliever/Pain Reliever Aid

**250  
200**  
CAPLETS<sup>™</sup>  
\*CAPSULE-SHAPED TABLETS

**25%  
MORE  
FREE**

See new warnings information

# EXCEDRIN

MIGRAINE  
Acetaminophen, Aspirin (NSAID)  
and Caffeine  
Pain Reliever/Pain Reliever Aid

**250  
200**  
CAPLETS<sup>™</sup>  
\*CAPSULE-SHAPED TABLETS

**25%  
MORE  
FREE**

TAMPER-EVIDENT BOTTLE  
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH  
"SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

### Drug Facts

**Active ingredients (in each caplet)**

Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)*	Pain reliever
Caffeine 65 mg	Pain reliever aid

\*nonsteroidal anti-inflammatory drug

### Purposes

### Use

treats migraine

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount

• with other drugs containing acetaminophen

• 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

• have 3 or more alcoholic drinks every day while using this product

• take more or for a longer time than directed

**Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

**Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

### Ask a doctor before use if

- you have never had migraines diagnosed by a health professional

### Drug Facts (continued)

- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fevers and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

### Ask a doctor or pharmacist before use if you are

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.

• you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools

• have stomach pain that does not get better

• your migraine is not relieved or worsens after first dose

• new or unexpected symptoms occur

- ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

**do not use more than directed**

- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor

• do not take more than 2 caplets in 24 hours, unless directed by a doctor

• under 18 years of age: ask a doctor

### Other information

• store at room temperature 20°-25°C (68°-77°F)

• close cap tightly after use

• read all product information before using. Keep this box for important information.

**Inactive ingredients** benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

**Questions?** 1-800-452-0051

46170647



46170647



DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

LOT/EXP AREA



©2017 GSK group of companies or its licensor. All rights reserved. 16/274-300 Visit us at www.excedrin.com Distributed by: GSK Consumer Healthcare, Warren, NJ 07059

46170261



46170261



TAMPER-EVIDENT BOTTLE  
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH  
"SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)*	Pain reliever
Caffeine 65 mg	Pain reliever aid

\*nonsteroidal anti-inflammatory drug

**Use** • treats migraine

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling • asthma (wheezing) • shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

**Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck

**Drug Facts (continued)**

- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint • vomit blood • have bloody or black stools
  - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not use more than directed
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

**Other information**

- store at controlled room temperature 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

**Inactive ingredients**

benzoic acid, camauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

**Questions?** 1-800-468-7746

Glue

CODE 128 FPO



46170261

CODE 128 FPO



46170261

See new warnings information

# EXCEDRIN®

## MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

300 CAPLETS

NOI 0067-2039-33

See new warnings information

# EXCEDRIN®

## MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

300 CAPLETS

NOI 0067-2039-33

See new warnings information

# EXCEDRIN®

## MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

300 CAPLETS

NOI 0067-2039-33

See new warnings information

# EXCEDRIN®

## MIGRAINE


Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

300 CAPLETS

NOI 0067-2039-33

DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	



DRUG FACT FORMATTING: MODIFIED		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	7.5 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	7 pt.	Helvetica Cond Bold Obl
HEADINGS	7 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	5.8 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	

LOT/EXP AREA

3 0067-2035-80 6



UPC FPO

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
 ©2017 GSK group of companies or its licensor. All rights reserved.  
 Visit us at www.excedrin.com 167287-80



TAMPER-EVIDENT BOTTLE  
 DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH  
 "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

**Drug Facts**

**Active ingredients (in each GelTab)** **Purposes**  
 Acetaminophen 250 mg.....Pain reliever  
 Aspirin 250 mg (NSAID)\*.....Pain reliever  
 Caffeine 65 mg.....Pain reliever aid  
 \*nonsteroidal anti-inflammatory drug

**Use** • treats migraine

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:  
 • hives • facial swelling • asthma (wheezing) • shock  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
 • more than 2 gels in 24 hours, which is the maximum daily amount  
 • with other drugs containing acetaminophen  
 • 3 or more alcoholic drinks every day while using this product  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
 • are age 60 or older  
 • have had stomach ulcers or bleeding problems  
 • take a blood thinning (anticoagulant) or steroid drug  
 • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 • have 3 or more alcoholic drinks every day while using this product  
 • take more or for a longer time than directed  
**Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month.  
**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking

**Drug Facts (continued)**

this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

**Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

**Ask a doctor before use if**

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:  
 • feel faint • vomit blood • have bloody or black stools  
 • have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

**Drug Facts (continued)**

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.**  
 In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not use more than directed
- adults: take 2 gels with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 gels in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor


**Other information**

- store at controlled room temperature 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

**Inactive ingredients**

benzoic acid, D&C yellow #10 aluminum lake, edetate disodium, FD&C blue #1, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hypromellose, iron oxide black, isopropyl alcohol, lactose monohydrate, light mineral oil, low-substituted hydroxypropyl cellulose, maltitol syrup, medium-chain triglycerides, microcrystalline cellulose, pepsin, polysorbate 20, povidone, propyl gallate, propylene glycol, simethicone, sorbitan monolaurate, stearic acid, titanium dioxide

**Questions?** 1-800-452-0051



CODE 128 FPO

46170725



CODE 128 FPO

46170725

DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold
SUBHEADS	6 pt.	Helvetica Cond
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	83%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	